


2011/11/10

Nicotine Transdermal System
PATCH STOP SMOKING AID
21 mg delivered over 24 hours



For your family's protection, patches are supplied in child-resistant pouches. Do not use if individual pouch is open or torn. Carton shrink wrapped for your protection. Do not use if shrink wrap is open or torn.

NDC 0067-0215-07



Nicotine Transdermal System


21 mg delivered over 24 hours

PATCH STOP SMOKING AID

21mg

with
Behavior
Support
Program


DISPOSABLE Compact disc, self-help guide, and disposal tray included.




IF YOU SMOKE **MORE** THAN 10 CIGARETTES PER DAY: START WITH STEP 1
IF YOU SMOKE 10 OR **LESS** CIGARETTES PER DAY: START WITH STEP 2

7

PATCHES



**Nicotine Transdermal System
PATCH** 21 mg delivered over 24 hours
STOP SMOKING AID



**OPEN
THIS SIDE**

The use of this product is covered by
U.S. Patent No. 4,597,961 and
5,834,011
Distributed by:
Novartis Consumer Health, Inc.
Parisippany, NJ 07054-0622
©2002
301311D

TO INCREASE YOUR SUCCESS IN QUITTING:

1. You must be motivated to quit.
2. Use one patch daily according to directions.
3. Complete the full treatment program.
4. Do not use patch for more than 8 weeks.

**Nicotine Transdermal System
PATCH** 21 mg delivered over 24 hours
STOP SMOKING AID



Drug Facts

Active Ingredient (in each patch)
Nicotine, 21 mg delivered over 24 hours.

Purpose
Stop smoking aid

Use reduces withdrawal symptoms, including nicotine craving, associated with quitting smoking

Warnings
If you are pregnant or breast-feeding, only use this medicine on the advice of your health care provider. Smoking can seriously harm your child. Try to stop smoking without using any nicotine replacement medicine. This medicine is believed to be safer than smoking. However, the risks to your child from this medicine are not fully known.

Do not use
• If you continue to smoke, chew tobacco, use snuff, use nicotine gum, or use nicotine patch or other nicotine containing products

Ask a doctor before use if you have
• heart disease, recent heart attack, or irregular heartbeat. Nicotine can increase your heart rate.
• high blood pressure not controlled with medication. Nicotine can increase your blood pressure.
• an allergy to adhesive tape or have skin problems, because you are more likely to get rashes.

Ask a doctor or pharmacist before use if you are
• taking a prescription medicine for depression or asthma. Your prescription dose may need to be adjusted.

When using this product:

• do not smoke even when not wearing the patch. The nicotine in your skin will still be entering your bloodstream for several hours after you take off the patch.

• if you have vivid dreams or other sleep disturbances remove this patch at bedtime

Stop use and ask a doctor if

• skin redness caused by the patch does not go away after four days, or if your skin swells, or you get a rash
• irregular heartbeat or palpitations occur

Keep out of reach of children and pets. Used patches have enough nicotine to poison children and pets. If swallowed, get medical help or contact a Poison Control Center right away. Discard used patches using the enclosed disposal tray.

Directions

- If you are under 18 years of age, ask a doctor before use
- before using this product, read the enclosed self-help guide for complete directions and other information.
- stop smoking completely when you begin using the patch
- if you smoke more than 10 cigarettes per day, use the following schedule below:

Weeks 1 through 4	Weeks 5 through 7	Weeks 8 through 10
STEP 1 use one 21 mg patch/day	STEP 2 use one 14 mg patch/day	STEP 3 use one 7 mg patch/day

- if you smoke 10 or less cigarettes per day, start with **Step 2** for 6 weeks, then **Step 3** for 2 weeks and then stop
- apply one new patch every 24 hours on skin that is dry, clean and hairless
- remove backing from patch and immediately press onto skin. Hold for 10 seconds.
- wash hands after applying or removing patch. Throw away the patch in the enclosed disposal tray.
- the used patch should be removed and a new one applied to a different skin site at the same time each day
- if you have vivid dreams, you may remove the patch at bedtime and apply a new one in the morning
- do not wear more than one patch at a time
- do not cut patch in half or into smaller pieces
- do not leave patch on for more than 24 hours because it may irritate your skin and lose strength after 24 hours
- stop using the patch at the end of 8 weeks. If you still feel the need to use the patch talk to your doctor.

Other Information - store at 20-25°C (68-77°F)


Inactive ingredients acrylic adhesive, aluminumized polyester, cotton viscose, methacrylic acid copolymer.

Comments or Questions? Call 1-800-585-8682 Weekdays (9am-8pm ET)

See bottom panel for lot number and expiration date.

- Not for sale to persons under 18 years of age.
- Proof of age required.
- Not for sale in vending machines or from any source where proof of age cannot be verified.

Nicotine Transdermal System
PATCH 14 mg delivered over 24 hours
STOP SMOKING AID



For your family's protection, patches are supplied in child-resistant pouches. Do not use if individual pouch is open or torn. Carton shrink wrap is for your protection. Do not use if shrink wrap is open or torn.

NDC 0067-0



Nicotine Transdermal System


PATCH

14 mg delivered over 24 hours
STOP SMOKING AID

14mg


with
Behavior Support Program

disc Compact disc, self-help guide, and disposal tray included.




IF YOU SMOKE **MORE** THAN 10 CIGARETTES PER DAY: USE AFTER COMPLETING STEP 1
IF YOU SMOKE 10 OR **LESS** CIGARETTES PER DAY: START WITH STEP 2

7 PATCHES



**Nicotine Transdermal System
PATCH STOP SMOKING AID**



**OPEN
THIS SIDE**


The use of this product is covered by
U.S. Patent No. 4,597,961 and
5,834,011
Distributed by:
Novartis Consumer Health, Inc.
Parsippany, NJ 07054-0622
©2002

130D

TO INCREASE YOUR SUCCESS IN QUITTING:

1. You must be motivated to quit.
2. Use one patch daily according to directions.
3. Complete the full treatment program.
4. Do not use patch for more than 8 weeks.
5. Use patch with a behavioral support program, such as the one described in the enclosed booklet and on the compact disc.

**Nicotine Transdermal System
PATCH STOP SMOKING AID**



Drug Facts

Active Ingredient (in each patch)
Nicotine, 14 mg delivered over 24 hours.

Use reduces withdrawal symptoms, including nicotine craving, associated with quitting smoking

Warnings
If you are pregnant or breast-feeding, only use this medicine on the advice of your health care provider. Smoking can seriously harm your child. Try to stop smoking without using any nicotine replacement medicine. This medicine is believed to be safer than smoking. However, the risks to your child from this medicine are not fully known.
Do not use

- if you continue to smoke, chew tobacco, use snuff, use nicotine gum, or use nicotine patch or other nicotine containing products

Ask a doctor before use if you have

- heart disease, recent heart attack, or irregular heartbeat. Nicotine can increase your heart rate.
- high blood pressure not controlled with medication. Nicotine can increase your blood pressure.
- an allergy to adhesive tape or have skin problems, because you are more likely to get rashes.

Ask a doctor or pharmacist before use if you are

- taking a prescription medicine for depression or asthma. Your prescription dose may need to be adjusted.

When using this product

- do not smoke even when not wearing the patch. The nicotine in your skin will still be entering your bloodstream for several hours after you take off the patch.
- if you have vivid dreams or other sleep disturbances remove this patch at bedtime

Stop use and ask a doctor if

- skin redness caused by the patch does not go away after four days, or if your skin swells, or you get a rash
- irregular heartbeat or palpitations occur
- you get symptoms of nicotine overdose, such as nausea, vomiting, dizziness, weakness and rapid heartbeat

Keep out of reach of children and pets. Used patches have enough nicotine to poison children and pets. If swallowed, get medical help or contact a Poison Control Center right away. Discard used patches using the enclosed disposal tray.

Directions

- If you are under 18 years of age, ask a doctor before use
- before using this product, read the enclosed self-help guide for complete directions and other information.
- stop smoking completely when you begin using the patch
- if you smoke more than 10 cigarettes per day, use the following schedule below:

Weeks	Days	Use
1-2	14	STEP 1 use one 21 mg patch/day
3-4	14	STEP 2 use one 14 mg patch/day
5-8	7	STEP 3 use one 7 mg patch/day

- if you smoke 10 or less cigarettes per day, start with Step 2 for 6 weeks, then Step 3 for 2 weeks and then stop
- apply one new patch every 24 hours on skin that is dry, clean and hairless
- remove backing from patch and immediately press onto skin. Hold for 10 seconds.
- wash hands after applying or removing patch. Throw away the patch in the enclosed disposal tray.
- the used patch should be removed and a new one applied to a different skin site at the same time each day
- if you have vivid dreams, you may remove the patch at bedtime and apply a new one in the morning
- do not wear more than one patch at a time
- do not cut patch in half or into smaller pieces
- do not leave patch on for more than 24 hours because it may irritate your skin and loses strength after 24 hours
- stop using the patch at the end of 8 weeks. If you still feel the need to use the patch talk to your doctor.

Other Information • store at 20-25°C (68-77°F)


Inactive ingredients acrylate adhesive, aluminumized polyester, cotton viscose, methacrylic acid copolymer.

Comments or Questions? Call 1-800-585-8682 Weekdays (9am-9pm ET)

See bottom panel for lot number and expiration date.

- Not for sale to persons under 18 years of age.
- Proof of age required.
- Not for sale in vending machines or from any source where proof of age cannot be verified.

Nicotine Transdermal System
7 mg delivered over 24 hours
PATCH STOP SMOKING AID



For your family's protection, patches are supplied in child-resistant pouches. Do not use if individual pouch is open or torn. Carton shrink wrapped for your protection. Do not use if shrink wrap is open or torn.

NDC 0067-0213-

HABITROL TAKE CONTROL



Nicotine Transdermal System


PATCH STOP SMOKING AID

7 mg delivered over 24 hours

7mg

with
Behavior Support Program


DISC Compact disc, self-help guide, and disposal tray included.



FOR USE AFTER COMPLETING STEP 2


7
PATCHES

9/15/02



3 0067-5036-077

Nicotine Transdermal System
PATCH 3 7 mg delivered over 24 hours
STOP SMOKING AID



OPEN
THIS SIDE

The use of this product is covered by
U.S. Patent No. 4,597,961 and
5,834,011
Distributed by:
Novartis Consumer Health, Inc.
Parlisperry, NJ 07054-0622
©2002
30129D

- TO INCREASE YOUR SUCCESS IN QUITTING:**
1. You must be motivated to quit.
 2. Use one patch daily according to directions.
 3. Complete the full treatment program.
 4. Do not use patch for more than 8 weeks.
 5. Use patch with a behavioral support program.

Nicotine Transdermal System
PATCH 3 7 mg delivered over 24 hours
STOP SMOKING AID



Drug Facts

Active Ingredient (in each patch)
Nicotine, 7 mg delivered over 24 hours.

Use reduces withdrawal symptoms, including nicotine craving, associated with quitting smoking

Warnings

If you are pregnant or breast-feeding, only use this medicine on the advice of your health care provider. Smoking can seriously harm your child. Try to stop smoking without using any nicotine replacement medicine. This medicine is believed to be safer than smoking. However, the risks to your child from this medicine are not fully known.

Do not use

- if you continue to smoke, chew tobacco, use snuff, use nicotine gum, or use nicotine patch or other nicotine containing products
- Ask a doctor before use if you have
 - heart disease, recent heart attack, or irregular heartbeat. Nicotine can increase your heart rate.
 - high blood pressure not controlled with medication. Nicotine can increase your blood pressure.
 - an allergy to adhesive tape or have skin problems, because you are more likely to get rashes.
- Ask a doctor or pharmacist before use if you are
 - using a non-nicotine stop smoking drug
 - taking a prescription medicine for depression or asthma. Your prescription dose may need to be adjusted.

When using this product

- do not smoke even when not wearing the patch. The nicotine in your skin will still be entering your bloodstream for several hours after you take off the patch.
- if you have vivid dreams or other sleep disturbances remove this patch at bedtime

Stop use and ask a doctor if

- skin redness caused by the patch does not go away after four days, or if your skin swells, or you get a rash
 - irregular heartbeat or palpitations occur
 - you get symptoms of nicotine overdose, such as nausea, vomiting, dizziness, weakness and rapid heartbeat.
- Keep out of reach of children and pets. Used patches have enough nicotine to poison children and pets. If swallowed, get medical help or contact a Poison Control Center right away. Discard used patches using the enclosed disposal tray.

Directions

- If you are under 18 years of age, ask a doctor before use
- before using this product, read the enclosed self-help guide for complete directions and other information.
- stop smoking completely when you begin using the patch
- if you smoke more than 10 cigarettes per day, use the following schedule below:

Week 1 and 2	Week 3 and 4	Week 5 and 6
STEP 1 use one 21 mg patch/day	STEP 2 use one 14 mg patch/day	STEP 3 use one 7 mg patch/day

- if you smoke 10 or less cigarettes per day, start with Step 2 for 6 weeks, then Step 3 for 2 weeks and then stop
- apply one new patch every 24 hours on skin that is dry, clean and hairless
- remove backing from patch and immediately press onto skin. Hold for 10 seconds.
- wash hands after applying or removing patch. Throw away the patch in the enclosed disposal tray.
- the used patch should be removed and a new one applied to a different skin site at the same time each day
- if you have vivid dreams, you may remove the patch at bedtime and apply a new one in the morning
- do not wear more than one patch at a time
- do not cut patch in half or into smaller pieces
- do not leave patch on for more than 24 hours because it may irritate your skin and loses strength after 24 hours
- stop using the patch at the end of 8 weeks. If you still feel the need to use the patch talk to your doctor.

Other Information - store at 20-25°C (68-77°F)

Inactive Ingredients acrylate adhesive, aluminumized polyester, cotton viscose, methacrylic acid copolymer.

Comments or Questions? Call 1-800-585-8692 Weekdays (9am-9pm ET)

See bottom panel for lot number and expiration date.

- Not for sale to persons under 18 years of age.
- Proof of age required.
- Not for sale in vending machines or from any source where proof of age cannot be verified.

Labeling: _____
NDA No. 20-076 Rev. 1-30-11
Reviewed by *[Signature]* 4/6/02

Nicotine Transdermal System
PATCH 21 mg delivered over 24 hours
STOP SMOKING AID
21mg

DO NOT OPEN POUCH UNTIL READY TO USE.

DIRECTIONS: Apply one patch to a dry, clean, hairless portion of upper body or arms. Refer to Self-Help Guide for detailed directions.

DO NOT CUT PATCH. KEEP OUT OF REACH OF CHILDREN AND PETS. Used patches have enough nicotine to poison children and pets. If swallowed, get medical help or contact a Poison Control Center right away. Throw away used patches using the enclosed disposal tray.

Store at 20-25°C (68-77°F).

Distributed by: **Novartis Consumer Health, Inc.**, Parsippany, NJ 07054-0622

DO NOT USE IF INDIVIDUAL POUCH IS OPEN OR TORN, OR IF PATCH IS CUT.

EXP _____ LOT _____

28016C



TO OPEN CUT ALONG DOTTED LINE

[Handwritten: 6/11/02]


Labeling: _____
NDA No. 20-076 Rcd. 1-30-02
Reviewed by Haltz 6/6/02

Nicotine Transdermal System
7 mg delivered over 24 hours
PATCH STOP SMOKING AID
7 mg

DO NOT OPEN POUCH UNTIL READY TO USE.
DIRECTIONS: Apply one patch to a dry, clean, hairless portion of upper body or arms. Refer to Self-Help Guide for detailed directions.
DO NOT CUT PATCH. KEEP OUT OF REACH OF CHILDREN AND PETS. Used patches have enough nicotine to poison children and pets. If swallowed, get medical help or contact a Poison Control Center right away.
Throw away used patches using the enclosed disposal tray.
Store at 20-25°C (68-77°F).
Distributed by: Novartis Consumer Health, Inc., Parsippany, NJ 07054-0622
DO NOT USE IF INDIVIDUAL POUCH IS OPEN OR TORN, OR IF PATCH IS CUT.

EXP LOT

28014C



TO OPEN CUT ALONG DOTTED LINE

b
1/11/02

**This is a representation of an electronic record that was signed electronically and
this page is the manifestation of the electronic signature.**

/s/

Charles Ganley
7/15/02 05:50:16 PM