



00000000

Drug Facts

Active ingredient (in each patch) **Purpose**
Nicotine, 7 mg delivered over 24 hours..... Stop smoking aid

Use reduces withdrawal symptoms, including nicotine craving, associated with quitting smoking.

Warnings
If you are pregnant or breast-feeding, only use this medicine on the advice of your health care provider. Smoking can seriously harm your child. Try to stop smoking without using any nicotine replacement medicine. This medicine is believed to be safer than smoking. However, the risks to your child from this medicine are not fully known.

Ask a doctor before use if you have

- heart disease, recent heart attack, or irregular heartbeat. Nicotine can increase your heart rate.
- high blood pressure not controlled with medication. Nicotine can increase your blood pressure.
- an allergy to adhesive tape or have skin problems, because you are more likely to get rashes.
- stomach ulcer or diabetes.
- history of seizures.

Ask a doctor or pharmacist before use if you are

- using a non-nicotine stop smoking drug
- taking a prescription medicine for depression or asthma. Your prescription dose may need to be adjusted.

When using this product

- if you have vivid dreams or other sleep disturbances remove this patch at bedtime

Stop use and ask a doctor if

- skin redness caused by the patch does not go away after four days, or if your skin swells, or you get a rash
- irregular heartbeat or palpitations occur
- you get symptoms of nicotine overdose, such as nausea, vomiting, dizziness, weakness and rapid heartbeat
- you have symptoms of an allergic reaction (such as difficulty breathing or rash)

Keep out of reach of children and pets. Used patches have enough nicotine to poison children and pets. If swallowed, get medical help or contact a Poison Control Center right away. Save pouch to use for patch disposal. Dispose of the used patches by folding sticky ends together and putting in pouch.

Directions

- if you are under 18 years of age, ask a doctor before use
- before using this product, read the enclosed self-help guide for complete directions and other information
- begin using the patch on your quit day
- if you smoke more than 10 cigarettes per day, use the following schedule below:

Weeks 1 thru 4	Weeks 5 and 6	Weeks 7 and 8
STEP 1 use one 21 mg patch/day	STEP 2 use one 14 mg patch/day	STEP 3 use one 7 mg patch/day

- if you smoke 10 or less cigarettes per day, start with Step 2 for 6 weeks, then Step 3 for 2 weeks
- apply one new patch every 24 hours on skin that is dry, clean and hairless
- remove backing from patch and immediately press onto skin. Hold for 10 seconds.
- wash hands after applying or removing patch. Save pouch to use for patch disposal. Dispose of the used patches by folding sticky ends together and putting in pouch.
- the used patch should be removed and a new one applied to a different skin site at the same time each day
- if you have vivid dreams, you may remove the patch at bedtime and apply a new one in the morning
- do not wear more than one patch at a time
- do not cut patch in half or into smaller pieces
- do not leave patch on for more than 24 hours because it may irritate your skin and loses strength after 24 hours
- to avoid possible burns, remove patch before undergoing any MRI (magnetic resonance imaging) procedures
- it is important to complete treatment. If you feel you need to use the patch for a longer period to keep from smoking, talk to your health care provider.

Other information • store at 20-25°C (68-77°F)

Inactive ingredients acrylate adhesive, aluminumized polyester, cellulose paper, methacrylic acid copolymer

Comments or Questions? Call 1-800-585-8682 Weekdays (9am-9pm ET) or visit us at www.habitrol.com

See bottom panel for lot number and expiration date.

NDC 43598-446-70

HABITROL® TAKE CONTROL™




Nicotine Transdermal System PATCH

7 mg delivered over 24 hours
STOP SMOKING AID

7 mg



Includes:
Behavior Support Program with self-help guide

FOR USE AFTER COMPLETING STEP 2

7 PATCHES

Distributed by:
Dr. Reddy's Laboratories, Inc.
Princeton, NJ 08540



0 000000 00000 0



Must remain black for Lot/Exp

• Not for sale to persons under 18 years of age.
 • Proof of age required.
 • Not for sale in vending machines or from any source where proof of age cannot be verified.

For your family's protection, patches are supplied in child-resistant pouches. Do not use if individual pouch is open or torn.

See self-help guide inside.

Font & Format Specifications		
Drug Facts (Title)	Font Name: Helvetica Neue LT Std 77 Bold Condensed Oblique	10 point type
Drug Facts (continued)	N/A	N/A
Headings	Font Name: Helvetica Neue LT Std 77 Bold Condensed Oblique	8 point type
Sub-headings	Font Name: Helvetica Neue LT Std 77 Bold Condensed	8 point type
Body text	Font Name: Helvetica Neue LT Std 57 Condensed	8 point type
Table	Font Name: Helvetica Neue LT Std 57 Condensed	5 point type
Spacing of the hair lines from edge of box—Lx	Minimum of 2 spaces either side of Drug Fact Box	Yes
Tracing	0	Horizontal Scale: 100%
Lead-in margin space in body of Drug Facts	4.5 pt.	Header: 0.5 pt.
Line spacing	1.5 pt.	



00000000

Drug Facts

Active ingredient (in each patch)
Nicotine, 7 mg delivered over 24 hours.....**Purpose**
.....Stop smoking aid

Use reduces withdrawal symptoms, including nicotine craving, associated with quitting smoking.

Warnings
If you are pregnant or breast-feeding, only use this medicine on the advice of your health care provider. Smoking can seriously harm your child. Try to stop smoking without using any nicotine replacement medicine. This medicine is believed to be safer than smoking. However, the risks to your child from this medicine are not fully known.

Ask a doctor before use if you have

- heart disease, recent heart attack, or irregular heartbeat. Nicotine can increase your heart rate.
- high blood pressure not controlled with medication. Nicotine can increase your blood pressure.
- an allergy to adhesive tape or have skin problems, because you are more likely to get rashes.
- stomach ulcer or diabetes.
- history of seizures.

Ask a doctor or pharmacist before use if you are

- using a non-nicotine stop smoking drug
- taking a prescription medicine for depression or asthma. Your prescription dose may need to be adjusted.

When using this product

- if you have vivid dreams or other sleep disturbances remove this patch at bedtime

Stop use and ask a doctor if

- skin redness caused by the patch does not go away after four days, or if your skin swells, or you get a rash
- irregular heartbeat or palpitations occur
- you get symptoms of nicotine overdose, such as nausea, vomiting, dizziness, weakness and rapid heartbeat
- you have symptoms of an allergic reaction (such as difficulty breathing or rash)

Keep out of reach of children and pets. Used patches have enough nicotine to poison children and pets. If swallowed, get medical help or contact a Poison Control Center right away. Save pouch to use for patch disposal. Dispose of the used patches by folding sticky ends together and putting in pouch.

Directions

- if you are under 18 years of age, ask a doctor before use
- before using this product, read the enclosed self-help guide for complete directions and other information
- begin using the patch on your quit day
- if you smoke more than 10 cigarettes per day, use the following schedule below:

Weeks 1 thru 4	Weeks 5 and 6	Weeks 7 and 8
STEP 1 use one 21 mg patch/day	STEP 2 use one 14 mg patch/day	STEP 3 use one 7 mg patch/day

- if you smoke 10 or less cigarettes per day, start with Step 2 for 6 weeks, then Step 3 for 2 weeks
- apply one new patch every 24 hours on skin that is dry, clean and hairless
- remove backing from patch and immediately press onto skin. Hold for 10 seconds.
- wash hands after applying or removing patch. Save pouch to use for patch disposal. Dispose of the used patches by folding sticky ends together and putting in pouch.
- the used patch should be removed and a new one applied to a different skin site at the same time each day
- if you have vivid dreams, you may remove the patch at bedtime and apply a new one in the morning
- do not wear more than one patch at a time
- do not cut patch in half or into smaller pieces
- do not leave patch on for more than 24 hours because it may irritate your skin and loses strength after 24 hours
- to avoid possible burns, remove patch before undergoing any MRI (magnetic resonance imaging) procedures
- it is important to complete treatment. If you feel you need to use the patch for a longer period to keep from smoking, talk to your health care provider.

Other information • store at 20-25°C (68-77°F)

Inactive ingredients acrylate adhesive, aluminumized polyester, cellulose paper, methacrylic acid copolymer

Comments or Questions? Call 1-800-585-8682 Weekdays (9am-9pm ET) or visit us at www.habitrol.com

See bottom panel for lot number and expiration date.

Nicotine Transdermal System PATCH
7 mg delivered over 24 hours
STOP SMOKING AID

STEP 3

Nicotine Transdermal System PATCH
7 mg delivered over 24 hours
STOP SMOKING AID

STEP 3

Nicotine Transdermal System PATCH
7 mg delivered over 24 hours
STOP SMOKING AID

NDC 43598-446-74

HABITROL® TAKE CONTROL™

SUPPORT PROGRAM

TO INCREASE YOUR SUCCESS IN QUITTING:

- You must be motivated to quit.
- Use one patch daily according to directions.
- It is important to complete treatment.
- If you feel you need to use the patch for a longer period to keep from smoking, talk to your health care provider.
- Use patch with a behavioral support program, such as the one described in the enclosed booklet.

Nicotine Transdermal System PATCH
7 mg delivered over 24 hours
STOP SMOKING AID

STEP 3

Nicotine Transdermal System PATCH
7 mg delivered over 24 hours
STOP SMOKING AID

STEP 3

7 mg

Includes:
Behavior Support Program with self-help guide

14 PATCHES

FOR USE AFTER COMPLETING STEP 2

See self-help guide inside.

FOR your family's protection, patches are supplied in child-resistant pouches. Do not use if individual pouch is open or torn.

Distributed by:
Dr. Reddy's Laboratories, Inc.
Princeton, NJ 08540



00000000



Must remain black for Lot/Exp



UPC
0 00000 00000 0

Font & Format Specifications		
Drug Facts (Title)	Font Name: Helvetica Neue LT Std 77 Bold Condensed Oblique	10 point type
Drug Facts (continued)	N/A	N/A
Headings	Font Name: Helvetica Neue LT Std 77 Bold Condensed Oblique	8 point type
Sub-headings	Font Name: Helvetica Neue LT Std 77 Bold Condensed	8 point type
Body text	Font Name: Helvetica Neue LT Std 57 Condensed	8 point type
Table	Font Name: Helvetica Neue LT Std 57 Condensed	5 point type
Spacing of the hair lines from edge of box—L4. Minimum of 2 spaces either side of Drug Fact Box	Yes	
Tracing	0	Horizontal Scale: 100%
Lead-in margin space in body of Drug Facts	0.5 pt.	Hardner
Line height	1.5 pt.	



Drug Facts

Active ingredient (in each patch) Nicotine, 14 mg delivered over 24 hours. **Purpose** Stop smoking aid

Use reduces withdrawal symptoms, including nicotine craving, associated with quitting smoking.

Warnings
If you are pregnant or breast-feeding, only use this medicine on the advice of your health care provider. Smoking can seriously harm your child. Try to stop smoking without using any nicotine replacement medicine. This medicine is believed to be safer than smoking. However, the risks to your child from this medicine are not fully known.

Ask a doctor before use if you have

- heart disease, recent heart attack, or irregular heartbeat. Nicotine can increase your heart rate.
- high blood pressure not controlled with medication. Nicotine can increase your blood pressure.
- an allergy to adhesive tape or have skin problems, because you are more likely to get rashes.
- stomach ulcer or diabetes.
- history of seizures.

Ask a doctor or pharmacist before use if you are

- using a non-nicotine stop smoking drug
- taking a prescription medicine for depression or asthma. Your prescription dose may need to be adjusted.

When using this product

- if you have vivid dreams or other sleep disturbances remove this patch at bedtime

Stop use and ask a doctor if

- skin redness caused by the patch does not go away after four days, or if your skin swells, or you get a rash
- irregular heartbeat or palpitations occur
- you get symptoms of nicotine overdose, such as nausea, vomiting, dizziness, weakness and rapid heartbeat
- you have symptoms of an allergic reaction (such as difficulty breathing or rash)

Keep out of reach of children and pets. Used patches have enough nicotine to poison children and pets. If swallowed, get medical help or contact a Poison Control Center right away. Save pouch to use for patch disposal. Dispose of the used patches by folding sticky ends together and putting in pouch.

Directions

- If you are under 18 years of age, ask a doctor before use
- before using this product, read the enclosed self-help guide for complete directions and other information
- begin using the patch on your quit day
- if you smoke more than 10 cigarettes per day, use the following schedule below:

Weeks 1 thru 4	Weeks 5 and 6	Weeks 7 and 8
STEP 1 use one 21 mg patch/day	STEP 2 use one 14 mg patch/day	STEP 3 use one 7 mg patch/day

- if you smoke 10 or less cigarettes per day, start with Step 2 for 6 weeks, then Step 3 for 2 weeks
- apply one new patch every 24 hours on skin that is dry, clean and hairless
- remove backing from patch and immediately press onto skin. Hold for 10 seconds.
- wash hands after applying or removing patch. Save pouch to use for patch disposal. Dispose of the used patches by folding sticky ends together and putting in pouch.
- the used patch should be removed and a new one applied to a different skin site at the same time each day
- if you have vivid dreams, you may remove the patch at bedtime and apply a new one in the morning
- do not wear more than one patch at a time
- do not cut patch in half or into smaller pieces
- do not leave patch on for more than 24 hours because it may irritate your skin and loses strength after 24 hours
- to avoid possible burns, remove patch before undergoing any MRI (magnetic resonance imaging) procedures
- it is important to complete treatment. If you feel you need to use the patch for a longer period to keep from smoking, talk to your health care provider.

Other information • store at 20-25°C (68-77°F)

Inactive ingredients acrylate adhesive, aluminumized polyester, cellulose paper, methacrylic acid copolymer

Comments or Questions? Call 1-800-585-8682 Weekdays (9am-8pm ET) or visit us at www.habitrol.com

See bottom panel for lot number and expiration date.

• Not for sale to persons under 18 years of age.
• Proof of age required.
• Not for sale in vending machines or from any source where proof of age cannot be verified.

Nicotine Transdermal System PATCH
14 mg delivered over 24 hours
STOP SMOKING AID



For your family's protection, patches are supplied in child-resistant pouches. Do not use if individual pouch is open or torn.

See self-help guide inside.

TO INCREASE YOUR SUCCESS IN QUITTING:

1. You must be motivated to quit.
2. Use one patch daily according to directions.
3. It is important to complete treatment.
4. If you feel you need to use the patch for a longer period to keep from smoking, talk to your health care provider.
5. Use patch with a behavioral support program, such as the one described in the enclosed booklet.

NDC 43598-447-70

HABITROL® TAKE CONTROL™



Nicotine Transdermal System PATCH
14 mg delivered over 24 hours
STOP SMOKING AID



Nicotine Transdermal System PATCH 14 mg delivered over 24 hours
STOP SMOKING AID

14 mg

Includes:
Behavior Support Program with self-help guide



IF YOU SMOKE **MORE** THAN 10 CIGARETTES PER DAY: USE AFTER COMPLETING STEP 1

IF YOU SMOKE 10 OR **LESS** CIGARETTES PER DAY: START WITH STEP 2

7 PATCHES

Distributed by:
Dr. Reddy's Laboratories, Inc.
Princeton, NJ 08540



Must remain black for Lot/Exp



Font & Format Specifications		
Drug Facts (Title)	Font Name: Helvetica Neue LT Std 77 Bold Condensed Oblique	10 point type
Drug Facts (Subtitle)	N/A	N/A
Headings	Font Name: Helvetica Neue LT Std 77 Bold Condensed Oblique	8 point type
Subheadings	Font Name: Helvetica Neue LT Std 77 Bold Condensed	8 point type
Body text	Font Name: Helvetica Neue LT Std 52 Condensed	6 point type
Sub-headers	Font Name: Helvetica Neue LT Std 52 Condensed	5 point type
Spacing of the hair lines from edge of box=LA, Minimum of 2 spaces either side of Drug Fact Box		Yes
Tracking	0	Horizontal Scale: 100%
Leading (Minimum space in body copy of Drug Facts)	6.5 pt.	Headlines: 0.5 pt.
Baselines	1.5 pt.	



Drug Facts

Active ingredient (in each patch) **Purpose**
Nicotine, 14 mg delivered over 24 hours..... Stop smoking aid

Use reduces withdrawal symptoms, including nicotine craving, associated with quitting smoking.

Warnings
If you are pregnant or breast-feeding, only use this medicine on the advice of your health care provider. Smoking can seriously harm your child. Try to stop smoking without using any nicotine replacement medicine. This medicine is believed to be safer than smoking. However, the risks to your child from this medicine are not fully known.

Ask a doctor before use if you have

- heart disease, recent heart attack, or irregular heartbeat. Nicotine can increase your heart rate.
- high blood pressure not controlled with medication. Nicotine can increase your blood pressure.
- an allergy to adhesive tape or have skin problems, because you are more likely to get rashes.
- stomach ulcer or diabetes.
- history of seizures.

Ask a doctor or pharmacist before use if you are

- using a non-nicotine stop smoking drug
- taking a prescription medicine for depression or asthma. Your prescription dose may need to be adjusted.

When using this product

- if you have vivid dreams or other sleep disturbances remove this patch at bedtime

Stop use and ask a doctor if

- skin redness caused by the patch does not go away after four days, or if your skin swells, or you get a rash
- irregular heartbeat or palpitations occur
- you get symptoms of nicotine overdose, such as nausea, vomiting, dizziness, weakness and rapid heartbeat
- you have symptoms of an allergic reaction (such as difficulty breathing or rash)

Keep out of reach of children and pets. Used patches have enough nicotine to poison children and pets. If swallowed, get medical help or contact a Poison Control Center right away. Save pouch to use for patch disposal. Dispose of the used patches by folding sticky ends together and putting in pouch.

Directions

- If you are under 18 years of age, ask a doctor before use
- before using this product, read the enclosed self-help guide for complete directions and other information
- begin using the patch on your quit day
- if you smoke more than 10 cigarettes per day, use the following schedule below:

Weeks 1 thru 4	Weeks 5 and 6	Weeks 7 and 8
STEP 1 use one 21 mg patch/day	STEP 2 use one 14 mg patch/day	STEP 3 use one 7 mg patch/day

- if you smoke 10 or less cigarettes per day, start with Step 2 for 6 weeks, then Step 3 for 2 weeks
- apply one new patch every 24 hours on skin that is dry, clean and hairless
- remove backing from patch and immediately press onto skin. Hold for 10 seconds.
- wash hands after applying or removing patch. Save pouch to use for patch disposal. Dispose of the used patches by folding sticky ends together and putting in pouch.
- the used patch should be removed and a new one applied to a different skin site at the same time each day
- if you have vivid dreams, you may remove the patch at bedtime and apply a new one in the morning
- do not wear more than one patch at a time
- do not cut patch in half or into smaller pieces
- do not leave patch on for more than 24 hours because it may irritate your skin and loses strength after 24 hours
- to avoid possible burns, remove patch before undergoing any MRI (magnetic resonance imaging) procedures
- it is important to complete treatment. If you feel you need to use the patch for a longer period to keep from smoking, talk to your health care provider.

Other information • store at 20-25°C (68-77°F)

Inactive ingredients acrylate adhesive, aluminumized polyester, cellulose paper, methacrylic acid copolymer

Comments or Questions? Call 1-800-585-8682 Weekdays (9am-8pm ET) or visit us at www.habitrol.com

See bottom panel for lot number and expiration date.



TO INCREASE YOUR SUCCESS IN QUITTING:

1. You must be motivated to quit.
2. Use one patch daily according to directions.
3. It is important to complete treatment.
4. If you feel you need to use the patch for a longer period to keep from smoking, talk to your health care provider.
5. Use patch with a behavioral support program, such as the one described in the enclosed booklet.

NDC 43598-447-74

HABITROL® TAKE CONTROL™



Nicotine Transdermal System PATCH
14 mg delivered over 24 hours
STOP SMOKING AID



Nicotine Transdermal System PATCH

14 mg delivered over 24 hours
STOP SMOKING AID

14 mg

Includes:
Behavior Support Program with self-help guide



For your family's protection, patches are supplied in child-resistant pouches.
Do not use if individual pouch is open or torn.

See self-help guide inside.

IF YOU SMOKE **MORE** THAN 10 CIGARETTES PER DAY: USE AFTER COMPLETING STEP 1

IF YOU SMOKE 10 OR **LESS** CIGARETTES PER DAY: START WITH STEP 2

14 PATCHES

Nicotine Transdermal System PATCH
14 mg delivered over 24 hours
STOP SMOKING AID



Distributed by:
Dr. Reddy's Laboratories, Inc.
Princeton, NJ 08540



Must remain black for Lot/Exp



Font & Format Specifications		
Drug Facts (Title)	Font Name: Helvetica Neue LT Std 77 Bold Condensed Oblique	10 point type
Drug Facts (Subtitle)	N/A	N/A
Headings	Font Name: Helvetica Neue LT Std 77 Bold Condensed Oblique	8 point type
Subheadings	Font Name: Helvetica Neue LT Std 77 Bold Condensed	8 point type
Body text	Font Name: Helvetica Neue LT Std 52 Condensed	6 point type
Sub-headers	Font Name: Helvetica Neue LT Std 52 Condensed	5 point type
Spacing of the hair lines from edge of box=LA, Minimum of 2 spaces either side of Drug Fact Box		Yes
Tracking	0	Horizontal Scale: 100%
Leading (Minimum space in body copy of Drug Facts)	6.5 pt.	Headlines: 0.5 pt.
Baselines	1.5 pt.	



Drug Facts

Active ingredient (in each patch)
Nicotine, 21 mg delivered over 24 hours. **Purpose**
Stop smoking aid

Use reduces withdrawal symptoms, including nicotine craving, associated with quitting smoking.

Warnings
If you are pregnant or breast-feeding, only use this medicine on the advice of your health care provider. Smoking can seriously harm your child. Try to stop smoking without using any nicotine replacement medicine. This medicine is believed to be safer than smoking. However, the risks to your child from this medicine are not fully known.

Ask a doctor before use if you have

- heart disease, recent heart attack, or irregular heartbeat. Nicotine can increase your heart rate.
- high blood pressure not controlled with medication. Nicotine can increase your blood pressure.
- an allergy to adhesive tape or have skin problems, because you are more likely to get rashes.
- stomach ulcer or diabetes.
- history of seizures.

Ask a doctor or pharmacist before use if you are

- using a non-nicotine stop smoking drug
- taking a prescription medicine for depression or asthma. Your prescription dose may need to be adjusted.

When using this product

- if you have vivid dreams or other sleep disturbances remove this patch at bedtime

Stop use and ask a doctor if

- skin redness caused by the patch does not go away after four days, or if your skin swells, or you get a rash
- irregular heartbeat or palpitations occur
- you get symptoms of nicotine overdose, such as nausea, vomiting, dizziness, weakness and rapid heartbeat
- you have symptoms of an allergic reaction (such as difficulty breathing or rash)

Keep out of reach of children and pets. Used patches have enough nicotine to poison children and pets. If swallowed, get medical help or contact a Poison Control Center right away. Save pouch to use for patch disposal. Dispose of the used patches by folding sticky ends together and putting in pouch.

Directions

- if you are under 18 years of age, ask a doctor before use
- before using this product, read the enclosed self-help guide for complete directions and other information
- begin using the patch on your quit day
- if you smoke more than 10 cigarettes per day, use the following schedule below:

Weeks 1 thru 4	Weeks 5 and 6	Weeks 7 and 8
STEP 1 use one 21 mg patch/day	STEP 2 use one 14 mg patch/day	STEP 3 use one 7 mg patch/day

- if you smoke 10 or less cigarettes per day, start with Step 2 for 6 weeks, then Step 3 for 2 weeks
- apply one new patch every 24 hours on skin that is dry, clean and hairless
- remove backing from patch and immediately press onto skin. Hold for 10 seconds.
- wash hands after applying or removing patch. Save pouch to use for patch disposal. Dispose of the used patches by folding sticky ends together and putting in pouch.
- the used patch should be removed and a new one applied to a different skin site at the same time each day
- if you have vivid dreams, you may remove the patch at bedtime and apply a new one in the morning
- do not wear more than one patch at a time
- do not cut patch in half or into smaller pieces
- do not leave patch on for more than 24 hours because it may irritate your skin and loses strength after 24 hours
- to avoid possible burns, remove patch before undergoing any MRI (magnetic resonance imaging) procedures
- it is important to complete treatment. If you feel you need to use the patch for a longer period to keep from smoking, talk to your health care provider.

Other information • store at 20-25°C (68-77°F)

Inactive ingredients acrylate adhesive, aluminumized polyester, cellulose paper, methacrylic acid copolymer

Comments or Questions? Call 1-800-585-8682 Weekdays (9am-8pm ET) or visit us at www.habitrol.com

Nicotine Transdermal System
PATCH
21 mg delivered over 24 hours
STOP SMOKING AID



For your family's protection, patches are supplied in child-resistant pouches.
Do not use if individual pouch is open or torn.

See self-help guide inside.

NDC 43598-448-70




Nicotine Transdermal System
PATCH
21 mg delivered over 24 hours
STOP SMOKING AID



Nicotine Transdermal System **PATCH** 21 mg delivered over 24 hours STOP SMOKING AID

21 mg



Includes:
Behavior Support Program with self-help guide

IF YOU SMOKE **MORE** THAN 10 CIGARETTES PER DAY: START WITH STEP 1

IF YOU SMOKE 10 OR **LESS** CIGARETTES PER DAY: START WITH STEP 2

7 PATCHES

Distributed by:
Dr. Reddy's Laboratories, Inc.
Princeton, NJ 08540



Must remain black for Lot/Exp

Font & Format Specifications		
Drug Facts (Title)	Font Name: Helvetica Neue LT Std 77 B0 d Condensed Oblique	10 point type
Drug Facts (continued)	N/A	N/A
Headings	Font Name: Helvetica Neue LT Std 77 Bold Condensed Oblique	8 point type
Subheadings	Font Name: Helvetica Neue LT Std 77 Bold Condensed	6 point type
Body text	Font Name: Helvetica Neue LT Std 57 Condensed	6 point type
Table	Font Name: Helvetica Neue LT Std 57 Condensed	5 point type
Spacing of the hair lines from edge of box—LA, Minimum of 2 spaces either side of Drug Facts Box		Yes
Tracking	0	Horizontal Scale
Leading (Minimum space in body copy of Drug Facts)	6.5 pt.	Halfline
Baseline	1.5 pt.	0.5 pt.



Drug Facts

Active ingredient (in each patch) Nicotine, 21 mg delivered over 24 hours. **Purpose** Stop smoking aid

Use reduces withdrawal symptoms, including nicotine craving, associated with quitting smoking.

Warnings
If you are pregnant or breast-feeding, only use this medicine on the advice of your health care provider. Smoking can seriously harm your child. Try to stop smoking without using any nicotine replacement medicine. This medicine is believed to be safer than smoking. However, the risks to your child from this medicine are not fully known.

Ask a doctor before use if you have

- heart disease, recent heart attack, or irregular heartbeat. Nicotine can increase your heart rate.
- high blood pressure not controlled with medication. Nicotine can increase your blood pressure.
- an allergy to adhesive tape or have skin problems, because you are more likely to get rashes.
- stomach ulcer or diabetes.
- history of seizures.

Ask a doctor or pharmacist before use if you are

- using a non-nicotine stop smoking drug
- taking a prescription medicine for depression or asthma. Your prescription dose may need to be adjusted.

When using this product

- if you have vivid dreams or other sleep disturbances remove this patch at bedtime

Stop use and ask a doctor if

- skin redness caused by the patch does not go away after four days, or if your skin swells, or you get a rash
- irregular heartbeat or palpitations occur
- you get symptoms of nicotine overdose, such as nausea, vomiting, dizziness, weakness and rapid heartbeat
- you have symptoms of an allergic reaction (such as difficulty breathing or rash)

Keep out of reach of children and pets. Used patches have enough nicotine to poison children and pets. If swallowed, get medical help or contact a Poison Control Center right away. Save pouch to use for patch disposal. Dispose of the used patches by folding sticky ends together and putting in pouch.

Directions

- if you are under 18 years of age, ask a doctor before use
- before using this product, read the enclosed self-help guide for complete directions and other information
- begin using the patch on your quit day
- if you smoke more than 10 cigarettes per day, use the following schedule below:

Weeks 1 thru 4	Weeks 5 and 6	Weeks 7 and 8
STEP 1 use one 21 mg patch/day	STEP 2 use one 14 mg patch/day	STEP 3 use one 7 mg patch/day

- if you smoke 10 or less cigarettes per day, start with Step 2 for 6 weeks, then Step 3 for 2 weeks
- apply one new patch every 24 hours on skin that is dry, clean and hairless
- remove backing from patch and immediately press onto skin. Hold for 10 seconds.
- wash hands after applying or removing patch. Save pouch to use for patch disposal. Dispose of the used patches by folding sticky ends together and putting in pouch.
- the used patch should be removed and a new one applied to a different skin site at the same time each day
- if you have vivid dreams, you may remove the patch at bedtime and apply a new one in the morning
- do not wear more than one patch at a time
- do not cut patch in half or into smaller pieces
- do not leave patch on for more than 24 hours because it may irritate your skin and loses strength after 24 hours
- to avoid possible burns, remove patch before undergoing any MRI (magnetic resonance imaging) procedures
- it is important to complete treatment. If you feel you need to use the patch for a longer period to keep from smoking, talk to your health care provider.

Other information • store at 20-25°C (68-77°F)

Inactive ingredients acrylate adhesive, aluminumized polyester, cellulose paper, methacrylic acid copolymer

Comments or Questions? Call 1-800-585-8682 Weekdays (9am-8pm ET) or visit us at www.habitrol.com

Nicotine Transdermal System PATCH 21 mg delivered over 24 hours STOP SMOKING AID



For your family's protection, patches are supplied in child-resistant pouches. Do not use if individual pouch is open or torn.

See self-help guide inside.



TO INCREASE YOUR SUCCESS IN QUITTING:

1. You must be motivated to quit.
2. Use one patch daily according to directions.
3. It is important to complete treatment.
4. If you feel you need to use the patch for a longer period to keep from smoking, talk to your health care provider.
5. Use patch with a behavioral support program, such as the one described in the enclosed booklet.

NDC 43598-448-74




Nicotine Transdermal System PATCH 21 mg delivered over 24 hours STOP SMOKING AID



Nicotine Transdermal System PATCH 21 mg delivered over 24 hours STOP SMOKING AID

21 mg



Includes:
Behavior Support Program with self-help guide

IF YOU SMOKE **MORE** THAN 10 CIGARETTES PER DAY: START WITH STEP 1

IF YOU SMOKE 10 OR **LESS** CIGARETTES PER DAY: START WITH STEP 2

14 PATCHES

Distributed by:
Dr. Reddy's Laboratories, Inc.
Princeton, NJ 08540



Must remain black for Lot/Exp

Font & Format Specifications		
Drug Facts (Title)	Font Name: Helvetica Neue LT Std 77 B0 d Condensed Oblique	10 point type
Drug Facts (continued)	N/A	N/A
Headings	Font Name: Helvetica Neue LT Std 77 Bold Condensed Oblique	8 point type
Subheadings	Font Name: Helvetica Neue LT Std 77 Bold Condensed	6 point type
Body text	Font Name: Helvetica Neue LT Std 57 Condensed	6 point type
Table	Font Name: Helvetica Neue LT Std 57 Condensed	5 point type
Spacing of the hair lines from edge of box—LA, Minimum of 2 spaces either side of Drug Facts Box		Yes
Tracking	0	Horizontal Scale: 100%
Leading (Minimum space in body copy of Drug Facts)	6.5 pt.	Halfline
Baseline	1.5 pt.	



Drug Facts

Active ingredient (in each patch)
Nicotine, 21 mg delivered over 24 hours.....**Purpose**
Stop smoking aid

Use reduces withdrawal symptoms, including nicotine craving, associated with quitting smoking.

Warnings
If you are pregnant or breast-feeding, only use this medicine on the advice of your health care provider. Smoking can seriously harm your child. Try to stop smoking without using any nicotine replacement medicine. This medicine is believed to be safer than smoking. However, the risks to your child from this medicine are not fully known.

Ask a doctor before use if you have

- heart disease, recent heart attack, or irregular heartbeat. Nicotine can increase your heart rate.
- high blood pressure not controlled with medication. Nicotine can increase your blood pressure.
- an allergy to adhesive tape or have skin problems, because you are more likely to get rashes.
- stomach ulcer or diabetes.
- history of seizures.

Ask a doctor or pharmacist before use if you are

- using a non-nicotine stop smoking drug
- taking a prescription medicine for depression or asthma. Your prescription dose may need to be adjusted.

When using this product

- if you have vivid dreams or other sleep disturbances remove this patch at bedtime

Stop use and ask a doctor if

- skin redness caused by the patch does not go away after four days, or if your skin swells, or you get a rash
- irregular heartbeat or palpitations occur
- you get symptoms of nicotine overdose, such as nausea, vomiting, dizziness, weakness and rapid heartbeat
- you have symptoms of an allergic reaction (such as difficulty breathing or rash)

Keep out of reach of children and pets. Used patches have enough nicotine to poison children and pets. If swallowed, get medical help or contact a Poison Control Center right away. Save pouch to use for patch disposal. Dispose of the used patches by folding sticky ends together and putting in pouch.

Directions

- if you are under 18 years of age, ask a doctor before use
- before using this product, read the enclosed self-help guide for complete directions and other information
- begin using the patch on your quit day
- if you smoke more than 10 cigarettes per day, use the following schedule below:

Weeks 1 thru 4	Weeks 5 and 6	Weeks 7 and 8
STEP 1 use one 21 mg patch/day	STEP 2 use one 14 mg patch/day	STEP 3 use one 7 mg patch/day

- if you smoke 10 or less cigarettes per day, start with Step 2 for 6 weeks, then Step 3 for 2 weeks
- apply one new patch every 24 hours on skin that is dry, clean and hairless
- remove backing from patch and immediately press onto skin. Hold for 10 seconds.
- wash hands after applying or removing patch. Save pouch to use for patch disposal. Dispose of the used patches by folding sticky ends together and putting in pouch.
- the used patch should be removed and a new one applied to a different skin site at the same time each day
- if you have vivid dreams, you may remove the patch at bedtime and apply a new one in the morning
- do not wear more than one patch at a time
- do not cut patch in half or into smaller pieces
- do not leave patch on for more than 24 hours because it may irritate your skin and loses strength after 24 hours
- to avoid possible burns, remove patch before undergoing any MRI (magnetic resonance imaging) procedures
- it is important to complete treatment. If you feel you need to use the patch for a longer period to keep from smoking, talk to your health care provider.

Other information • store at 20-25°C (68-77°F)

Inactive ingredients acrylate adhesive, aluminumized polyester, cellulose paper, methacrylic acid copolymer

Comments or Questions? Call 1-800-585-8682 Weekdays (9am-8pm ET) or visit us at www.habitrol.com

See bottom panel for lot number and expiration date.

• Not for sale to persons under 18 years of age.
• Proof of age required.
• Not for sale in vending machines or from any source where proof of age cannot be verified.



TO INCREASE YOUR SUCCESS IN QUITTING:

1. You must be motivated to quit.
2. Use one patch daily according to directions.
3. It is important to complete treatment.
4. If you feel you need to use the patch for a longer period to keep from smoking, talk to your health care provider.
5. Use patch with a behavioral support program, such as the one described in the enclosed booklet.

NDC 43598-448-28



Nicotine Transdermal System PATCH
21 mg delivered over 24 hours
STOP SMOKING AID



Nicotine Transdermal System PATCH
21 mg delivered over 24 hours
STOP SMOKING AID

Nicotine Transdermal System PATCH
21 mg delivered over 24 hours
STOP SMOKING AID



For your family's protection, patches are supplied in child-resistant pouches.
Do not use if individual pouch is open or torn.

21 mg



Includes:
Behavior Support Program with self-help guide

IF YOU SMOKE **MORE** THAN 10 CIGARETTES PER DAY: START WITH STEP 1
IF YOU SMOKE 10 OR **LESS** CIGARETTES PER DAY: START WITH STEP 2

28 PATCHES


Distributed by:
Dr. Reddy's Laboratories, Inc.
Princeton, NJ 08540

See self-help guide inside.



Must remain black for Lot/Exp

Font & Format Specifications		
Drug Facts (Title)	Font Name: Helvetica Neue LT Std 77 Bb d Condensed Oblique	10 point type
Drug Facts (continued)	N/A	N/A
Heading	Font Name: Helvetica Neue LT Std 77 Bold Condensed Oblique	8 point type
Subheading	Font Name: Helvetica Neue LT Std 77 Bold Condensed	8 point type
Body text	Font Name: Helvetica Neue LT Std 57 Condensed	8 point type
Table	Font Name: Helvetica Neue LT Std 57 Condensed	5 point type
Spacing of the hair lines from edge of bar—La. Minimum of 2 spaces either side of Drug Fact Bar	Yes	
Tracking	0	Horizontal Scale: 100%
Leading (Minimum space in body copy of Drug Facts)	6.5 pt.	Halfline 0.5 pt.
Baseline	1.5 pt.	



Drug Facts

Active ingredient (in each patch)

Nicotine, 21 mg delivered over 24 hours.....	Stop smoking aid
Nicotine, 14 mg delivered over 24 hours.....	Stop smoking aid
Nicotine, 7 mg delivered over 24 hours.....	Stop smoking aid

Purpose

Stop smoking aid

Use reduces withdrawal symptoms, including nicotine craving, associated with quitting smoking.

Warnings

If you are pregnant or breast-feeding, only use this medicine on the advice of your health care provider. Smoking can seriously harm your child. Try to stop smoking without using any nicotine replacement medicine. This medicine is believed to be safer than smoking. However, the risks to your child from this medicine are not fully known.

Ask a doctor before use if you have

- heart disease, recent heart attack, or irregular heartbeat. Nicotine can increase your heart rate.
- high blood pressure not controlled with medication. Nicotine can increase your blood pressure.
- an allergy to adhesive tape or have skin problems, because you are more likely to get rashes.
- stomach ulcer or diabetes.
- history of seizures.

Ask a doctor or pharmacist before use if you are

- using a non-nicotine stop smoking drug
- taking a prescription medicine for depression or asthma. Your prescription dose may need to be adjusted.

When using this product

- if you have vivid dreams or other sleep disturbances remove this patch at bedtime

Stop use and ask a doctor if

- skin redness caused by the patch does not go away after four days, or if your skin swells, or you get a rash
- irregular heartbeat or palpitations occur
- you get symptoms of nicotine overdose, such as nausea, vomiting, dizziness, weakness and rapid heartbeat
- you have symptoms of an allergic reaction (such as difficulty breathing or rash)

Keep out of reach of children and pets. Used patches have enough nicotine to poison children and pets. If swallowed, get medical help or contact a Poison Control Center right away. Save pouch to use for patch disposal. Dispose of the used patches by folding sticky ends together and putting in pouch.

Drug Facts (continued)

Directions

- If you are under 18 years of age, ask a doctor before use
- before using this product, read the enclosed self-help guide for complete directions and other information
- begin using the patch on your quit day
- if you smoke more than 10 cigarettes per day, use the following schedule below:

Weeks 1 thru 4	Weeks 5 and 6	Weeks 7 and 8
STEP 1 use one 21 mg patch/day	STEP 2 use one 14 mg patch/day	STEP 3 use one 7 mg patch/day

- if you smoke 10 or less cigarettes per day, start with Step 2 for 6 weeks, then Step 3 for 2 weeks
- apply one new patch every 24 hours on skin that is dry, clean and hairless
- remove backing from patch and immediately press onto skin. Hold for 10 seconds.
- wash hands after applying or removing patch. Save pouch to use for patch disposal.
- Dispose of the used patches by folding sticky ends together and putting in pouch.
- the used patch should be removed and a new one applied to a different skin site at the same time each day
- if you have vivid dreams, you may remove the patch at bedtime and apply a new one in the morning
- do not wear more than one patch at a time
- do not cut patch in half or into smaller pieces
- do not leave patch on for more than 24 hours because it may irritate your skin and loses strength after 24 hours
- to avoid possible burns, remove patch before undergoing any MRI (magnetic resonance imaging) procedures
- it is important to complete treatment. If you feel you need to use the patch for a longer period to keep from smoking, talk to your health care provider.

Other information • store at 20-25°C (68-77°F)

Inactive ingredients acrylate adhesive, aluminumized polyester, cellulose paper, methacrylic acid copolymer

Comments or Questions? Call 1-800-585-8682 Weekdays (8am-8pm ET) or visit us at www.habitrol.com

TO INCREASE YOUR SUCCESS IN QUITTING:

- You must be motivated to quit.
- Use one patch daily according to directions.
- It is important to complete treatment.
- If you feel you need to use the patch for a longer period to keep from smoking, talk to your health care provider.
- Use patch with a behavioral support program, such as the one described in the enclosed booklet.

NDC 43598-445-56

HABITROL™ TAKE CONTROL™

Nicotine Transdermal System PATCH KIT STOP SMOKING AID

Contains 21 mg, 14 mg and 7 mg delivered over 24 hours

INCLUDES ALL 3 STEPS

STEP 1
21 mg delivered over 24 hours
Contains 28 Patches (4 week supply)

STEP 2
14 mg delivered over 24 hours
Contains 14 Patches (2 week supply)

STEP 3
7 mg delivered over 24 hours
Contains 14 Patches (2 week supply)

56 PATCHES

COMPLETE STOP SMOKING AID KIT – USE ONLY IF YOU SMOKE MORE THAN 10 CIGARETTES PER DAY

Includes: Behavior Support Program with self-help guide

Nicotine Transdermal System PATCH KIT STOP SMOKING AID

Contains 21 mg, 14 mg and 7 mg delivered over 24 hours

For your family's protection, patches are supplied in child-resistant pouches.
Do not use if individual pouch is open or torn.

See self-help guide inside.

Distributed by:
Dr. Reddy's Laboratories, Inc.
Princeton, NJ 08540

See bottom panel for lot number and expiration date.

• Not for sale to persons under 18 years of age.
• Proof of age required.
• Not for sale in vending machines or from any source where proof of age cannot be verified.

Font & Format Specifications

Drug Facts (Title)	Font Name: Helvetica Neue LT Std 77 Bold Condensed Oblique	10 point type
Drug Facts (continued)	Font Name: Helvetica Neue LT Std 77 Bold Condensed Oblique (Drug Facts Helvetica Neue LT Std 77 Condensed (continued))	8 point type
Warnings	Font Name: Helvetica Neue LT Std 77 Bold Condensed Oblique	8 point type
Directions	Font Name: Helvetica Neue LT Std 77 Bold Condensed	8 point type
Body text	Font Name: Helvetica Neue LT Std 57 Condensed	6 point type
Buttons	Font Name: Helvetica Neue LT Std 57 Condensed	5 point type
Spacing of the hair lines from edge of box – i.e., Minimum of 2 spaces either side of Drug Facts Box		Yes
Tracking	0 to 25 Horizontal Scale	0 to 90%
Leading (Minimum space in body copy of Drug Facts)	6.5 pt Hairline	0.5 pt.
Buttons	1.5 pt.	

I: Thinking About Quitting
Nicotine Transdermal System Patch Stop Smoking Aid

Why the Habitrol® Take Control™ Support Program Leads to Success

usually have to try to quit several times before they completely stop.

Your own chances to stop smoking depend on how strongly you are addicted to nicotine, how much you want to quit, and how closely you follow a quitting program like the one that comes with this product. If you find you cannot stop or if you start smoking again after using this product, please talk to a health care professional who can help you find a program that may work better for you.

First, there is the 3-step nicotine patch that delivers controlled amounts of nicotine to help reduce your withdrawal cravings for nicotine. The patch utilizes nicotine replacement therapy, and through the use of step-down dosing, gradually reduces the amount of nicotine in your system.

Second, the Habitrol® Take Control™ Support Program, developed by behavioral change and smoking cessation experts, recognizes that quitting smoking is a personal journey that occurs in six stages, known as the “Stages of Change.”

1. This *guide* will teach you how to move through the quit process by helping you to identify your reasons for quitting, manage urges to smoke, recover from slips and control your weight.
2. **1-888-HABITROL — our toll-free telephone support center**, staffed by smoking cessation professionals, will offer you friendly support and information. Call Monday through Friday between 9 a.m. and 8 p.m. ET or visit us at www.habitrol.com. You can use this resource to help

you get ready to quit, manage urges, cope with withdrawal symptoms, recover from slips and deal with smoking spouses and friends.

Working Towards Success — The Stages of Change

Quitting smoking is a process that begins long before your quit day. Researchers have determined that smokers go through the following six “Stages of Change” while on the road to becoming smoke-free.



II: Getting Started: Your Personal Quit Plan

Setting Your Quit Date

Now that you are motivated and committed, the next step is to choose a quit date within the next 2–3 weeks. Decide whether it will be easier to quit on a weekday when your smoking may already be restricted, or on the weekend when you are more relaxed. You may want to consider quitting at the beginning of the week, on a Sunday or Monday. It's best to choose a date when:

- Your stress level is low
- You'll be confronted with a minimum of smoking triggers
- You will not be in a social situation with other smokers, especially those that include alcohol

After completing this list, refer to it every day to reinforce your commitment to becoming a nonsmoker.

Rewarding Yourself

Many people get this far and forget to reward themselves. We know that we are more likely to keep going if we get rewarded. So, in the space below, write ways that you can reward yourself at least once a day.

If You Slip

Imagine that you go out with friends and end up having a cigarette. In other words, you “slipped.” This is not a relapse back to smoking. Don't let this mistake make you feel like a failure or like giving up. Get

Pre-Contemplation Stage:

If you are in this stage, you don't have any intention to quit smoking in the near future (i.e., within the next 6 months). You feel that the benefits you receive from smoking outweigh the costs and risks of smoking. This is called the *Pre-Contemplation Stage*. If you are in this stage right now, you are not ready to use the patch.

Contemplation Stage:

After a period of time, often years, you may begin to realize that you are “hooked,” and that smoking is affecting your health and the health of those around you. You begin to think about the benefits of quitting, but you know that quitting will be difficult. You are seriously considering quitting smoking sometime within the next six months, but you are ambivalent. This is called the *Contemplation Stage*. In this stage, you may read articles on ways to quit or on the health effects of smoking, while you would have ignored this information in the *Pre-Contemplation Stage*. You begin to imagine your life without cigarettes. You also begin to experiment with making changes. For example, you may be trying to delay your first cigarette of the day, smoke only half of some cigarettes, or not smoke in your house or car. If you are in this stage right now, you are still not ready to use the patch.

Preparation Stage:

When you've made the decision to quit within the next month, and you have experimented with making changes, you've entered the *Preparation Stage*. Most people who have purchased the patch for the first time are in this stage. Preparation is a good name for this stage because *thorough preparation greatly improves your chances of success*. Just “winging it” or relying on willpower alone is not enough. *You need a game plan that includes setting a quit date and committing to at least one person that you will make this change*. Using this guide, or visiting www.habitrol.com, as well as talking to the professionals at 1-888-HABITROL, can help you become fully prepared for your quit day and beyond.

Common withdrawal symptoms include intense cravings for nicotine, irritability, anxiety, depression, restlessness, difficulty concentrating, difficulty sleeping and increased appetite. Physical withdrawal symptoms usually peak within 24 to 72 hours after quitting, then decline over the next several weeks. Some smokers, however, may experience withdrawal symptoms for several months. Staying on this patch for the full eight weeks can reduce the withdrawal symptoms you experience. It is important to complete treatment. If you feel you need to use the patch for a longer period to keep from smoking, talk to your health care provider.

Smoking and your mind
The physical need for nicotine isn't the only reason you may find it difficult to quit smoking. You can also be psychologically dependent. Over time you've created strong associations or “triggers” to light



A quick tip — If you bought a pack of cigarettes to “just have one,” run the rest under cold water and then throw them out! Every single one! Don't allow yourself to think that you can keep cigarettes stashed away and still resist them. Remember, the only reason to keep cigarettes around is to smoke them!

The patch helps smokers quit smoking by reducing nicotine withdrawal symptoms. Many patch users will be able to stop smoking for at least a few days, but many will start smoking again. Most smokers usually have to try to quit several times before they completely stop.

Your own chances to stop smoking depend on how strongly you are addicted to nicotine, how much you want to quit, and how closely you follow a quitting program like the one that comes with this product. If you find you cannot stop or if you start smoking again after using this product, please talk to a health care professional who can help you find a program that may work better for you.

And, if for some reason you are not ready to try again right now, keep these materials until you are ready. Before you set your quit date, call us to help you prepare for a successful quit.

Nicotine Transdermal System
STOP SMOKING AID
PATCH



Your guide to help you successfully quit smoking

Keep this card in your wallet, for easy access anytime.



Nicotine Transdermal System
STOP SMOKING AID
PATCH



Call 1-888-HABITROL (1-888-422-4876) or visit us at www.habitrol.com.

Dr. Reddy's Laboratories, Inc. ©2015

Table of Contents

I: Thinking About Quitting.....3
 Why the Habitrol® Take Control™ Support Program Leads to Success.....3
 Working Towards Success — The Stages of Change5
 Which Way Now?.....8
 Why I Want to Stop Smoking.....8
 Health Benefits of Quitting.....9

II: Getting Started: Your Personal Quit Plan.....10
 Setting Your Quit Date.....10
 Understanding Nicotine Addiction.....10
 Smoking and your body.....10
 Smoking and your mind.....11
 Knowing your “triggers”.....12
 Taming Your “Triggers”.....12
 Reducing the Urge to Smoke.....12
 Building Your Supporting Cast.....14
 Countdown to Quit Day — 10 Steps to Success.....15
 Rewarding Yourself.....16
 If You Slip.....16

III: The Patch.....18
 Important Information About This Nicotine Transdermal System18
 How the Patch Works.....19
 How to Use the Patch.....20

IV: Weight Control Guide.....27

V: You Are On Your Way!.....29

VI: Your Daily Success Calendar.....30

VII: The Habitrol® Take Control™ Support Program.....34

Action Stage:

On the day that you quit smoking, you are in the *Action Stage*. It's time to set your plan in motion. You will probably experience cravings for nicotine and urges to have a cigarette throughout the day, but through preparation, you have developed multiple strategies for dealing with your “triggers,” withdrawal symptoms and those cravings or urges to smoke. It's important to start using the patch on the morning of your quit day to help reduce urges to smoke and other withdrawal symptoms, such as irritability and difficulty concentrating.

Maintenance and Termination Stages:

If you are able to remain smoke-free for 6 months, you enter the *Maintenance Stage*. You like your new life, but still have occasional urges. Watch out for overconfidence. Thinking you can smoke “just one cigarette” should be a blinking neon

warning sign. For most people, that one cigarette leads to two, then three, then full-time smoking again. Constantly remind yourself of the benefits you now enjoy as a nonsmoker. Have a plan ready to help you manage unexpected situations that may cause you stress and challenge your resolve to remain smoke-free. Most successful quitters will have occasional urges for many years.

Only about 1 in 5 quitters reach the *Termination Stage*, in which they have absolutely no temptation to smoke, and are 100% confident that they will never smoke again.

Your own chances to stop smoking depend on how strongly you are addicted to nicotine, how much you want to quit, and how closely you follow a quitting program like the one that comes with this product. If you find you cannot stop or if you start smoking again after using this product, please talk to a health care professional who can

help you find a program that may work better for you.

Which Way Now?

If you are planning on quitting within the next month or have recently quit, continue working through this guide to develop your personal quit plan, including the use of the patch. If you have any questions or need additional support, call the professionals at 1-888-HABITROL or visit us at www.habitrol.com. You will be encouraged by the wealth of information and support that is readily available.

If you are not planning on quitting soon, this isn't the time for you to use the patch, but the professionals at 1-888-HABITROL can still help you. They can offer you personalized feedback about your smoking, help you set a quit date, assist you with your ambivalence about quitting and provide you information about the resources available to you, even

before you're ready to quit. You can also help a friend or family member that is trying to quit by passing along this information about the Habitrol® Take Control™ Support Program.

Why I Want to Stop Smoking

Knowing that there are benefits to quitting is essential to your success. Writing down the reasons you want to quit on a card and keeping it with you to review can help you resist a temptation to smoke. Check off the reasons below that apply to you.

- I want control back — I'm almost always thinking about or smoking a cigarette. I feel like the cigarette is controlling me.
- I want to improve my chances of living a longer, healthier life.
- I want to spend my cigarette money on other things — maybe a vacation.

emotionally and physically. Be patient with them. Explain that you might be a little (or very) irritable for a while.

Building Your Supporting Cast

Quitting isn't easy. Sometimes you can feel all alone. This is where friends and family, both smokers and nonsmokers, can help. Take your friends and family through your quit plan. Once they understand why you are changing some of your old habits, they'll be in a better position to understand and support you.

Tell your smoking friends of your desire to quit (truth is, many of them may have that same desire). You might be pleasantly surprised how supportive they can be. If your smoking friends express an interest in quitting, give them our number, 1-888-HABITROL or visit us at www.habitrol.com, and we'll help get them started. You might even choose to quit at the same time so that you can work as a team supporting one another.

It's hard for people who have never smoked to understand what you're going through, both



2. Switch your focus. Deliberately switch your attention from having a cigarette onto something else, like reading or stretching.
3. Use the mental imagery to transform the urge into something manageable. For example,

My Trigger (In the past, I smoked...)	My Solution (Now, I will...)
<input type="checkbox"/> To concentrate	_____
<input type="checkbox"/> To relax	_____
<input type="checkbox"/> To give myself a lift	_____
<input type="checkbox"/> Because I was angry	_____
<input type="checkbox"/> Because I was bored	_____
<input type="checkbox"/> Because I felt stressed	_____
<input type="checkbox"/> _____	_____
<input type="checkbox"/> _____	_____

Remember when you first tried a cigarette? Remember how awful it tasted? Remember your initial dizziness? Nausea? The point is, it took some time for you to learn how to smoke. Now you can learn not to smoke. That takes time too. Sticking with the Habitrol® Take Control™ Support program can help. Call 1-888-HABITROL or visit us at www.habitrol.com. We can help you develop customized solutions to your personal triggers.

Knowing your “triggers”

Listed to the right are some of the common situations or activities that “trigger” smoking in many people and some suggestions on how to change your habits in order to reduce your urge to smoke.

Taming Your “Triggers”

Here are some other common “triggers.” Check off the ones that apply to you and write down how you will cope in the spaces below. Also, record other personal “triggers” and the things you plan to do instead.

Reducing the Urge to Smoke

Urges to smoke only last for a few seconds to a few minutes. Believe it or not, the urge to smoke will pass whether you smoke or not. When an urge strikes, try the following:

1. The “Three-Second Breathing” Exercise: Hold your breath for 3 seconds. Then slowly exhale through your mouth.

pursed lips. Hold your breath for 3 seconds. Then slowly exhale through your mouth.

If you smoke...Then try...

- With your morning coffee...**
Drinking hot chocolate, tea or orange juice
- After a meal...**
Leaving the table and doing something else or brushing your teeth
- After a stressful situation...**
Taking a walk or a 5 minute break
- Before bed...**
Reading or deep breathing
- On the telephone...**
Doodling on a notepad or talking while standing or walking
- At parties...**
Avoiding alcohol, and munch on healthy snacks
- Watching TV...**
Watching TV while holding something else, like a stress ball, or sitting in a different chair

III: The Patch

Important Information About This Nicotine Transdermal System

you find a program that may work better for you.

Ask your doctor before use if you:

- Have heart disease or an irregular heartbeat, or if you had a recent heart attack. Nicotine can increase your heart rate.
- Have high blood pressure not controlled with medication.
- Are allergic to any adhesives or patch ingredients or have skin problems, because you are more likely to get rashes.
- Are using a non-nicotine stop smoking drug.
- Take prescription medicine for depression or asthma. Your prescription dose may need to be adjusted.

18

only. Nicotine on your hands could get into your eyes and nose and could cause stinging, redness or irritation.

6. After 24 hours, remove the patch you have been wearing.

If you remove and apply the patch at about the same time each day, it will help you to remember when to put on a new patch. Choose a different place on your skin to apply the next patch and repeat



24

V: You Are On Your Way!

Congratulations again on your decision to quit smoking. By incorporating the behavior-changing tools discussed in this guide into your everyday life, you are on your way to successfully becoming and remaining smoke-free. Continue to reinforce your commitment to remain smoke-free by reading this guide again, or calling 1-888-HABITROL. While you are taking control with the Habitrol® Take Control™ Support Program, don't forget to celebrate your successes along the way.



Final Package Insert

Reference ID: 3954735

Keep out of reach of children and pets. Use patches have enough nicotine to poison children and pets. If swallowed, get medical help or contact a Poison Control Center right away.

How the Patch Works

The patch is a nicotine transdermal system skin patch containing nicotine, the addictive agent in tobacco products. When you wear a patch, it gradually releases nicotine into your bloodstream through the skin. By replacing some of the nicotine to which your body has become accustomed from smoking, the patch helps reduce the nicotine withdrawal symptoms many people normally feel when they stop smoking. By offering three patches with different nicotine dose levels, this patch uses a step-down dosing system that allows you to gradually reduce your nicotine level by changing the patch you wear (moving to a lower dose) over

an eight-week period. Nicotine replacement therapy can reduce nicotine withdrawal symptoms such as irritability, anxiety, restlessness, headaches, difficulty sleeping and concentrating, increased appetite, and craving for nicotine. By helping to reduce your physical nicotine withdrawal symptoms, the patch helps you to concentrate on the psychological aspects of quitting and to change your habits that "trigger" your nicotine cravings.

This patch program takes 8 weeks to complete. It is important to complete treatment. If you feel you need to use the patch for a longer period to keep from smoking, talk to your health care provider.

Why wearing a nicotine patch isn't as bad as smoking

By placing a nicotine patch on your skin, you are NOT inhaling the harmful tars, toxins and chemicals found in cigarettes. These are the most dangerous parts of the cigarette.

steps 1 through 5. Do not reapply a patch to a previously used skin site for at least 1 week. Do not leave the patch on for more than 24 hours, because it may irritate your skin and loses strength after 24 hours. Do not wear more than one patch at the same time, and do not cut a patch in half or into smaller pieces to wear.

If you have trouble sleeping

You should wear the patch 24 hours a day. This may help overcome your morning cravings for nicotine. However, if you have vivid dreams or other sleep disturbances, you may remove the patch at bedtime and apply a new patch the following morning. The patch should be applied at approximately the same time each day.

If your patch comes off

If your patch falls off, put on a new one. Then remove this new patch at your regular time to keep your schedule the same. When

applying the patch, be sure to press it firmly onto your skin with the palm of your hand for about 10 seconds, making sure that the patch sticks well, especially around the edges.

If your patch gets wet

Water will not harm or affect the patch you are wearing. You can take a bath or a shower, or you can swim while you are wearing the patch.

Disposing of the patch

Save pouch to use for patch disposal. Dispose of the used patches by folding sticky ends together and putting in pouch. Keep all used patches out of reach of children and pets.

Storage instructions

Keep the patch in its protective pouch until you are ready to use it. Store your patches between 20-25°C (68-77°F), because the patch is sensitive to heat. The inside of your car, for example, can reach much

And, because you're not smoking while wearing the patch, there's no second-hand smoke or odors to affect your family and friends.

How to Use the Patch

It is important that you are firmly committed to giving up smoking.

Warnings:

- If you are pregnant or breast-feeding, only use this medicine on the advice of your health care provider. Smoking can seriously harm your child. Try to stop smoking without using any nicotine replacement medicine. This medicine is believed to be safer than smoking. However, the risks to your child from this medicine are not fully known.
- Ask a doctor before use if you have heart disease, have had a recent heart attack, or have an irregular heartbeat. Nicotine can increase your heart rate.
- Ask a doctor before use if you have high blood pressure not controlled with medication. Nicotine can increase your blood pressure.
- Ask a doctor before use if you have an allergy to adhesive tape or have skin problems because you are more likely to get rashes with patch use.
- Ask a doctor before use if you have stomach ulcer or diabetes.
- Ask a doctor before use if you have history of seizures.
- Ask a doctor or pharmacist before use if you are using a non-nicotine stop smoking drug.
- Ask a doctor or pharmacist before use if you are taking a prescription medicine for depression or asthma. Your prescription dose may need to be adjusted.
- When using this product, if you have vivid dreams or other sleep disturbances, you may remove the patch at bedtime and apply a new one in the morning.
- Stop use and ask a doctor if skin redness caused by the patch does

19

higher temperatures in the summer. Keep all unused patches out of the reach of children and pets.

If your skin reacts to the patch

When you first put on a patch, mild itching, burning, or tingling at the patch application site is normal and should go away within an hour. After you remove a patch, the skin under the patch might be somewhat red. Your skin should not stay red for more than a day. If you have a skin rash or redness caused by the patch that does not go away after 4 days, or your skin swells, call your doctor. Do not put on a new patch; you may be allergic to one of the components of the patch.

Other side effects you may experience

While wearing a nicotine replacement patch, you may experience one or more of the following side effects: nausea, dizziness, dry mouth, diarrhea, nervousness or restlessness,

controlled with medication. Nicotine can increase your blood pressure.

- Ask a doctor before use if you have an allergy to adhesive tape or have skin problems because you are more likely to get rashes with patch use.
- Ask a doctor before use if you have stomach ulcer or diabetes.
- Ask a doctor before use if you have history of seizures.
- Ask a doctor or pharmacist before use if you are using a non-nicotine stop smoking drug.
- Ask a doctor or pharmacist before use if you are taking a prescription medicine for depression or asthma. Your prescription dose may need to be adjusted.
- When using this product, if you have vivid dreams or other sleep disturbances, you may remove the patch at bedtime and apply a new one in the morning.
- Stop use and ask a doctor if skin redness caused by the patch does

IV: Weight Control Guide

Why You May Gain Weight After You Quit Smoking

It's perfectly normal to gain some weight when you stop smoking. But don't panic. Sometimes the weight change is simply a result of a change in your body's metabolism. This is caused by your body's withdrawal from nicotine. Your metabolism is simply returning to that of a nonsmoker. Besides, the average weight gain is only about five to ten pounds, and as you will see, there are things you can do to help control it.

Another source of extra pounds is the need for oral gratification. If you use food as a temporary substitute for cigarettes, you may gain weight. If you feel some concern about your weight, remember: a small weight gain is better than the high health risks associated with smoking.

- Do not leave the patch on for more than 24 hours because it may irritate your skin and it loses strength after 24 hours.
- To avoid possible burns, remove patch before undergoing any MRI (Magnetic Resonance Imaging) procedures.
- It is important to complete treatment. If you feel you need to use the patch for a longer period to keep from smoking, talk to your health care provider. The patch has been tested in 3 months studies and long term studies have not been conducted.

See chart on following page.

First, it is important that you are well prepared to give up smoking. If you are under 18 years of age, ask a doctor before use.

If you smoke more than 10 cigarettes per day: Begin with Step 1. Use the 21 mg patch for 4 weeks, then 14 mg patch for 2 weeks, and 7 mg patch for 2 weeks.		
Weeks 1-4	Weeks 5 and 6	Weeks 7 and 8
Step 1 21 mg/day	Step 2 14 mg/day	Step 3 7 mg/day

If you smoke 10 or less cigarettes per day: Do not begin with Step 1. Begin with Step 2, 14 mg patch for 6 weeks, and 7 mg patch for 2 weeks.	
Weeks 1-6	Weeks 7 and 8
Step 2 14 mg/day	Step 3 7 mg/day

21

How to control your weight through the benefits of exercise

One of the best ways to control your weight is to move more. So if you already exercise regularly, keep it up. If not, start with small increases in your daily activity.

We're NOT talking about running a marathon. We're talking about walking to the store instead of hopping in the car. Or, walking up three flights of stairs instead of taking the elevator. Maybe you should rediscover your bicycle, take up golf or go for a swim.

A little exercise will do more than you think:

- It reduces possible weight gain while quitting smoking
- It can improve your sleeping
- It can make you feel better in general

The following tips will help you minimize the weight gain while you quit smoking.

How to control your weight by watching how and what you eat

Some general tips:

- Replace fattening snack foods with healthy, low-calorie snacks
- Keep low-fat, low-calorie snacks in the car
- Broil, boil, poach, or steam food instead of frying
- Remove fat from meat and skin from poultry
- Eat slowly and you'll feel full sooner on less food
- Try eating several small meals a day instead of three big ones
- Drink a large glass of water before a meal

27

28

cut or irritated in any way. Immediately before applying the patch, wash your hands and the skin area with plain soap and water and dry completely. Avoid using any soap, lotion, hand cream, tanning lotion or oil, bath oil or insect repellent that contains alcohol, lanolin or glycerin as a moisturizer. These products can leave a moisturizing film on your skin, which can interfere with the adherence of the patch.



2. Do not remove the patch from its sealed, protective pouch until you are ready to use it. Carefully cut open the child-resistant pouch along the dotted line, as indicated. If the new patch is cut, throw it away. Save pouch to use for patch disposal. Dispose of the used patches by folding

cut or irritated in any way. Immediately before applying the patch, wash your hands and the skin area with plain soap and water and dry completely. Avoid using any soap, lotion, hand cream, tanning lotion or oil, bath oil or insect repellent that contains alcohol, lanolin or glycerin as a moisturizer. These products can leave a moisturizing film on your skin, which can interfere with the adherence of the patch.

3. A shiny protective liner covers the sticky side of the patch where it contacts the skin. The liner has a precut slit to help you remove it from the patch. With the silver side facing you, pull the liner away from the patch, starting at the precut slit. Hold the patch at the edge (touch the sticky side as little as possible) and pull off the other piece of the protective liner. Throw this liner away.

22

23



Keep this card in your wallet, for easy access anytime.

10 Steps to Success

1. Tape a list of your reasons for quitting on the mirror. Go over them daily.
2. Tell one or two friends and your family. Ask for the specific help that you think you will need from each of them.
3. Use the Habitrol® Take Control™ Support Program.
4. Know your "triggers," and practice your coping strategies.
5. Change your routines (for example, sit in a different chair, not your smoking chair).
6. Plan how you will spend all the money you save from not smoking.
7. Freshen your environment. On the day before your quit date, clean your clothes, car and any rooms where you spend a lot of time.
8. Throw away any remaining cigarettes, ashtrays and lighters the night before your quit day.
9. Have plenty of low-calorie snacks available.
10. Use the patch as directed.



CHECK OFF EACH DAY AS A NONSMOKER AND WATCH THOSE SUCCESSFUL DAY'S ACCUMULATE

WEEK 5	DAY 30	DAY 31	DAY 32	DAY 33	DAY 34	DAY 35
DAY 29 Can you believe it? Four whole weeks! Step down to the 14 mg/day patch, which is Step 2	Take one day at a time	Admit it, you're feeling great!	Don't forget to call a friend if you need some extra TLC	Ever think about how much money you'll save being a nonsmoker?	Keep exercising	List any surprise triggers and how you tamed them
WEEK 6	DAY 37	DAY 38	DAY 39	DAY 40	DAY 41	DAY 42
DAY 36 Expecting a high pressure day? Prepare yourself by reviewing Taming Your "Triggers" (page 12)	Try something new Take up a hobby	Don't forget to switch your focus!	Have you rewarded yourself lately?	See how many successful smoke free days you've had!	Take a walk and renew your commitment	Is it time to buy more patches?
WEEK 7	DAY 44	DAY 45	DAY 46	DAY 47	DAY 48	DAY 49
DAY 43 A big congratulations! You're coming down the home stretch. Step down to the 7 mg/day patch, which is Step 3	Think about celebrating a friend to stop smoking!	Review your reasons for quitting	Remember to keep a healthy snack on hand	Need some added support? Call a friend or 1 888 HABITROL or visit www.habitrol.com	Daily exercise is still your best defense	Stronger than you thought you were. Congratulations!
WEEK 8	DAY 51	DAY 52	DAY 53	DAY 54	DAY 55	DAY 56
DAY 50 You hit week on the patch!	You're a hero!	Think of a way to celebrate	Visit us at www.habitrol.com	Are you feeling a great sense of freedom?	It is important to complete treatment Buy yourself a gift today	If you feel you need to use the patch for a longer period to keep from smoking, talk to your health care provider You are a success!

Toll Free: 1-888-HABITROL (1-888-422-4876) or visit us at www.habitrol.com

CHECK OFF EACH DAY AS A NONSMOKER AND WATCH THOSE SUCCESSFUL DAY'S ACCUMULATE

WEEK 1: QUIT	DAY 2	DAY 3	DAY 4	DAY 5	DAY 6	DAY 7
DAY 1 It's your first day as a non-smoker! Congratulations! Start off the day with your first Step 1 (21 mg) patch instead of a cigarette!	Avoid your triggers today	Remember to switch your focus when you think of a cigarette. Think of something else, like how much healthier you'll be.	Are you rewarding yourself?	Dry clean or wash clothes to get rid of the cigarette smell	Try the Three Second Breathing Exercise	It's a good idea to review Taming Your "Triggers" (page 12)
WEEK 2	DAY 9	DAY 10	DAY 11	DAY 12	DAY 13	DAY 14
DAY 8 You made it through the crucial first week!	Don't forget to change your patch every day	What triggers did you tame today?	Have you reached out to your support?	Did you remember to pick up more patches?	Make sure you have healthy snacks on hand	Way to go! Two whole weeks without smoking!
WEEK 3	DAY 16	DAY 17	DAY 18	DAY 19	DAY 20	DAY 21
DAY 15 Treat yourself today! Go to the movies or the park.	Remember to switch your focus when the urge strikes	List the triggers you've tamed. Are you proud of yourself?	Check your patch supply. Do you need more patches?	Are you exercising enough? Exercise is a great stress reliever!	Think about encouraging a friend to stop smoking!	Celebrate! Three weeks of no smoking!
WEEK 4	DAY 23	DAY 24	DAY 25	DAY 26	DAY 27	DAY 28
DAY 22 You deserve a reward! Do something special for yourself	Doesn't food taste better these days?	Was today trigger free?	The urge to smoke will pass. Stay committed!	Remember to get up and move. Try a new exercise today	Treat yourself to a bottle of cologne to celebrate your improved sense of smell.	Tomorrow, you step down to the next lower strength patch

Toll Free: 1-888-HABITROL (1-888-422-4876) or visit us at www.habitrol.com

You may want to note how you're feeling, too. Each day renew your commitment to being a nonsmoker.

*If you smoke 10 or less cigarettes per day, start with the Step 2, 14 mg patch for 6 weeks, then step 3, 7 mg patch for 2 weeks.

29

30

31

32

33

34

This is a representation of an electronic record that was signed electronically and this page is the manifestation of the electronic signature.

/s/

THERESA M MICHELE
07/05/2016