

**SPORANOX<sup>®</sup>**  
**(itraconazole)**  
**Capsules**

**BOXED WARNING**


**Congestive Heart Failure, Cardiac Effects and Drug Interactions:**

**SPORANOX<sup>®</sup> (itraconazole) Capsules should not be administered for the treatment of onychomycosis in patients with evidence of ventricular dysfunction such as congestive heart failure (CHF) or a history of CHF.** If signs or symptoms of congestive heart failure occur during administration of SPORANOX<sup>®</sup> Capsules, discontinue administration. When itraconazole was administered intravenously to dogs and healthy human volunteers, negative inotropic effects were seen. (See CONTRAINDICATIONS, WARNINGS, PRECAUTIONS: Drug Interactions, ADVERSE REACTIONS: Post-marketing Experience, and CLINICAL PHARMACOLOGY: Special Populations for more information.)

**Drug Interactions: Coadministration of the following drugs are contraindicated with SPORANOX<sup>®</sup> Capsules: methadone, disopyramide, dofetilide, dronedarone, quinidine, ergot alkaloids (such as dihydroergotamine, ergometrine (ergonovine), ergotamine, methylergometrine (methylergonovine)), irinotecan, lurasidone, oral midazolam, pimozide, triazolam, felodipine, nisoldipine, ranolazine, eplerenone, cisapride, lovastatin, simvastatin, ticagrelor and, in subjects with varying degrees of renal or hepatic impairment, colchicine, fesoterodine, telithromycin and solifenacin.** See PRECAUTIONS: Drug Interactions Section for specific examples. Coadministration with itraconazole can cause elevated plasma concentrations of these drugs and may increase or prolong both the pharmacologic effects and/or adverse reactions to these drugs. For example, increased plasma concentrations of some of these drugs can lead to QT prolongation and ventricular tachyarrhythmias including occurrences of torsades de pointes, a potentially fatal arrhythmia. See CONTRAINDICATIONS and WARNINGS Sections, and PRECAUTIONS: Drug Interactions Section for specific examples.

**DESCRIPTION**

SPORANOX<sup>®</sup> is the brand name for itraconazole, an azole antifungal agent. Itraconazole is a 1:1:1:1 racemic mixture of four diastereomers (two enantiomeric pairs), each possessing three chiral centers. It may be represented by the following structural formula and nomenclature:



(±)-1-[(R\*)-sec-butyl]-4-[p-[4-[p-[(2R\*,4S\*)-2-(2,4-dichlorophenyl)-2-(1H-1,2,4-triazol-1-ylmethyl)-1,3-dioxolan-4-yl]methoxy]phenyl]-1-piperazinyl]phenyl]-Δ<sup>2</sup>-1,2,4-triazolin-5-one mixture with (±)-1-[(R\*)-sec-butyl]-4-[p-[4-[p-[(2S\*,4R\*)-2-(2,4-dichlorophenyl)-2-(1H-1,2,4-triazol-1-ylmethyl)-1,3-dioxolan-4-yl]methoxy]phenyl]-1-piperazinyl]phenyl]-Δ<sup>2</sup>-1,2,4-triazolin-5-one

or

(±)-1-[(RS)-sec-butyl]-4-[p-[4-[p-[(2R,4S)-2-(2,4-dichlorophenyl)-2-(1H-1,2,4-triazol-1-ylmethyl)-1,3-dioxolan-4-yl]methoxy]phenyl]-1-piperazinyl]phenyl]-Δ<sup>2</sup>-1,2,4-triazolin-5-one

Itraconazole has a molecular formula of C<sub>35</sub>H<sub>38</sub>Cl<sub>2</sub>N<sub>8</sub>O<sub>4</sub> and a molecular weight of 705.64. It is a white to slightly yellowish powder. It is insoluble in water, very slightly soluble in alcohols, and freely soluble in dichloromethane. It has a pKa of 3.70 (based on extrapolation of values obtained from methanolic solutions) and a log (n-octanol/water) partition coefficient of 5.66 at pH 8.1.

SPORANOX<sup>®</sup> Capsules contain 100 mg of itraconazole coated on sugar spheres (composed of sucrose, maize starch, and purified water). Inactive ingredients are hard gelatin capsule, hypromellose, polyethylene glycol (PEG) 20,000, titanium dioxide, FD&C Blue No. 1, FD&C Blue No. 2, D&C Red No. 22 and D&C Red No. 28.

## CLINICAL PHARMACOLOGY

### Pharmacokinetics and Metabolism:

#### General Pharmacokinetic Characteristics

Peak plasma concentrations of itraconazole are reached within 2 to 5 hours following oral administration. As a consequence of non-linear pharmacokinetics, itraconazole accumulates in plasma during multiple dosing. Steady-state concentrations are generally reached within about 15 days, with C<sub>max</sub> values of 0.5 µg/ml, 1.1 µg/ml and 2.0 µg/ml after oral administration of 100 mg once daily, 200 mg once daily and 200 mg b.i.d., respectively. The terminal half-life of itraconazole generally ranges from 16 to 28 hours after single dose and increases to 34 to 42 hours with repeated dosing. Once treatment is stopped, itraconazole plasma concentrations decrease to an almost undetectable concentration within 7 to 14 days, depending on the dose and duration of treatment. Itraconazole mean total plasma clearance following

intravenous administration is 278 ml/min. Itraconazole clearance decreases at higher doses due to saturable hepatic metabolism.

### Absorption

Itraconazole is rapidly absorbed after oral administration. Peak plasma concentrations of itraconazole are reached within 2 to 5 hours following an oral capsule dose. The observed absolute oral bioavailability of itraconazole is about 55%.

The oral bioavailability of itraconazole is maximal when SPORANOX<sup>®</sup> (itraconazole) Capsules are taken immediately after a full meal. Absorption of itraconazole capsules is reduced in subjects with reduced gastric acidity, such as subjects taking medications known as gastric acid secretion suppressors (e.g., H<sub>2</sub>-receptor antagonists, proton pump inhibitors) or subjects with achlorhydria caused by certain diseases. (See PRECAUTIONS: Drug Interactions.) Absorption of itraconazole under fasted conditions in these subjects is increased when SPORANOX<sup>®</sup> Capsules are administered with an acidic beverage (such as a non-diet cola). When SPORANOX<sup>®</sup> Capsules were administered as a single 200-mg dose under fasted conditions with non-diet cola after ranitidine pretreatment, a H<sub>2</sub>-receptor antagonist, itraconazole absorption was comparable to that observed when SPORANOX<sup>®</sup> Capsules were administered alone. (See PRECAUTIONS: Drug Interactions.)

Itraconazole exposure is lower with the Capsule formulation than with the Oral Solution when the same dose of drug is given. (See WARNINGS)

### Distribution

Most of the itraconazole in plasma is bound to protein (99.8%), with albumin being the main binding component (99.6% for the hydroxy-metabolite). It has also a marked affinity for lipids. Only 0.2% of the itraconazole in plasma is present as free drug. Itraconazole is distributed in a large apparent volume in the body (>700 L), suggesting extensive distribution into tissues. Concentrations in lung, kidney, liver, bone, stomach, spleen and muscle were found to be two to three times higher than corresponding concentrations in plasma, and the uptake into keratinous tissues, skin in particular, up to four times higher. Concentrations in the cerebrospinal fluid are much lower than in plasma.

### Metabolism

Itraconazole is extensively metabolized by the liver into a large number of metabolites. *In vitro* studies have shown that CYP3A4 is the major enzyme involved in the metabolism of itraconazole. The main metabolite is hydroxy-itraconazole, which has *in vitro* antifungal activity comparable to itraconazole; trough plasma concentrations of this metabolite are about twice those of itraconazole.

## Excretion

Itraconazole is excreted mainly as inactive metabolites in urine (35%) and in feces (54%) within one week of an oral solution dose. Renal excretion of itraconazole and the active metabolite hydroxy-itraconazole account for less than 1% of an intravenous dose. Based on an oral radiolabeled dose, fecal excretion of unchanged drug ranges from 3% to 18% of the dose.

As re-distribution of itraconazole from keratinous tissues appears to be negligible, elimination of itraconazole from these tissues is related to epidermal regeneration. Contrary to plasma, the concentration in skin persists for 2 to 4 weeks after discontinuation of a 4-week treatment and in nail keratin – where itraconazole can be detected as early as 1 week after start of treatment – for at least six months after the end of a 3-month treatment period.

## Special Populations:

### Renal Impairment:

Limited data are available on the use of oral itraconazole in patients with renal impairment. A pharmacokinetic study using a single 200-mg oral dose of itraconazole was conducted in three groups of patients with renal impairment (uremia: n=7; hemodialysis: n=7; and continuous ambulatory peritoneal dialysis: n=5). In uremic subjects with a mean creatinine clearance of 13 mL/min.  $\times 1.73 \text{ m}^2$ , the exposure, based on AUC, was slightly reduced compared with normal population parameters. This study did not demonstrate any significant effect of hemodialysis or continuous ambulatory peritoneal dialysis on the pharmacokinetics of itraconazole ( $T_{\max}$ ,  $C_{\max}$ , and  $AUC_{0-8h}$ ). Plasma concentration-versus-time profiles showed wide intersubject variation in all three groups. After a single intravenous dose, the mean terminal half-lives of itraconazole in patients with mild (defined in this study as CrCl 50-79 ml/min), moderate (defined in this study as CrCl 20-49 ml/min), and severe renal impairment (defined in this study as CrCl <20 ml/min) were similar to that in healthy subjects (range of means 42-49 hours vs 48 hours in renally impaired patients and healthy subjects, respectively). Overall exposure to itraconazole, based on AUC, was decreased in patients with moderate and severe renal impairment by approximately 30% and 40%, respectively, as compared with subjects with normal renal function. Data are not available in renally impaired patients during long-term use of itraconazole. Dialysis has no effect on the half-life or clearance of itraconazole or hydroxy-itraconazole. (See PRECAUTIONS and DOSAGE AND ADMINISTRATION.)

### Hepatic Impairment:

Itraconazole is predominantly metabolized in the liver. A pharmacokinetic study was conducted in 6 healthy and 12 cirrhotic subjects who were administered a single 100-mg dose of itraconazole as capsule. A statistically significant reduction in mean  $C_{\max}$  (47%) and a twofold increase in the elimination half-life ( $37 \pm 17$  hours vs.  $16 \pm 5$  hours) of itraconazole were noted in cirrhotic subjects compared with healthy subjects. However, overall exposure to

itraconazole, based on AUC, was similar in cirrhotic patients and in healthy subjects. Data are not available in cirrhotic patients during long-term use of itraconazole. (See CONTRAINDICATIONS, PRECAUTIONS: Drug Interactions and DOSAGE AND ADMINISTRATION.)

#### **Decreased Cardiac Contractility:**

When itraconazole was administered intravenously to anesthetized dogs, a dose-related negative inotropic effect was documented. In a healthy volunteer study of itraconazole intravenous infusion, transient, asymptomatic decreases in left ventricular ejection fraction were observed using gated SPECT imaging; these resolved before the next infusion, 12 hours later. If signs or symptoms of congestive heart failure appear during administration of SPORANOX<sup>®</sup> Capsules, SPORANOX<sup>®</sup> should be discontinued. (See BOXED WARNING, CONTRAINDICATIONS, WARNINGS, PRECAUTIONS: Drug Interactions and ADVERSE REACTIONS: Post-marketing Experience for more information.)

### **MICROBIOLOGY**

#### **Mechanism of Action:**

*In vitro* studies have demonstrated that itraconazole inhibits the cytochrome P450-dependent synthesis of ergosterol, which is a vital component of fungal cell membranes.

#### **Activity In Vitro and in Clinical Infections:**

Itraconazole exhibits *in vitro* activity against *Blastomyces dermatitidis*, *Histoplasma capsulatum*, *Histoplasma duboisii*, *Aspergillus flavus*, *Aspergillus fumigatus*, and *Trichophyton* species (See INDICATIONS AND USAGE, Description of Clinical Studies).

Correlation between minimum inhibitory concentration (MIC) results *in vitro* and clinical outcome has yet to be established for azole antifungal agents.

#### **Drug Resistance:**

Isolates from several fungal species with decreased susceptibility to itraconazole have been isolated *in vitro* and from patients receiving prolonged therapy.

Itraconazole is not active against *Zygomycetes* (e.g., *Rhizopus* spp., *Rhizomucor* spp., *Mucor* spp. and *Absidia* spp.), *Fusarium* spp., *Scedosporium* spp. and *Scopulariopsis* spp.

#### **Cross-resistance:**

Several *in vitro* studies have reported that some fungal clinical isolates with reduced susceptibility to one azole antifungal agent may also be less susceptible to other azole derivatives. The finding of cross-resistance is dependent on a number of factors, including the

species evaluated, its clinical history, the particular azole compounds compared, and the type of susceptibility test that is performed.

Studies (both *in vitro* and *in vivo*) suggest that the activity of amphotericin B may be suppressed by prior azole antifungal therapy. As with other azoles, itraconazole inhibits the <sup>14</sup>C-demethylation step in the synthesis of ergosterol, a cell wall component of fungi. Ergosterol is the active site for amphotericin B. In one study the antifungal activity of amphotericin B against *Aspergillus fumigatus* infections in mice was inhibited by ketoconazole therapy. The clinical significance of test results obtained in this study is unknown.

## INDICATIONS AND USAGE

SPORANOX<sup>®</sup> (itraconazole) Capsules are indicated for the treatment of the following fungal infections in immunocompromised and non-immunocompromised patients:

1. Blastomycosis, pulmonary and extrapulmonary
2. Histoplasmosis, including chronic cavitory pulmonary disease and disseminated, non-meningeal histoplasmosis, and
3. Aspergillosis, pulmonary and extrapulmonary, in patients who are intolerant of or who are refractory to amphotericin B therapy.

Specimens for fungal cultures and other relevant laboratory studies (wet mount, histopathology, serology) should be obtained before therapy to isolate and identify causative organisms. Therapy may be instituted before the results of the cultures and other laboratory studies are known; however, once these results become available, anti-infective therapy should be adjusted accordingly.

SPORANOX<sup>®</sup> Capsules are also indicated for the treatment of the following fungal infections in non-immunocompromised patients:

1. Onychomycosis of the toenail, with or without fingernail involvement, due to dermatophytes (tinea unguium), and
2. Onychomycosis of the fingernail due to dermatophytes (tinea unguium).

Prior to initiating treatment, appropriate nail specimens for laboratory testing (KOH preparation, fungal culture, or nail biopsy) should be obtained to confirm the diagnosis of onychomycosis.

(See CLINICAL PHARMACOLOGY: Special Populations, CONTRAINDICATIONS, WARNINGS, and ADVERSE REACTIONS: Post-marketing Experience for more information.)

### **Description of Clinical Studies:**

#### **Blastomycosis:**

Analyses were conducted on data from two open-label, non-concurrently controlled studies (N=73 combined) in patients with normal or abnormal immune status. The median dose was 200 mg/day. A response for most signs and symptoms was observed within the first 2 weeks, and all signs and symptoms cleared between 3 and 6 months. Results of these two studies demonstrated substantial evidence of the effectiveness of itraconazole for the treatment of blastomycosis compared with the natural history of untreated cases.

#### **Histoplasmosis:**

Analyses were conducted on data from two open-label, non-concurrently controlled studies (N=34 combined) in patients with normal or abnormal immune status (not including HIV-infected patients). The median dose was 200 mg/day. A response for most signs and symptoms was observed within the first 2 weeks, and all signs and symptoms cleared between 3 and 12 months. Results of these two studies demonstrated substantial evidence of the effectiveness of itraconazole for the treatment of histoplasmosis, compared with the natural history of untreated cases.

#### **Histoplasmosis in HIV-infected patients:**

Data from a small number of HIV-infected patients suggested that the response rate of histoplasmosis in HIV-infected patients is similar to that of non-HIV-infected patients. The clinical course of histoplasmosis in HIV-infected patients is more severe and usually requires maintenance therapy to prevent relapse.

#### **Aspergillosis:**

Analyses were conducted on data from an open-label, “single-patient-use” protocol designed to make itraconazole available in the U.S. for patients who either failed or were intolerant of amphotericin B therapy (N=190). The findings were corroborated by two smaller open-label studies (N=31 combined) in the same patient population. Most adult patients were treated with a daily dose of 200 to 400 mg, with a median duration of 3 months. Results of these studies demonstrated substantial evidence of effectiveness of itraconazole as a second-line therapy for the treatment of aspergillosis compared with the natural history of the disease in patients who either failed or were intolerant of amphotericin B therapy.

### **Onychomycosis of the toenail:**

Analyses were conducted on data from three double-blind, placebo-controlled studies (N=214 total; 110 given SPORANOX<sup>®</sup> Capsules) in which patients with onychomycosis of the toenails received 200 mg of SPORANOX<sup>®</sup> Capsules once daily for 12 consecutive weeks. Results of these studies demonstrated mycologic cure, defined as simultaneous occurrence of negative KOH plus negative culture, in 54% of patients. Thirty-five percent (35%) of patients were considered an overall success (mycologic cure plus clear or minimal nail involvement with significantly decreased signs) and 14% of patients demonstrated mycologic cure plus clinical cure (clearance of all signs, with or without residual nail deformity). The mean time to overall success was approximately 10 months. Twenty-one percent (21%) of the overall success group had a relapse (worsening of the global score or conversion of KOH or culture from negative to positive).

### **Onychomycosis of the fingernail:**

Analyses were conducted on data from a double-blind, placebo-controlled study (N=73 total; 37 given SPORANOX<sup>®</sup> Capsules) in which patients with onychomycosis of the fingernails received a 1-week course (pulse) of 200 mg of SPORANOX<sup>®</sup> Capsules b.i.d., followed by a 3-week period without SPORANOX<sup>®</sup>, which was followed by a second 1-week pulse of 200 mg of SPORANOX<sup>®</sup> Capsules b.i.d. Results demonstrated mycologic cure in 61% of patients. Fifty-six percent (56%) of patients were considered an overall success and 47% of patients demonstrated mycologic cure plus clinical cure. The mean time to overall success was approximately 5 months. None of the patients who achieved overall success relapsed.

## **CONTRAINDICATIONS**

### **Congestive Heart Failure:**

SPORANOX<sup>®</sup> (itraconazole) Capsules should not be administered for the treatment of onychomycosis in patients with evidence of ventricular dysfunction such as congestive heart failure (CHF) or a history of CHF. (See BOXED WARNING, WARNINGS, PRECAUTIONS: Drug Interactions-Calcium Channel Blockers, ADVERSE REACTIONS: Post-marketing Experience, and CLINICAL PHARMACOLOGY: Special Populations.)

### **Drug Interactions:**

Coadministration of a number of CYP3A4 substrates are contraindicated with SPORANOX<sup>®</sup>. Plasma concentrations increase for the following drugs: methadone, disopyramide, dofetilide, dronedarone, quinidine, ergot alkaloids (such as dihydroergotamine, ergometrine (ergonovine), ergotamine, methylergometrine (methylergonovine)), irinotecan, lurasidone, oral midazolam, pimozide, triazolam, felodipine, nisoldipine, ranolazine, eplerenone, cisapride, lovastatin, simvastatin, ticagrelor, and, in subjects with varying degrees of renal or hepatic impairment, colchicine, fesoterodine, telithromycin and solifenacin. (See PRECAUTIONS: Drug

Interactions Section for specific examples.) This increase in drug concentrations caused by coadministration with itraconazole may increase or prolong both the pharmacologic effect and/or adverse reactions to these drugs. For example, increased plasma concentrations of some of these drugs can lead to QT prolongation and ventricular tachyarrhythmias including occurrences of torsade de pointes, a potentially fatal arrhythmia. Specific examples are listed in PRECAUTIONS: Drug Interactions.

SPORANOX<sup>®</sup> should not be administered for the treatment of onychomycosis to pregnant patients or to women contemplating pregnancy.

SPORANOX<sup>®</sup> is contraindicated for patients who have shown hypersensitivity to itraconazole. There is limited information regarding cross-hypersensitivity between itraconazole and other azole antifungal agents. Caution should be used when prescribing SPORANOX<sup>®</sup> to patients with hypersensitivity to other azoles.

## **WARNINGS**

### **Hepatic Effects:**

**SPORANOX<sup>®</sup> has been associated with rare cases of serious hepatotoxicity, including liver failure and death. Some of these cases had neither pre-existing liver disease nor a serious underlying medical condition, and some of these cases developed within the first week of treatment. If clinical signs or symptoms develop that are consistent with liver disease, treatment should be discontinued and liver function testing performed. Continued SPORANOX<sup>®</sup> use or reinstatement of treatment with SPORANOX<sup>®</sup> is strongly discouraged unless there is a serious or life-threatening situation where the expected benefit exceeds the risk. (See PRECAUTIONS: Information for Patients and ADVERSE REACTIONS.)**

### **Cardiac Dysrhythmias:**

Life-threatening cardiac dysrhythmias and/or sudden death have occurred in patients using drugs such as cisapride, pimozide, methadone, or quinidine concomitantly with SPORANOX<sup>®</sup> and/or other CYP3A4 inhibitors. Concomitant administration of these drugs with SPORANOX<sup>®</sup> is contraindicated. (See BOXED WARNING, CONTRAINDICATIONS, and PRECAUTIONS: Drug Interactions.)

### **Cardiac Disease:**

**SPORANOX<sup>®</sup> Capsules should not be administered for the treatment of onychomycosis in patients with evidence of ventricular dysfunction such as congestive heart failure (CHF) or a history of CHF. SPORANOX<sup>®</sup> Capsules should not be used for other indications**

in patients with evidence of ventricular dysfunction unless the benefit clearly outweighs the risk.

For patients with risk factors for congestive heart failure, physicians should carefully review the risks and benefits of SPORANOX<sup>®</sup> therapy. These risk factors include cardiac disease such as ischemic and valvular disease; significant pulmonary disease such as chronic obstructive pulmonary disease; and renal failure and other edematous disorders. Such patients should be informed of the signs and symptoms of CHF, should be treated with caution, and should be monitored for signs and symptoms of CHF during treatment. If signs or symptoms of CHF appear during administration of SPORANOX<sup>®</sup> Capsules, discontinue administration.

Itraconazole has been shown to have a negative inotropic effect. When itraconazole was administered intravenously to anesthetized dogs, a dose-related negative inotropic effect was documented. In a healthy volunteer study of itraconazole intravenous infusion, transient, asymptomatic decreases in left ventricular ejection fraction were observed using gated SPECT imaging; these resolved before the next infusion, 12 hours later.

SPORANOX<sup>®</sup> has been associated with reports of congestive heart failure. In post-marketing experience, heart failure was more frequently reported in patients receiving a total daily dose of 400 mg although there were also cases reported among those receiving lower total daily doses.

Calcium channel blockers can have negative inotropic effects which may be additive to those of itraconazole. In addition, itraconazole can inhibit the metabolism of calcium channel blockers. Therefore, caution should be used when co-administering itraconazole and calcium channel blockers due to an increased risk of CHF. Concomitant administration of SPORANOX<sup>®</sup> and felodipine or nisoldipine is contraindicated.

Cases of CHF, peripheral edema, and pulmonary edema have been reported in the post-marketing period among patients being treated for onychomycosis and/or systemic fungal infections. (See CLINICAL PHARMACOLOGY: Special Populations, CONTRAINDICATIONS, PRECAUTIONS: Drug Interactions, and ADVERSE REACTIONS: Post-marketing Experience for more information.)

**Interaction potential:**

SPORANOX<sup>®</sup> has a potential for clinically important drug interactions. Coadministration of specific drugs with itraconazole may result in changes in efficacy of itraconazole and/or the coadministered drug, life-threatening effects and/or sudden death. Drugs that are contraindicated, not recommended or recommended for use with caution in combination with itraconazole are listed in PRECAUTIONS: Drug Interactions.

**Interchangeability:**

SPORANOX<sup>®</sup> (itraconazole) Capsules and SPORANOX<sup>®</sup> Oral Solution should not be used interchangeably. This is because drug exposure is greater with the Oral Solution than with the Capsules when the same dose of drug is given. In addition, the topical effects of mucosal exposure may be different between the two formulations. Only the Oral Solution has been demonstrated effective for oral and/or esophageal candidiasis.

**PRECAUTIONS**

**General:**

SPORANOX<sup>®</sup> (itraconazole) Capsules should be administered after a full meal. (See CLINICAL PHARMACOLOGY: Pharmacokinetics and Metabolism).

Under fasted conditions, itraconazole absorption was decreased in the presence of decreased gastric acidity. The absorption of itraconazole may be decreased with the concomitant administration of antacids or gastric acid secretion suppressors. Studies conducted under fasted conditions demonstrated that administration with 8 ounces of a non-diet cola beverage resulted in increased absorption of itraconazole in AIDS patients with relative or absolute achlorhydria. This increase relative to the effects of a full meal is unknown. (See CLINICAL PHARMACOLOGY: Pharmacokinetics and Metabolism).

**Hepatotoxicity:**

Rare cases of serious hepatotoxicity have been observed with SPORANOX<sup>®</sup> treatment, including some cases within the first week. It is recommended that liver function monitoring be considered in all patients receiving SPORANOX<sup>®</sup>. Treatment should be stopped immediately and liver function testing should be conducted in patients who develop signs and symptoms suggestive of liver dysfunction.

**Neuropathy:**

If neuropathy occurs that may be attributable to SPORANOX<sup>®</sup> Capsules, the treatment should be discontinued.

**Cystic Fibrosis:**

If a cystic fibrosis patient does not respond to SPORANOX<sup>®</sup> Capsules, consideration should be given to switching to alternative therapy. For more information concerning the use of itraconazole in cystic fibrosis patients see the prescribing information for SPORANOX<sup>®</sup> Oral Solution.

**Hearing Loss:**

Transient or permanent hearing loss has been reported in patients receiving treatment with itraconazole. Several of these reports included concurrent administration of quinidine which is

contraindicated (see BOXED WARNING: Drug Interactions, CONTRAINDICATIONS: Drug Interactions and PRECAUTIONS: Drug Interactions). The hearing loss usually resolves when treatment is stopped, but can persist in some patients.

### Information for Patients:

- The topical effects of mucosal exposure may be different between the SPORANOX<sup>®</sup> Capsules and Oral Solution. Only the Oral Solution has been demonstrated effective for oral and/or esophageal candidiasis. SPORANOX<sup>®</sup> Capsules should not be used interchangeably with SPORANOX<sup>®</sup> Oral Solution.
- Instruct patients to take SPORANOX<sup>®</sup> Capsules with a full meal. SPORANOX<sup>®</sup> Capsules must be swallowed whole.
- Instruct patients about the signs and symptoms of congestive heart failure, and if these signs or symptoms occur during SPORANOX<sup>®</sup> administration, they should discontinue SPORANOX<sup>®</sup> and contact their healthcare provider immediately.
- Instruct patients to stop SPORANOX<sup>®</sup> treatment immediately and contact their healthcare provider if any signs and symptoms suggestive of liver dysfunction develop. Such signs and symptoms may include unusual fatigue, anorexia, nausea and/or vomiting, jaundice, dark urine, or pale stools.
- Instruct patients to contact their physician before taking any concomitant medications with itraconazole to ensure there are no potential drug interactions.
- Instruct patients that hearing loss can occur with the use of itraconazole. The hearing loss usually resolves when treatment is stopped, but can persist in some patients. Advise patients to discontinue therapy and inform their physicians if any hearing loss symptoms occur.
- Instruct patients that dizziness or blurred/double vision can sometimes occur with itraconazole. Advise patients that if they experience these events, they should not drive or use machines.

### Drug Interactions:

Itraconazole is mainly metabolized through CYP3A4. Other drugs that either share this metabolic pathway or modify CYP3A4 activity may influence the pharmacokinetics of itraconazole. Similarly, itraconazole may modify the pharmacokinetics of other drugs that share this metabolic pathway. Itraconazole is a potent CYP3A4 inhibitor and a P-glycoprotein inhibitor. When using concomitant medication, it is recommended that the corresponding label be consulted for information on the route of metabolism and the possible need to adjust dosages.

### Drugs that may decrease itraconazole plasma concentrations

Drugs that reduce the gastric acidity (e.g. acid neutralizing medicines such as aluminum hydroxide, or acid secretion suppressors such as H<sub>2</sub>-receptor antagonists and proton pump

inhibitors) impair the absorption of itraconazole from itraconazole capsules. It is recommended that these drugs be used with caution when coadministered with itraconazole capsules:

- It is recommended that itraconazole capsules be administered with an acidic beverage (such as non-diet cola) upon co-treatment with drugs reducing gastric acidity.
- It is recommended that acid neutralizing medicines (e.g. aluminum hydroxide) be administered at least 1 hour before or 2 hours after the intake of SPORANOX<sup>®</sup> Capsules.
- Upon coadministration, it is recommended that the antifungal activity be monitored and the itraconazole dose increased as deemed necessary.

Coadministration of itraconazole with potent enzyme inducers of CYP3A4 may decrease the bioavailability of itraconazole and hydroxy-itraconazole to such an extent that efficacy may be reduced. Examples include:

- Antibacterials: isoniazid, rifabutin (see also under ‘Drugs that may have their plasma concentrations increased by itraconazole’), rifampicin
- Anticonvulsants: carbamazepine, (see also under ‘Drugs that may have their plasma concentrations increased by itraconazole’), phenobarbital, phenytoin
- Antivirals: efavirenz, nevirapine

Therefore, administration of potent enzyme inducers of CYP3A4 with itraconazole is not recommended. It is recommended that the use of these drugs be avoided from 2 weeks before and during treatment with itraconazole, unless the benefits outweigh the risk of potentially reduced itraconazole efficacy. Upon coadministration, it is recommended that the antifungal activity be monitored and the itraconazole dose increased as deemed necessary.

#### Drugs that may increase itraconazole plasma concentrations

Potent inhibitors of CYP3A4 may increase the bioavailability of itraconazole. Examples include:

- Antibacterials: ciprofloxacin, clarithromycin, erythromycin
- Antivirals: ritonavir-boosted darunavir, ritonavir-boosted fosamprenavir, indinavir (see also under ‘Drugs that may have their plasma concentrations increased by itraconazole’), ritonavir (see also under ‘Drugs that may have their plasma concentrations increased by itraconazole’) and telaprevir.

It is recommended that these drugs be used with caution when coadministered with itraconazole capsules. It is recommended that patients who must take itraconazole concomitantly with potent inhibitors of CYP3A4 be monitored closely for signs or symptoms of increased or prolonged pharmacologic effects of itraconazole, and the itraconazole dose be decreased as deemed necessary.

### Drugs that may have their plasma concentrations increased by itraconazole

Itraconazole and its major metabolite, hydroxy-itraconazole, can inhibit the metabolism of drugs metabolized by CYP3A4 and can inhibit the drug transport by P-glycoprotein, which may result in increased plasma concentrations of these drugs and/or their active metabolite(s) when they are administered with itraconazole. These elevated plasma concentrations may increase or prolong both therapeutic and adverse effects of these drugs. CYP3A4-metabolized drugs known to prolong the QT interval may be contraindicated with itraconazole, since the combination may lead to ventricular tachyarrhythmias including occurrences of torsade de pointes, a potentially fatal arrhythmia. Once treatment is stopped, itraconazole plasma concentrations decrease to an almost undetectable concentration within 7 to 14 days, depending on the dose and duration of treatment. In patients with hepatic cirrhosis or in subjects receiving CYP3A4 inhibitors, the decline in plasma concentrations may be even more gradual. This is particularly important when initiating therapy with drugs whose metabolism is affected by itraconazole.

Examples of drugs that may have their plasma concentrations increased by itraconazole presented by drug class with advice regarding coadministration with itraconazole:

<b>Table 1: Drugs that may have their plasma concentrations increased by itraconazole</b>				
<b>Drug Class</b>	<b>Contraindicated</b>	<b>Not Recommended</b>	<b>Use with Caution</b>	<b>Comments</b>
	<i>Under no circumstances is the drug to be coadministered with itraconazole, and up to two weeks after discontinuation of treatment with itraconazole.</i>	<i>It is recommended that the use of the drug be avoided during and up to two weeks after discontinuation of treatment with itraconazole, unless the benefits outweigh the potentially increased risks of side effects. If coadministration cannot be avoided, clinical monitoring for signs or symptoms of increased or prolonged effects or side effects of the interacting drug is recommended, and its dosage be reduced or interrupted as deemed necessary. When appropriate, it is recommended that plasma concentrations be measured. The label of the coadministered drug should be consulted for information on dose adjustment and adverse effects.</i>	<i>Careful monitoring is recommended when the drug is coadministered with itraconazole. Upon coadministration, it is recommended that patients be monitored closely for signs or symptoms of increased or prolonged effects or side effects of the interacting drug, and its dosage be reduced as deemed necessary. When appropriate, it is recommended that plasma concentrations be measured. The label of the coadministered drug should be consulted for information on dose adjustment and adverse effects.</i>	
Alpha Blockers		tamsulosin		

<b>Table 1: Drugs that may have their plasma concentrations increased by itraconazole</b>				
<b>Drug Class</b>	<b>Contraindicated</b>	<b>Not Recommended</b>	<b>Use with Caution</b>	<b>Comments</b>
Analgesics	methadone		alfentanil, buprenorphine IV and sublingual, fentanyl, oxycodone, sufentanil	<p><b>Methadone:</b> The potential increase in plasma concentrations of methadone when coadministered with SPORANOX<sup>®</sup> may increase the risk of serious cardiovascular events including QTc prolongation and <i>torsade de pointes</i>.</p> <p><b>Fentanyl:</b> The potential increase in plasma concentrations of fentanyl when coadministered with SPORANOX<sup>®</sup> may increase the risk of potentially fatal respiratory depression.</p> <p><b>Sufentanil:</b> No human pharmacokinetic data of an interaction with itraconazole are available. <i>In vitro</i> data suggest that sufentanil is metabolized by CYP3A4 and so potentially increased sufentanil plasma concentrations would be expected when coadministered with SPORANOX<sup>®</sup>.</p>
Antiarrhythmics	disopyramide, dofetilide, dronedarone, quinidine		digoxin	<p><b>Disopyramide, dofetilide, dronedarone, quinidine:</b> The potential increase in plasma concentrations of these drugs when coadministered with SPORANOX<sup>®</sup> may increase the risk of serious cardiovascular events including QTc prolongation.</p>

<b>Table 1: Drugs that may have their plasma concentrations increased by itraconazole</b>				
<b>Drug Class</b>	<b>Contraindicated</b>	<b>Not Recommended</b>	<b>Use with Caution</b>	<b>Comments</b>
Antibacterials	telithromycin, in subjects with severe renal impairment or severe hepatic impairment	rifabutin	telithromycin	<p><b>Telithromycin:</b> The potential increase in plasma concentrations of telithromycin in subjects with severe renal impairment or severe hepatic impairment, when coadministered with SPORANOX® may increase the risk of serious cardiovascular events including QT prolongation and torsade de pointes.</p> <p><b>Rifabutin:</b> See also under ‘Drugs that may decrease itraconazole plasma concentrations’.</p>
Anticoagulants and Antiplatelet Drugs	ticagrelor	apixaban, rivaroxaban	coumarins, cilostazol, dabigatran	<p><b>Ticagrelor:</b> The potential increase in plasma concentrations of ticagrelor may increase the risk of bleeding.</p> <p><b>Coumarins:</b> SPORANOX® may enhance the anticoagulant effect of coumarin-like drugs, such as warfarin.</p>

<b>Table 1: Drugs that may have their plasma concentrations increased by itraconazole</b>				
<b>Drug Class</b>	<b>Contraindicated</b>	<b>Not Recommended</b>	<b>Use with Caution</b>	<b>Comments</b>
Anticonvulsants		carbamazepine		<b>Carbamazepine:</b> <i>In vivo</i> studies have demonstrated an increase in plasma carbamazepine concentrations in subjects concomitantly receiving ketoconazole. Although there are no data regarding the effect of itraconazole on carbamazepine metabolism, because of the similarities between ketoconazole and itraconazole, concomitant administration of SPORANOX <sup>®</sup> and carbamazepine may inhibit the metabolism of carbamazepine. See also under 'Drugs that may decrease itraconazole plasma concentrations'.
Antidiabetics			repaglinide, saxagliptin	
Anthelmintics and Antiprotozoals			praziquantel	
Antimigraine Drugs	ergot alkaloids, such as dihydroergotamine, ergometrine (ergonovine), ergotamine, methylergometrine (methylergonovine)		eletriptan	<b>Ergot Alkaloids:</b> The potential increase in plasma concentrations of ergot alkaloids when coadministered with SPORANOX <sup>®</sup> may increase the risk of ergotism, ie. a risk for vasospasm potentially leading to cerebral ischemia and/or ischemia of the extremities.

<b>Table 1: Drugs that may have their plasma concentrations increased by itraconazole</b>				
<b>Drug Class</b>	<b>Contraindicated</b>	<b>Not Recommended</b>	<b>Use with Caution</b>	<b>Comments</b>
Antineoplastics	irinotecan	axitinib, dabrafenib, dasatinib, ibrutinib, nilotinib, sunitinib	bortezomib, busulphan, docetaxel, erlotinib, imatinib, ixabepilone, lapatinib, ponatinib, trimetrexate, vinca alkaloids	<b>Irinotecan:</b> The potential increase in plasma concentrations of irinotecan when coadministered with SPORANOX <sup>®</sup> may increase the risk of potentially fatal adverse events.
Antipsychotics, Anxiolytics and Hypnotics	lurasidone, oral midazolam, pimozide, triazolam		alprazolam, aripiprazole, buspirone, diazepam, haloperidol, midazolam IV, perospirone, quetiapine, ramelteon, risperidone	<b>Midazolam, triazolam:</b> Coadministration of SPORANOX <sup>®</sup> and oral midazolam, or triazolam may cause several-fold increases in plasma concentrations of these drugs. This may potentiate and prolong hypnotic and sedative effects, especially with repeated dosing or chronic administration of these agents.  <b>Pimozide:</b> The potential increase in plasma concentrations of pimozide when coadministered with SPORANOX <sup>®</sup> may increase the risk of serious cardiovascular events including QTc prolongation and <i>torsade de pointes</i> .
Antivirals		simeprevir	maraviroc, indinavir, ritonavir, saquinavir	<b>Indinavir, ritonavir:</b> See also under 'Drugs that may increase itraconazole plasma concentrations'.
Beta Blockers			nadolol	

<b>Table 1: Drugs that may have their plasma concentrations increased by itraconazole</b>				
<b>Drug Class</b>	<b>Contraindicated</b>	<b>Not Recommended</b>	<b>Use with Caution</b>	<b>Comments</b>
Calcium Channel Blockers	felodipine, nisoldipine		other dihydropyridines, verapamil	<b>Calcium channel blockers</b> can have a negative inotropic effect which may be additive to those of itraconazole. The potential increase in plasma concentrations of calcium channel blockers when co-administered with SPORANOX <sup>®</sup> may increase the risk of congestive heart failure. <b>Dihydropyridines:</b> Concomitant administration of SPORANOX <sup>®</sup> may cause several-fold increases in plasma concentrations of dihydropyridines. Edema has been reported in patients concomitantly receiving SPORANOX <sup>®</sup> and dihydropyridine calcium channel blockers.
Cardiovascular Drugs, Miscellaneous	ranolazine	aliskiren, sildenafil, for the treatment of pulmonary hypertension	bosentan, riociguat	<b>Ranolazine:</b> The potential increase in plasma concentrations of ranolazine when coadministered with SPORANOX <sup>®</sup> may increase the risk of serious cardiovascular events including QTc prolongation.
Diuretics	eplerenone			<b>Eplerenone:</b> The potential increase in plasma concentrations of eplerenone when coadministered with SPORANOX <sup>®</sup> may increase the risk of hyperkalemia and hypotension.

<b>Table 1: Drugs that may have their plasma concentrations increased by itraconazole</b>				
<b>Drug Class</b>	<b>Contraindicated</b>	<b>Not Recommended</b>	<b>Use with Caution</b>	<b>Comments</b>
Gastrointestinal Drugs	cisapride		aprepitant	<b>Cisapride:</b> The potential increase in plasma concentrations of cisapride when coadministered with SPORANOX <sup>®</sup> may increase the risk of serious cardiovascular events including QTc prolongation.
Immunosuppressants		everolimus, temsirolimus	budesonide, ciclesonide, cyclosporine, dexamethasone, fluticasone, methylprednisolone, rapamycin (also known as sirolimus), tacrolimus	
Lipid Regulating Drugs	lovastatin, simvastatin		atorvastatin	The potential increase in plasma concentrations of atorvastatin, lovastatin, and simvastatin when coadministered with SPORANOX <sup>®</sup> may increase the risk of skeletal muscle toxicity, including rhabdomyolysis.
Respiratory Drugs		salmeterol		

<b>Table 1: Drugs that may have their plasma concentrations increased by itraconazole</b>				
<b>Drug Class</b>	<b>Contraindicated</b>	<b>Not Recommended</b>	<b>Use with Caution</b>	<b>Comments</b>
Urological Drugs	fesoterodine, in subjects with moderate to severe renal impairment, or moderate to severe hepatic impairment, solifenacin, in subjects with severe renal impairment or moderate to severe hepatic impairment	darifenacin, vardenafil	fesoterodine, oxybutynin, sildenafil, for the treatment of erectile dysfunction, solifenacin, tadalafil, tolterodine	<p><b>Fesoterodine:</b> The potential increase in plasma concentrations of the fesoterodine active metabolite may be greater in subjects with moderate to severe renal impairment, or moderate to severe hepatic impairment, which may lead to an increased risk of adverse reactions.</p> <p><b>Solifenacin:</b> The potential increase in plasma concentrations of solifenacin in subjects with severe renal impairment or moderate to severe hepatic impairment, when coadministered with SPORANOX<sup>®</sup> may increase the risk of serious cardiovascular events including QT prolongation.</p>

<b>Table 1: Drugs that may have their plasma concentrations increased by itraconazole</b>				
<b>Drug Class</b>	<b>Contraindicated</b>	<b>Not Recommended</b>	<b>Use with Caution</b>	<b>Comments</b>
Other	colchicine, in subjects with renal or hepatic impairment	colchicine, conivaptan, tolvaptan	cinacalcet	<p><b>Colchicine:</b> The potential increase in plasma concentrations of colchicine when coadministered with SPORANOX<sup>®</sup> may increase the risk of potentially fatal adverse events.</p> <p><b>Conivaptan and Tolvaptan:</b> A safe and effective dose of either conivaptan or tolvaptan has not been established when coadministered with SPORANOX<sup>®</sup>.</p>

## Drugs that may have their plasma concentrations decreased by itraconazole

Coadministration of itraconazole with the NSAID meloxicam may decrease the plasma concentration of meloxicam. It is recommended that meloxicam be used with caution when coadministered with itraconazole, and its effects or side effects be monitored. It is recommended that the dosage of meloxicam, if coadministered with itraconazole, be adjusted if necessary.

## Pediatric Population

Interaction studies have only been performed in adults.

## **Carcinogenesis, Mutagenesis, and Impairment of Fertility:**

Itraconazole showed no evidence of carcinogenicity potential in mice treated orally for 23 months at dosage levels up to 80 mg/kg/day (approximately 10x the maximum recommended human dose [MRHD]). Male rats treated with 25 mg/kg/day (3.1x MRHD) had a slightly increased incidence of soft tissue sarcoma. These sarcomas may have been a consequence of hypercholesterolemia, which is a response of rats, but not dogs or humans, to chronic itraconazole administration. Female rats treated with 50 mg/kg/day (6.25x MRHD) had an increased incidence of squamous cell carcinoma of the lung (2/50) as compared to the untreated group. Although the occurrence of squamous cell carcinoma in the lung is extremely uncommon in untreated rats, the increase in this study was not statistically significant.

Itraconazole produced no mutagenic effects when assayed in DNA repair test (unscheduled DNA synthesis) in primary rat hepatocytes, in Ames tests with *Salmonella typhimurium* (6 strains) and *Escherichia coli*, in the mouse lymphoma gene mutation tests, in a sex-linked recessive lethal mutation (*Drosophila melanogaster*) test, in chromosome aberration tests in human lymphocytes, in a cell transformation test with C3H/10T½ C18 mouse embryo fibroblasts cells, in a dominant lethal mutation test in male and female mice, and in micronucleus tests in mice and rats.

Itraconazole did not affect the fertility of male or female rats treated orally with dosage levels of up to 40 mg/kg/day (5x MRHD), even though parental toxicity was present at this dosage level. More severe signs of parental toxicity, including death, were present in the next higher dosage level, 160 mg/kg/day (20x MRHD).

## **Pregnancy: Teratogenic effects. Pregnancy Category C:**

Itraconazole was found to cause a dose-related increase in maternal toxicity, embryotoxicity, and teratogenicity in rats at dosage levels of approximately 40-160 mg/kg/day (5-20x MRHD), and in mice at dosage levels of approximately 80 mg/kg/day (10x MRHD). Itraconazole has been shown to cross the placenta in a rat model. In rats, the teratogenicity consisted of major skeletal defects; in mice, it consisted of encephaloceles and/or macroglossia.

There are no studies in pregnant women. SPORANOX<sup>®</sup> should be used for the treatment of systemic fungal infections in pregnancy only if the benefit outweighs the potential risk.

SPORANOX<sup>®</sup> should not be administered for the treatment of onychomycosis to pregnant patients or to women contemplating pregnancy. SPORANOX<sup>®</sup> should not be administered to women of childbearing potential for the treatment of onychomycosis unless they are using effective measures to prevent pregnancy and they begin therapy on the second or third day following the onset of menses. Effective contraception should be continued throughout SPORANOX<sup>®</sup> therapy and for 2 months following the end of treatment.

During post-marketing experience, cases of congenital abnormalities have been reported. (See ADVERSE REACTIONS: Post-marketing Experience.)

### **Nursing Mothers:**

Itraconazole is excreted in human milk; therefore, the expected benefits of SPORANOX<sup>®</sup> therapy for the mother should be weighed against the potential risk from exposure of itraconazole to the infant. The U.S. Public Health Service Centers for Disease Control and Prevention advises HIV-infected women not to breast-feed to avoid potential transmission of HIV to uninfected infants.

### **Pediatric Use:**

The efficacy and safety of SPORANOX<sup>®</sup> have not been established in pediatric patients.

The long-term effects of itraconazole on bone growth in children are unknown. In three toxicology studies using rats, itraconazole induced bone defects at dosage levels as low as 20 mg/kg/day (2.5x MRHD). The induced defects included reduced bone plate activity, thinning of the zona compacta of the large bones, and increased bone fragility. At a dosage level of 80 mg/kg/day (10x MRHD) over 1 year or 160 mg/kg/day (20x MRHD) for 6 months, itraconazole induced small tooth pulp with hypocellular appearance in some rats.

### **Geriatric Use:**

Clinical studies of SPORANOX<sup>®</sup> Capsules did not include sufficient numbers of subjects aged 65 years and over to determine whether they respond differently from younger subjects. It is advised to use SPORANOX<sup>®</sup> Capsules in these patients only if it is determined that the potential benefit outweighs the potential risks. In general, it is recommended that the dose selection for an elderly patient should be taken into consideration, reflecting the greater frequency of decreased hepatic, renal, or cardiac function, and of concomitant disease or other drug therapy.

Transient or permanent hearing loss has been reported in elderly patients receiving treatment with itraconazole. Several of these reports included concurrent administration of quinidine which is contraindicated (see **BOXED WARNING: Drug Interactions**, **CONTRAINDICATIONS: Drug Interactions** and **PRECAUTIONS: Drug Interactions**).

#### **HIV-Infected Patients:**

Because hypochlorhydria has been reported in HIV-infected individuals, the absorption of itraconazole in these patients may be decreased.

#### **Renal Impairment:**

Limited data are available on the use of oral itraconazole in patients with renal impairment. The exposure of itraconazole may be lower in some patients with renal impairment. Caution should be exercised when itraconazole is administered in this patient population and dose adjustment may be needed. (See **CLINICAL PHARMACOLOGY: Special Populations** and **DOSAGE AND ADMINISTRATION**.)

#### **Hepatic Impairment:**

Limited data are available on the use of oral itraconazole in patients with hepatic impairment. Caution should be exercised when this drug is administered in this patient population. It is recommended that patients with impaired hepatic function be carefully monitored when taking SPORANOX<sup>®</sup>. It is recommended that the prolonged elimination half-life of itraconazole observed in the single oral dose clinical trial with itraconazole capsules in cirrhotic patients be considered when deciding to initiate therapy with other medications metabolized by CYP3A4.

In patients with elevated or abnormal liver enzymes or active liver disease, or who have experienced liver toxicity with other drugs, treatment with SPORANOX<sup>®</sup> is strongly discouraged unless there is a serious or life-threatening situation where the expected benefit exceeds the risk. It is recommended that liver function monitoring be done in patients with pre-existing hepatic function abnormalities or those who have experienced liver toxicity with other medications. (See **CLINICAL PHARMACOLOGY: Special Populations** and **DOSAGE AND ADMINISTRATION**.)

#### **ADVERSE REACTIONS**

Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a drug cannot be directly compared to rates in the clinical trials of another drug and may not reflect the rates observed in clinical practice.

SPORANOX<sup>®</sup> has been associated with rare cases of serious hepatotoxicity, including liver failure and death. Some of these cases had neither pre-existing liver disease nor a serious underlying medical condition. If clinical signs or symptoms develop that are consistent with

liver disease, treatment should be discontinued and liver function testing performed. The risks and benefits of SPORANOX<sup>®</sup> use should be reassessed. (See WARNINGS: Hepatic Effects and PRECAUTIONS: Hepatotoxicity and Information for Patients.)

### Adverse Events in the Treatment of Systemic Fungal Infections

Adverse event data were derived from 602 patients treated for systemic fungal disease in U.S. clinical trials who were immunocompromised or receiving multiple concomitant medications. Treatment was discontinued in 10.5% of patients due to adverse events. The median duration before discontinuation of therapy was 81 days (range: 2 to 776 days). The table lists adverse events reported by at least 1% of patients.

**Table 2: Clinical Trials of Systemic Fungal Infections: Adverse Events Occurring with an Incidence of Greater than or Equal to 1%**

<b>Body System/Adverse Event</b>	<b>Incidence (%) (N=602)</b>
<b>Gastrointestinal</b>	
Nausea	11
Vomiting	5
Diarrhea	3
Abdominal Pain	2
Anorexia	1
<b>Body as a Whole</b>	
Edema	4
Fatigue	3
Fever	3
Malaise	1
<b>Skin and Appendages</b>	
Rash*	9
Pruritus	3
<b>Central/Peripheral Nervous System</b>	
Headache	4
Dizziness	2
<b>Psychiatric</b>	
Libido Decreased	1
Somnolence	1
<b>Cardiovascular</b>	
Hypertension	3
<b>Metabolic/Nutritional</b>	
Hypokalemia	2
<b>Urinary System</b>	
Albuminuria	1
<b>Liver and Biliary System</b>	
Hepatic Function Abnormal	3
<b>Reproductive System, Male</b>	
Impotence	1

**Table 2: Clinical Trials of Systemic Fungal Infections: Adverse Events Occurring with an Incidence of Greater than or Equal to 1%**

Body System/Adverse Event	Incidence (%) (N=602)
---------------------------	-----------------------

\* Rash tends to occur more frequently in immunocompromised patients receiving immunosuppressive medications.

Adverse events infrequently reported in all studies included constipation, gastritis, depression, insomnia, tinnitus, menstrual disorder, adrenal insufficiency, gynecomastia, and male breast pain.

### Adverse Events Reported in Toenail Onychomycosis Clinical Trials

Patients in these trials were on a continuous dosing regimen of 200 mg once daily for 12 consecutive weeks.

The following adverse events led to temporary or permanent discontinuation of therapy.

**Table 3: Clinical Trials of Onychomycosis of the Toenail: Adverse Events Leading to Temporary or Permanent Discontinuation of Therapy**

Adverse Event	Incidence (%) Itraconazole (N=112)
Elevated Liver Enzymes (greater than twice the upper limit of normal)	4
Gastrointestinal Disorders	4
Rash	3
Hypertension	2
Orthostatic Hypotension	1
Headache	1
Malaise	1
Myalgia	1
Vasculitis	1
Vertigo	1

The following adverse events occurred with an incidence of greater than or equal to 1% (N=112): headache: 10%; rhinitis: 9%; upper respiratory tract infection: 8%; sinusitis, injury: 7%; diarrhea, dyspepsia, flatulence, abdominal pain, dizziness, rash: 4%; cystitis, urinary tract infection, liver function abnormality, myalgia, nausea: 3%; appetite increased, constipation, gastritis, gastroenteritis, pharyngitis, asthenia, fever, pain, tremor, herpes zoster, abnormal dreaming: 2%.

### Adverse Events Reported in Fingernail Onychomycosis Clinical Trials

Patients in these trials were on a pulse regimen consisting of two 1-week treatment periods of 200 mg twice daily, separated by a 3-week period without drug.

The following adverse events led to temporary or permanent discontinuation of therapy.

**Table 4: Clinical Trials of Onychomycosis of the Fingernail: Adverse Events Leading to Temporary or Permanent Discontinuation of Therapy**

Adverse Event	Incidence (%) Itraconazole (N=37)
Rash/Pruritus	3
Hypertriglyceridemia	3

The following adverse events occurred with an incidence of greater than or equal to 1% (N=37): headache: 8%; pruritus, nausea, rhinitis: 5%; rash, bursitis, anxiety, depression, constipation, abdominal pain, dyspepsia, ulcerative stomatitis, gingivitis, hypertriglyceridemia, sinusitis, fatigue, malaise, pain, injury: 3%.

#### **Adverse Events Reported from Other Clinical Trials**

In addition, the following adverse drug reaction was reported in patients who participated in SPORANOX<sup>®</sup> Capsules clinical trials: *Hepatobiliary Disorders*: hyperbilirubinemia.

The following is a list of additional adverse drug reactions associated with itraconazole that have been reported in clinical trials of SPORANOX<sup>®</sup> Oral Solution and itraconazole IV excluding the adverse reaction term “Injection site inflammation” which is specific to the injection route of administration:

*Cardiac Disorders*: cardiac failure, left ventricular failure, tachycardia;

*General Disorders and Administration Site Conditions*: face edema, chest pain, chills;

*Hepatobiliary Disorders*: hepatic failure, jaundice;

*Investigations*: alanine aminotransferase increased, aspartate aminotransferase increased, blood alkaline phosphatase increased, blood lactate dehydrogenase increased, blood urea increased, gamma-glutamyltransferase increased, urine analysis abnormal;

*Metabolism and Nutrition Disorders*: hyperglycemia, hyperkalemia, hypomagnesemia;

*Psychiatric Disorders*: confusional state;

*Renal and Urinary Disorders*: renal impairment;

*Respiratory, Thoracic and Mediastinal Disorders*: dysphonia, cough;

*Skin and Subcutaneous Tissue Disorders*: rash erythematous, hyperhidrosis;

*Vascular Disorders:* hypotension

### Post-marketing Experience

Adverse drug reactions that have been first identified during post-marketing experience with SPORANOX<sup>®</sup> (all formulations) are listed in the table below. Because these reactions are reported voluntarily from a population of uncertain size, reliably estimating their frequency or establishing a causal relationship to drug exposure is not always possible.

**Table 5: Postmarketing Reports of Adverse Drug Reactions**

<b>Blood and Lymphatic System Disorders:</b>	Leukopenia, neutropenia, thrombocytopenia
<b>Immune System Disorders:</b>	Anaphylaxis; anaphylactic, anaphylactoid and allergic reactions; serum sickness; angioneurotic edema
<b>Nervous System Disorders:</b>	Peripheral neuropathy, paresthesia, hypoesthesia, tremor
<b>Eye Disorders:</b>	Visual disturbances, including vision blurred and diplopia
<b>Ear and Labyrinth Disorders:</b>	Transient or permanent hearing loss
<b>Cardiac Disorders:</b>	Congestive heart failure
<b>Respiratory, Thoracic and Mediastinal Disorders:</b>	Pulmonary edema, dyspnea
<b>Gastrointestinal Disorders:</b>	Pancreatitis, dysgeusia
<b>Hepatobiliary Disorders:</b>	Serious hepatotoxicity (including some cases of fatal acute liver failure), hepatitis
<b>Skin and Subcutaneous Tissue Disorders:</b>	Toxic epidermal necrolysis, Stevens-Johnson syndrome, acute generalized exanthematous pustulosis, erythema multiforme, exfoliative dermatitis, leukocytoclastic vasculitis, alopecia, photosensitivity, urticaria
<b>Musculoskeletal and Connective Tissue Disorders:</b>	Arthralgia
<b>Renal and Urinary Disorders:</b>	Urinary incontinence, pollakiuria
<b>Reproductive System and Breast Disorders:</b>	Erectile dysfunction
<b>General Disorders and Administration Site</b>	Peripheral edema

**Table 5: Postmarketing Reports of Adverse Drug Reactions**

**Conditions:**

**Investigations:** Blood creatine phosphokinase increased

---

There is limited information on the use of SPORANOX<sup>®</sup> during pregnancy. Cases of congenital abnormalities including skeletal, genitourinary tract, cardiovascular and ophthalmic malformations as well as chromosomal and multiple malformations have been reported during post-marketing experience. A causal relationship with SPORANOX<sup>®</sup> has not been established. (See CLINICAL PHARMACOLOGY: Special Populations, CONTRAINDICATIONS, WARNINGS, and PRECAUTIONS: Drug Interactions for more information.)

**OVERDOSAGE**

Itraconazole is not removed by dialysis. In the event of accidental overdose, supportive measures should be employed. Activated charcoal may be given if considered appropriate. In general, adverse events reported with overdose have been consistent with adverse drug reactions already listed in this package insert for itraconazole. (See ADVERSE REACTIONS.)

**DOSAGE AND ADMINISTRATION**

SPORANOX<sup>®</sup> (itraconazole) Capsules should be taken with a full meal to ensure maximal absorption. SPORANOX<sup>®</sup> (itraconazole) Capsules must be swallowed whole.

SPORANOX<sup>®</sup> Capsules is a different preparation than SPORANOX<sup>®</sup> Oral Solution and should not be used interchangeably.

**Treatment of Blastomycosis and Histoplasmosis:**

The recommended dose is 200 mg once daily (2 capsules). If there is no obvious improvement, or there is evidence of progressive fungal disease, the dose should be increased in 100-mg increments to a maximum of 400 mg daily. Doses above 200 mg/day should be given in two divided doses.

**Treatment of Aspergillosis:**

A daily dose of 200 to 400 mg is recommended.

**Treatment in Life-Threatening Situations:**

In life-threatening situations, a loading dose should be used.

Although clinical studies did not provide for a loading dose, it is recommended, based on pharmacokinetic data, that a loading dose of 200 mg (2 capsules) three times daily (600 mg/day) be given for the first 3 days of treatment.

Treatment should be continued for a minimum of three months and until clinical parameters and laboratory tests indicate that the active fungal infection has subsided. An inadequate period of treatment may lead to recurrence of active infection.

SPORANOX<sup>®</sup> Capsules and SPORANOX<sup>®</sup> Oral Solution should not be used interchangeably. Only the oral solution has been demonstrated effective for oral and/or esophageal candidiasis.

**Treatment of Onychomycosis:**

Toenails with or without fingernail involvement: The recommended dose is 200 mg (2 capsules) once daily for 12 consecutive weeks.

**Treatment of Onychomycosis:**

Fingernails only: The recommended dosing regimen is 2 treatment pulses, each consisting of 200 mg (2 capsules) b.i.d. (400 mg/day) for 1 week. The pulses are separated by a 3-week period without SPORANOX<sup>®</sup>.

**Use in Patients with Renal Impairment:**

Limited data are available on the use of oral itraconazole in patients with renal impairment. Caution should be exercised when this drug is administered in this patient population. (See CLINICAL PHARMACOLOGY: Special Populations and PRECAUTIONS.)

**Use in Patients with Hepatic Impairment:**

Limited data are available on the use of oral itraconazole in patients with hepatic impairment. Caution should be exercised when this drug is administered in this patient population. (See CLINICAL PHARMACOLOGY: Special Populations, WARNINGS, and PRECAUTIONS.)

**HOW SUPPLIED**

SPORANOX<sup>®</sup> (itraconazole) Capsules are available containing 100 mg of itraconazole, with a blue opaque cap and pink transparent body, imprinted with “JANSSEN” and “SPORANOX 100.” The capsules are supplied in unit-dose blister packs of 3 × 10 capsules (NDC 50458-290-01), bottles of 30 capsules (NDC 50458-290-04) and in the PulsePak<sup>®</sup> containing 7 blister packs × 4 capsules each (NDC 50458-290-28).

Store at controlled room temperature 15°-25°C (59°-77°F). Protect from light and moisture.

Keep out of reach of children.

© Janssen Pharmaceuticals, Inc 2001

Revised: 04/2015

Product of Ireland

Capsule contents manufactured by:

Janssen Pharmaceutica, NV.

Olen, Belgium

Manufactured by:

JOLLC, Gurabo, Puerto Rico 00778

Manufactured for:

Janssen Pharmaceuticals, Inc.

Titusville, NJ 08560

## PATIENT INFORMATION

### 100 mg SPORANOX<sup>®</sup> (itraconazole) Capsules

**This summary contains important information about SPORANOX<sup>®</sup> (SPOR-ah-nox).** This information is for patients who have been prescribed SPORANOX<sup>®</sup> to treat fungal nail infections. **If your doctor prescribed SPORANOX<sup>®</sup> for medical problems other than fungal nail infections, ask your doctor if there is any information in this summary that does not apply to you.** Read this information carefully each time you start to use SPORANOX<sup>®</sup>. This information does not take the place of discussion between you and your doctor. Only your doctor can decide if SPORANOX<sup>®</sup> is the right treatment for you. If you do not understand some of this information or have any questions, talk with your doctor or pharmacist.

#### **WHAT IS THE MOST IMPORTANT INFORMATION I SHOULD KNOW ABOUT SPORANOX<sup>®</sup>?**

SPORANOX<sup>®</sup> is used to treat fungal nail infections. However, SPORANOX<sup>®</sup> is not for everyone. **Do not take SPORANOX<sup>®</sup> for fungal nail infections if you have had heart failure, including congestive heart failure. You should not take SPORANOX<sup>®</sup> if you are taking certain medicines that could lead to serious or life-threatening medical problems.** (See “Who Should Not Take SPORANOX<sup>®</sup>?” below.)

If you have had heart, lung, liver, kidney or other serious health problems, ask your doctor if it is safe for you to take SPORANOX<sup>®</sup>.

#### **WHAT HAPPENS IF I HAVE A FUNGAL NAIL INFECTION?**

Anyone can have a fungal nail infection, but it is usually found in adults. When a fungus infects the tip or sides of a nail, the infected part of the nail may turn yellow or brown. If not treated, the fungus may spread under the nail towards the cuticle. If the fungus spreads, more of the nail may change color, may become thick or brittle, and the tip of the nail may become raised. In some patients, this can cause pain and discomfort.

#### **WHAT IS SPORANOX<sup>®</sup>?**

SPORANOX<sup>®</sup> is a prescription medicine used to treat fungal infections of the toenails and fingernails. It is also used to treat some types of fungal infections in other areas of your body. We do not know if SPORANOX<sup>®</sup> works in children with fungal nail infections or if it is safe for children to take.

SPORANOX<sup>®</sup> comes in the form of capsules and liquid (oral solution). The capsule and liquid forms work differently, so you should not use one in place of the other. This Patient Information discusses only the capsule form of SPORANOX<sup>®</sup>. You will get these capsules in a medicine bottle or a SPORANOX *PulsePak*<sup>®</sup>. The *PulsePak*<sup>®</sup> contains 28 capsules for treatment of your fungal nail infection.

SPORANOX<sup>®</sup> goes into your bloodstream and travels to the source of the infection underneath the nail so that it can fight the infection there. Improved nails may not be obvious for several months after the treatment period is finished because it usually takes about 6 months to grow a new fingernail and 12 months to grow a new toenail.

### **WHO SHOULD NOT TAKE SPORANOX<sup>®</sup>?**

SPORANOX<sup>®</sup> is not for everyone. Your doctor will decide if SPORANOX<sup>®</sup> is the right treatment for you. **Some patients should not take SPORANOX<sup>®</sup> because they may have certain health problems or may be taking certain medicines that could lead to serious or life-threatening medical problems.**

Tell your doctor and pharmacist the name of all the prescription and non-prescription medicines you are taking, including dietary supplements and herbal remedies. Also tell your doctor about any other medical conditions you have had, especially heart, lung, liver or kidney conditions; or if you have cystic fibrosis, or have had an allergic reaction to SPORANOX<sup>®</sup> or any other antifungal medicines.

**Never** take SPORANOX<sup>®</sup> if you:

- have had heart failure, including congestive heart failure.
- are taking any of the medicines listed below. Dangerous or even life-threatening side effects could result:
  - cisapride (such as Propulsid<sup>®</sup>)
  - colchicine (such as Colcris<sup>™</sup>) [if you also have pre-existing kidney or liver impairment]
  - disopyramide (such as Norpace<sup>®</sup>)
  - dofetilide (such as Tikosyn<sup>™</sup>)
  - dronedarone (such as Multaq<sup>®</sup>)
  - eplerenone (such as Inspra<sup>®</sup>)
  - ergot alkaloids (such as Migranal<sup>®</sup>, Ergonovine, Cafergot<sup>®</sup>, Methergine<sup>®</sup>)
  - felodipine (such as Plendil<sup>®</sup>)

- fesoterodine (such as Toviaz<sup>®</sup>) [if you also have pre-existing kidney or liver impairment]
  - irinotecan (such as Camptosar<sup>®</sup>)
  - lovastatin (such as Mevacor<sup>®</sup>, Advicor<sup>®</sup>, Altacor<sup>™</sup>)
  - lurasidone (such as Latuda<sup>®</sup>)
  - methadone (such as Dolophine<sup>®</sup>)
  - midazolam (such as Versed<sup>®</sup>)
  - nisoldipine (such as Sular<sup>®</sup>)
  - pimozide (such as Orap<sup>®</sup>)
  - quinidine (such as Cardioquin<sup>®</sup>, Quinaglute<sup>®</sup>, Quinidex<sup>®</sup>)
  - ranolazine (such as Ranexa<sup>®</sup>)
  - simvastatin (such as Zocor<sup>®</sup>)
  - solifenacin (such as Vesicare<sup>®</sup>) [if you also have pre-existing kidney or liver impairment]
  - telithromycin (such as Ketek<sup>®</sup>) [if you also have pre-existing kidney or liver impairment]
  - ticagrelor (such as Brilinta<sup>®</sup>)
  - triazolam (such as Hacion<sup>®</sup>)
- have ever had an allergic reaction to itraconazole.

Taking SPORANOX<sup>®</sup> with certain other medicines may lead to serious or life-threatening medical problems. Tell your doctor and pharmacist the name of all the prescription and non-prescription medicines you are taking, including dietary supplements and herbal remedies. Your doctor will decide if SPORANOX<sup>®</sup> is the right treatment for you.

### **WHAT SHOULD I KNOW ABOUT SPORANOX<sup>®</sup> AND PREGNANCY OR BREAST FEEDING?**

Never take SPORANOX<sup>®</sup> if you have a fungal nail infection and are pregnant or planning to become pregnant within 2 months after you have finished your treatment.

If you are able to become pregnant, you should use effective birth control during SPORANOX<sup>®</sup> treatment and for 2 months after finishing treatment. Ask your doctor about effective types of birth control.

If you are breast-feeding, talk with your doctor about whether you should take SPORANOX<sup>®</sup>.

## HOW SHOULD I TAKE SPORANOX<sup>®</sup>?

Always take SPORANOX<sup>®</sup> Capsules during or right after a full meal.

Your doctor will decide the right dose for you. Depending on your infection, you will take SPORANOX<sup>®</sup> once a day for 12 weeks, or twice a day for 1 week in a “pulse” dosing schedule. You will receive either a bottle of capsules or a *PulsePak*<sup>®</sup>. Do not skip any doses. Be sure to finish all your SPORANOX<sup>®</sup> as prescribed by your doctor.

If you have ever had liver problems, your doctor should do a blood test to check your condition. If you haven’t had liver problems, your doctor may recommend blood tests to check the condition of your liver because patients taking SPORANOX<sup>®</sup> can develop liver problems.

SPORANOX<sup>®</sup> can sometimes cause dizziness or blurred/double vision. If you have these symptoms, do not drive or use machines.

If you forget to take or miss doses of SPORANOX<sup>®</sup>, ask your doctor what you should do with the missed doses.

## THE SPORANOX *PulsePak*<sup>®</sup>

If you use the *PulsePak*<sup>®</sup>, you will take SPORANOX<sup>®</sup> for 1 week and then take no SPORANOX<sup>®</sup> for the next 3 weeks before repeating the 1-week treatment. This is called “pulse dosing.” The SPORANOX *PulsePak*<sup>®</sup> contains enough medicine for one “pulse” (1 week of treatment).

The SPORANOX *PulsePak*<sup>®</sup> comes with special instructions. It contains 7 pouches—one for each day of treatment. Inside each pouch is a card containing 4 capsules. Looking at the back of the card, fold it back along the dashed line and peel away the backing so that you can remove 2 capsules.

- Take 2 capsules in the morning and 2 capsules in the evening. This means you will take 4 capsules a day for 7 days. At the end of 7 days, you will have taken all of the capsules in the *PulsePak*<sup>®</sup> box.
- After you finish the *PulsePak*<sup>®</sup>, do not take any SPORANOX<sup>®</sup> for the next 3 weeks. Even though you are not taking any capsules during this time, SPORANOX<sup>®</sup> keeps working inside your nails to help fight the fungal infection.
- You will need more than one “pulse” to treat your fungal nail infection. When your doctor prescribes another pulse treatment, be sure to get your refill before the end of week 4.

SPORANOX <sup>®</sup> Pulse Dosing														
Take 2 SPORANOX <sup>®</sup> capsules twice a day for 1 week														
	Day 1		Day 2		Day 3		Day 4		Day 5		Day 6		Day 7	
	AM	PM	AM	PM	AM	PM	AM	PM	AM	PM	AM	PM	AM	PM
<b>Week 1</b>	//	//	//	//	//	//	//	//	//	//	//	//	//	//
<b>Week 2</b>	<p><b>For the next 3 weeks, do not take any SPORANOX<sup>®</sup> capsules.</b></p> <p>Remember to get a refill before the end of Week 4 when your doctor prescribes another <i>PulsePak</i><sup>®</sup>.</p>													
<b>Week 3</b>														
<b>Week 4</b>														

### WHAT ARE THE POSSIBLE SIDE EFFECTS OF SPORANOX<sup>®</sup>?

The most common side effects include: headache, and digestive system problems (such as nausea, and abdominal pain).

**Stop SPORANOX<sup>®</sup> and call your doctor or get medical assistance right away** if you have a severe allergic reaction. Symptoms of an allergic reaction may include skin rash, itching, hives, shortness of breath or difficulty breathing, and/or swelling of the face. Very rarely, an oversensitivity to sunlight, a tingling sensation in the limbs or a severe skin disorder can occur. If any of these symptoms occur, stop taking SPORANOX<sup>®</sup> and contact your doctor.

**Stop SPORANOX<sup>®</sup> and call your doctor right away** if you develop shortness of breath; have unusual swelling of your feet, ankles or legs; suddenly gain weight; are unusually tired; cough up white or pink phlegm; have unusual fast heartbeats; or begin to wake up at night. In rare cases, patients taking SPORANOX<sup>®</sup> could develop serious heart problems, and these could be warning signs of heart failure.

**Stop SPORANOX<sup>®</sup> and call your doctor right away** if you become unusually tired; lose your appetite; or develop nausea, abdominal pain, or vomiting, a yellow color to your skin or eyes, or dark colored urine or pale stools (bowel movements). In rare cases, patients taking SPORANOX<sup>®</sup> could develop serious liver problems and these could be warning signs.

**Stop SPORANOX<sup>®</sup> and call your doctor right away** if you experience any hearing loss symptoms. In very rare cases, patients taking SPORANOX<sup>®</sup> have reported temporary or permanent hearing loss.

**Call your doctor right away** if you develop tingling or numbness in your extremities (hands or feet), if your vision gets blurry or you see double, if you hear a ringing in your ears, if you lose the ability to control your urine or urinate much more than usual.

Additional possible side effects include upset stomach, vomiting, constipation, fever, inflammation of the pancreas, menstrual disorder, erectile dysfunction, dizziness, muscle pain, painful joints, unpleasant taste, or hair loss. These are not all the side effects of SPORANOX<sup>®</sup>. Your doctor or pharmacist can give you a more complete list.

### **WHAT SHOULD I DO IF I TAKE AN OVERDOSE OF SPORANOX<sup>®</sup>?**

If you think you took too much SPORANOX<sup>®</sup>, call your doctor or local poison control center, or go to the nearest hospital emergency room right away.

### **HOW SHOULD I STORE SPORANOX<sup>®</sup>?**

Keep all medicines, including SPORANOX<sup>®</sup>, out of the reach of children.

Store SPORANOX<sup>®</sup> Capsules and the *PulsePak*<sup>®</sup> at room temperature in a dry place away from light.

### **GENERAL ADVICE ABOUT SPORANOX<sup>®</sup>**

Medicines are sometimes prescribed for conditions that are not mentioned in patient information leaflets. Do not use SPORANOX<sup>®</sup> for a condition for which it was not prescribed. Do not give SPORANOX<sup>®</sup> to other people, even if they have the same symptoms you have. It may harm them.

This leaflet summarizes the most important information about SPORANOX<sup>®</sup>. If you would like more information, talk with your doctor. You can ask your doctor or pharmacist for information about SPORANOX<sup>®</sup> that is written for health professionals or you can call 1-800-526-7736.

This patient information has been approved by the U.S. Food and Drug Administration.

Product of Ireland

© Janssen Pharmaceuticals, Inc 2001

Printed in USA/ Revised: 04/2015

**SPORANOX<sup>®</sup>**  
**(itraconazole)**  
**Oral Solution**


**BOXED WARNING**

**Congestive Heart Failure, Cardiac Effects and Drug Interactions:** If signs or symptoms of congestive heart failure occur during administration of SPORANOX<sup>®</sup> (itraconazole) Oral Solution, continued SPORANOX<sup>®</sup> use should be reassessed. When itraconazole was administered intravenously to dogs and healthy human volunteers, negative inotropic effects were seen. (See CONTRAINDICATIONS, WARNINGS, PRECAUTIONS: Drug Interactions, ADVERSE REACTIONS: Post-marketing Experience, and CLINICAL PHARMACOLOGY: Special Populations for more information.)

**Drug Interactions: Coadministration of the following drugs are contraindicated with SPORANOX<sup>®</sup> Oral Solution:** methadone, disopyramide, dofetilide, dronedarone, quinidine, ergot alkaloids (such as dihydroergotamine, ergometrine (ergonovine), ergotamine, methylergometrine (methylergonovine)), irinotecan, lurasidone, oral midazolam, pimozide, triazolam, felodipine, nisoldipine, ranolazine, eplerenone, cisapride, lovastatin, simvastatin, ticagrelor, and, in subjects with varying degrees of renal or hepatic impairment, colchicine, fesoterodine, telithromycin and solifenacin. See PRECAUTIONS: Drug Interactions Section for specific examples. Coadministration with itraconazole can cause elevated plasma concentrations of these drugs and may increase or prolong both the pharmacologic effects and/or adverse reactions to these drugs. For example, increased plasma concentrations of some of these drugs can lead to QT prolongation and ventricular tachyarrhythmias including occurrences of torsades de pointes, a potentially fatal arrhythmia. See CONTRAINDICATIONS and WARNINGS Sections, and PRECAUTIONS: Drug Interactions Section for specific examples.

**DESCRIPTION**

SPORANOX<sup>®</sup> is the brand name for itraconazole, an azole antifungal agent. Itraconazole is a 1:1:1:1 racemic mixture of four diastereomers (two enantiomeric pairs), each possessing three chiral centers. It may be represented by the following structural formula and nomenclature:



(±)-1-[(R\*)-sec-butyl]-4-[p-[4-[p-[(2R\*,4S\*)-2-(2,4-dichlorophenyl)-2-(1H-1,2,4-triazol-1-ylmethyl)-1,3-dioxolan-4-yl]methoxy]phenyl]-1-piperazinyl]phenyl]-Δ<sup>2</sup>-1,2,4-triazolin-5-one mixture with (±)-1-[(R\*)-sec-butyl]-4-[p-[4-[p-[(2S\*,4R\*)-2-(2,4-dichlorophenyl)-2-(1H-1,2,4-triazol-1-ylmethyl)-1,3-dioxolan-4-yl]methoxy]phenyl]-1-piperazinyl]phenyl]-Δ<sup>2</sup>-1,2,4-triazolin-5-one

or

(±)-1-[(RS)-sec-butyl]-4-[p-[4-[p-[(2R,4S)-2-(2,4-dichlorophenyl)-2-(1H-1,2,4-triazol-1-ylmethyl)-1,3-dioxolan-4-yl]methoxy]phenyl]-1-piperazinyl]phenyl]-Δ<sup>2</sup>-1,2,4-triazolin-5-one.

Itraconazole has a molecular formula of C<sub>35</sub>H<sub>38</sub>Cl<sub>2</sub>N<sub>8</sub>O<sub>4</sub> and a molecular weight of 705.64. It is a white to slightly yellowish powder. It is insoluble in water, very slightly soluble in alcohols, and freely soluble in dichloromethane. It has a pKa of 3.70 (based on extrapolation of values obtained from methanolic solutions) and a log (n-octanol/water) partition coefficient of 5.66 at pH 8.1.

SPORANOX<sup>®</sup> (itraconazole) Oral Solution contains 10 mg of itraconazole per mL, solubilized by hydroxypropyl-β-cyclodextrin (400 mg/mL) as a molecular inclusion complex. SPORANOX<sup>®</sup> Oral Solution is clear and yellowish in color with a target pH of 2. Other ingredients are hydrochloric acid, propylene glycol, purified water, sodium hydroxide, sodium saccharin, sorbitol, cherry flavor 1, cherry flavor 2 and caramel flavor.

## CLINICAL PHARMACOLOGY

### Pharmacokinetics and Metabolism:

Itraconazole

#### *General Pharmacokinetic Characteristics*

Peak plasma concentrations are reached within 2.5 hours following administration of the oral solution. As a consequence of non-linear pharmacokinetics, itraconazole accumulates in plasma during multiple dosing. Steady-state concentrations are generally reached within about 15 days, with C<sub>max</sub> and AUC values 4 to 7-fold higher than those seen after a single dose. Steady-state C<sub>max</sub> values of about 2 µg/ml are reached after oral administration of 200 mg once daily. The terminal half-life of itraconazole generally ranges from 16 to 28 hours after single dose and increases to 34 to 42 hours with repeated dosing. Once treatment is stopped, itraconazole plasma concentrations decrease to an almost undetectable concentration within 7 to 14 days, depending on the dose and duration of treatment. Itraconazole mean total plasma clearance following intravenous administration is 278 ml/min. Itraconazole clearance decreases at higher doses due to saturable hepatic metabolism.

### *Absorption*

Itraconazole is rapidly absorbed after administration of the oral solution. Peak plasma concentrations of itraconazole are reached within 2.5 hours following administration of the oral solution under fasting conditions. The observed absolute bioavailability of itraconazole under fed conditions is about 55% and increases by 30% when the oral solution is taken in fasting conditions. Itraconazole exposure is greater with the oral solution than with the capsule formulation when the same dose of drug is given. (see WARNINGS)

### *Distribution*

Most of the itraconazole in plasma is bound to protein (99.8%), with albumin being the main binding component (99.6% for the hydroxy-metabolite). It has also a marked affinity for lipids. Only 0.2% of the itraconazole in plasma is present as free drug. Itraconazole is distributed in a large apparent volume in the body (>700 L), suggesting extensive distribution into tissues. Concentrations in lung, kidney, liver, bone, stomach, spleen and muscle were found to be two to three times higher than corresponding concentrations in plasma, and the uptake into keratinous tissues, skin in particular, up to four times higher. Concentrations in the cerebrospinal fluid are much lower than in plasma.

### *Metabolism*

Itraconazole is extensively metabolized by the liver into a large number of metabolites. *In vitro* studies have shown that CYP3A4 is the major enzyme involved in the metabolism of itraconazole. The main metabolite is hydroxy-itraconazole, which has *in vitro* antifungal activity comparable to itraconazole; trough plasma concentrations of this metabolite are about twice those of itraconazole.

### *Excretion*

Itraconazole is excreted mainly as inactive metabolites in urine (35%) and in feces (54%) within one week of an oral solution dose. Renal excretion of itraconazole and the active metabolite hydroxy-itraconazole account for less than 1% of an intravenous dose. Based on an oral radiolabeled dose, fecal excretion of unchanged drug ranges from 3% to 18% of the dose.

### **Special Populations:**

#### **Renal Impairment:**

Limited data are available on the use of oral itraconazole in patients with renal impairment. A pharmacokinetic study using a single 200-mg oral dose of itraconazole was conducted in three groups of patients with renal impairment (uremia: n=7; hemodialysis: n=7; and continuous ambulatory peritoneal dialysis: n=5). In uremic subjects with a mean creatinine clearance of 13 mL/min.  $\times$  1.73 m<sup>2</sup>, the exposure, based on AUC, was slightly reduced compared with normal population parameters. This study did not demonstrate any significant effect of hemodialysis or

continuous ambulatory peritoneal dialysis on the pharmacokinetics of itraconazole ( $T_{max}$ ,  $C_{max}$ , and  $AUC_{0-8h}$ ). Plasma concentration-versus-time profiles showed wide intersubject variation in all three groups.

After a single intravenous dose, the mean terminal half-lives of itraconazole in patients with mild (defined in this study as  $CrCl$  50-79 ml/min), moderate (defined in this study as  $CrCl$  20-49 ml/min), and severe renal impairment (defined in this study as  $CrCl$  <20 ml/min) were similar to that in healthy subjects (range of means 42-49 hours vs 48 hours in renally impaired patients and healthy subjects, respectively). Overall exposure to itraconazole, based on AUC, was decreased in patients with moderate and severe renal impairment by approximately 30% and 40%, respectively, as compared with subjects with normal renal function.

Data are not available in renally impaired patients during long-term use of itraconazole. Dialysis has no effect on the half-life or clearance of itraconazole or hydroxy-itraconazole. (See PRECAUTIONS and DOSAGE AND ADMINISTRATION.)

#### **Hepatic Impairment:**

Itraconazole is predominantly metabolized in the liver. A pharmacokinetic study was conducted in 6 healthy and 12 cirrhotic subjects who were administered a single 100-mg dose of itraconazole as capsule. A statistically significant reduction in mean  $C_{max}$  (47%) and a twofold increase in the elimination half-life ( $37 \pm 17$  hours vs.  $16 \pm 5$  hours) of itraconazole were noted in cirrhotic subjects compared with healthy subjects. However, overall exposure to itraconazole, based on AUC, was similar in cirrhotic patients and in healthy subjects. Data are not available in cirrhotic patients during long-term use of itraconazole. (See CONTRAINDICATIONS, PRECAUTIONS: Drug Interactions and DOSAGE AND ADMINISTRATION.)

#### **Decreased Cardiac Contractility:**

When itraconazole was administered intravenously to anesthetized dogs, a dose-related negative inotropic effect was documented. In a healthy volunteer study of itraconazole intravenous infusion, transient, asymptomatic decreases in left ventricular ejection fraction were observed using gated SPECT imaging; these resolved before the next infusion, 12 hours later. If signs or symptoms of congestive heart failure appear during administration of SPORANOX<sup>®</sup> Oral Solution, monitor carefully and consider other treatment alternatives which may include discontinuation of SPORANOX<sup>®</sup> Oral Solution administration. (See BOXED WARNING, CONTRAINDICATIONS, WARNINGS, PRECAUTIONS: Drug Interactions and ADVERSE REACTIONS: Post-marketing Experience for more information.)

### **Cystic Fibrosis:**

Seventeen cystic fibrosis patients, ages 7 to 28 years old, were administered itraconazole oral solution 2.5 mg/kg b.i.d. for 14 days in a pharmacokinetic study. Sixteen patients completed the study. Steady state trough concentrations >250 ng/mL were achieved in 6 out of 11 patients  $\geq 16$  years of age but in none of the 5 patients <16 years of age. Large variability was observed in the pharmacokinetic data (%CV for trough concentrations = 98% and 70% for  $\geq 16$  and <16 years, respectively; %CV for AUC = 75% and 58% for  $\geq 16$  and <16 years, respectively). If a patient with cystic fibrosis does not respond to SPORANOX<sup>®</sup> Oral Solution, consideration should be given to switching to alternative therapy.

### **Hydroxypropyl- $\beta$ -Cyclodextrin:**

The oral bioavailability of hydroxypropyl- $\beta$ -cyclodextrin given as a solubilizer of itraconazole in oral solution is on average lower than 0.5% and is similar to that of hydroxypropyl- $\beta$ -cyclodextrin alone. This low oral bioavailability of hydroxypropyl- $\beta$ -cyclodextrin is not modified by the presence of food and is similar after single and repeated administrations.

## **MICROBIOLOGY**

### **Mechanism of Action:**

*In vitro* studies have demonstrated that itraconazole inhibits the cytochrome P450-dependent synthesis of ergosterol, which is a vital component of fungal cell membranes.

### **Drug Resistance:**

Isolates from several fungal species with decreased susceptibility to itraconazole have been isolated *in vitro* and from patients receiving prolonged therapy.

*Candida krusei*, *Candida glabrata* and *Candida tropicalis* are generally the least susceptible *Candida* species, with some isolates showing unequivocal resistance to itraconazole *in vitro*.

Itraconazole is not active against *Zygomycetes* (e.g., *Rhizopus* spp., *Rhizomucor* spp., *Mucor* spp. and *Absidia* spp.), *Fusarium* spp., *Scedosporium* spp. and *Scopulariopsis* spp.

### **Cross-resistance:**

In systemic candidosis, if fluconazole-resistant strains of *Candida* species are suspected, it cannot be assumed that these are sensitive to itraconazole, hence their sensitivity should be tested before the start of itraconazole therapy.

Several *in vitro* studies have reported that some fungal clinical isolates, including *Candida* species, with reduced susceptibility to one azole antifungal agent may also be less susceptible to other azole derivatives. The finding of cross-resistance is dependent on a number of factors,

including the species evaluated, its clinical history, the particular azole compounds compared, and the type of susceptibility test that is performed.

Studies (both *in vitro* and *in vivo*) suggest that the activity of amphotericin B may be suppressed by prior azole antifungal therapy. As with other azoles, itraconazole inhibits the <sup>14</sup>C-demethylation step in the synthesis of ergosterol, a cell wall component of fungi. Ergosterol is the active site for amphotericin B. In one study the antifungal activity of amphotericin B against *Aspergillus fumigatus* infections in mice was inhibited by ketoconazole therapy. The clinical significance of test results obtained in this study is unknown.

### **Activity In Vitro and in Clinical Infections:**

Itraconazole has been shown to be active against most strains of the following microorganism, **both *in vitro* and in clinical infections.**

*Candida albicans*

### **Susceptibility Testing Methods**

*(Applicable to Candida isolates from patients with oropharyngeal or esophageal candidiasis)*

*Candida albicans*

The interpretive criteria and breakpoints for itraconazole against *Candida albicans* are applicable to tests performed using Clinical Laboratory and Standards Institute (CLSI) microbroth dilution reference method M27A for MIC (partial inhibition endpoint) read at 48 hours.

### **Broth Microdilution Techniques**

Quantitative methods are used to determine antifungal minimum inhibitory concentrations (MICs). These MICs provide estimates of the susceptibility of *Candida* spp. to antifungal agents. MICs should be determined using a standardized procedure at 48 hours. Standardized procedures are based on a microdilution method (broth)<sup>1,2</sup> with standardized inoculum concentrations and standardized concentrations of itraconazole powder. The MIC values should be interpreted according to the criteria provided in Table below:

<b>Susceptibility Interpretive Criteria for Itraconazole<sup>1,2</sup></b>			
<b>Pathogen</b>	<b>Broth Microdilution MIC* (µg/mL) at 48 Hours</b>		
	<b>S</b>	<b>I</b>	<b>R</b>
<i>Candida albicans</i>	≤ 0.125	0.25 – 0.5	≥ 1

\* A report of “Susceptible” indicates that the antimicrobial drug is likely to inhibit growth of the microorganism if the antimicrobial drug reaches the concentration usually achievable at the site of infection. A report of *intermediate* (I) category implies that an infection due to the isolate may be appropriately treated in the body sites where the drugs are physiologically concentrated or when a high dosage of drug is used. The *intermediate* category is sometimes called *Susceptible-Dose Dependent* (SDD) and both categories are equivalent for itraconazole.

A report of *Resistant* (R) indicates that the antimicrobial is not likely to inhibit growth of the pathogen if the antimicrobial drug reaches the concentration usually achievable at the infection site; other therapy should be selected.

### Quality Control

Standardized susceptibility test procedures require the use of quality control organisms to control the technical aspects of the test procedures. Standard itraconazole powder should provide the following range of values noted in the table below.

NOTE: Quality control microorganisms are specific strains of organisms with intrinsic biological properties relating to resistance mechanisms and their genetic expression within fungi; the specific strains used for microbiological control are not clinically significant.

<b>Acceptable Quality Control Ranges for Itraconazole to be used in Validation of Susceptibility Test Results<sup>1,2</sup></b>	
<b>QC Strain</b>	<b>Broth Microdilution MIC (µg/mL) at 48 Hours</b>
<i>Candida parapsilosis</i> ATCC† 22019	0.06-0.25
<i>Candida krusei</i> ATCC 6258	0.12-0.5

† ATCC is the registered trademark of the American Type Culture Collection.

## CLINICAL STUDIES

### Oropharyngeal Candidiasis:

Two randomized, controlled studies for the treatment of oropharyngeal candidiasis have been conducted (total n=344). In one trial, clinical response to either 7 or 14 days of itraconazole oral solution, 200 mg/day, was similar to fluconazole tablets and averaged 84% across all arms. Clinical response in this study was defined as cured or improved (only minimal signs and symptoms with no visible lesions). Approximately 5% of subjects were lost to follow-up before any evaluations could be performed. Response to 14 days therapy of itraconazole oral solution was associated with a lower relapse rate than 7 days of itraconazole therapy. In another trial, the clinical response rate (defined as cured or improved) for itraconazole oral solution was

similar to clotrimazole troches and averaged approximately 71% across both arms, with approximately 3% of subjects lost to follow-up before any evaluations could be performed. Ninety-two percent of the patients in these studies were HIV seropositive.

In an uncontrolled, open-label study of selected patients clinically unresponsive to fluconazole tablets (n=74, all patients HIV seropositive), patients were treated with itraconazole oral solution 100 mg b.i.d. (Clinically unresponsive to fluconazole in this study was defined as having received a dose of fluconazole tablets at least 200 mg/day for a minimum of 14 days.) Treatment duration was 14-28 days based on response. Approximately 55% of patients had complete resolution of oral lesions. Of patients who responded and then entered a follow-up phase (n=22), all relapsed within 1 month (median 14 days) when treatment was discontinued. Although baseline endoscopies had not been performed, several patients in this study developed symptoms of esophageal candidiasis while receiving therapy with itraconazole oral solution. Itraconazole oral solution has not been directly compared to other agents in a controlled trial of similar patients.

### **Esophageal Candidiasis:**

A double-blind randomized study (n=119, 111 of whom were HIV seropositive) compared itraconazole oral solution (100 mg/day) to fluconazole tablets (100 mg/day). The dose of each was increased to 200 mg/day for patients not responding initially. Treatment continued for 2 weeks following resolution of symptoms, for a total duration of treatment of 3-8 weeks. Clinical response (a global assessment of cured or improved) was not significantly different between the two study arms, and averaged approximately 86% with 8% lost to follow-up. Six of 53 (11%) itraconazole-treated patients and 12/57 (21%) fluconazole-treated patients were escalated to the 200 mg dose in this trial. Of the subgroup of patients who responded and entered a follow-up phase (n=88), approximately 23% relapsed across both arms within 4 weeks.

## **INDICATIONS AND USAGE**

SPORANOX<sup>®</sup> (itraconazole) Oral Solution is indicated for the treatment of oropharyngeal and esophageal candidiasis.

(See CLINICAL PHARMACOLOGY: Special Populations, WARNINGS, and ADVERSE REACTIONS: Post-marketing Experience for more information.)

## **CONTRAINDICATIONS**

### **Congestive Heart Failure:**

SPORANOX<sup>®</sup> (itraconazole) Oral Solution should not be administered to patients with evidence of ventricular dysfunction such as congestive heart failure (CHF) or a history of CHF

except for the treatment of life-threatening or other serious infections. (See BOXED WARNING, WARNINGS, PRECAUTIONS: Drug Interactions-Calcium Channel Blockers, ADVERSE REACTIONS: Post-marketing Experience, and CLINICAL PHARMACOLOGY: Special Populations.)

### **Drug Interactions:**

Coadministration of a number of CYP3A4 substrates are contraindicated with SPORANOX<sup>®</sup>. Plasma concentrations increase for the following drugs: methadone, disopyramide, dofetilide, dronedarone, quinidine, ergot alkaloids (such as dihydroergotamine, ergometrine (ergonovine), ergotamine, methylergometrine (methylergonovine)), irinotecan, lurasidone, oral midazolam, pimozide, triazolam, felodipine, nisoldipine, ranolazine, eplerenone, cisapride, lovastatin, simvastatin, ticagrelor and, in subjects with varying degrees of renal or hepatic impairment, colchicine, fesoterodine, telithromycin and solifenacin. (See PRECAUTIONS: Drug Interactions Section for specific examples.) This increase in drug concentrations caused by coadministration with itraconazole may increase or prolong both the pharmacologic effects and/or adverse reactions to these drugs. For example, increased plasma concentrations of some of these drugs can lead to QT prolongation and ventricular tachyarrhythmias including occurrences of torsade de pointes, a potentially fatal arrhythmia. Specific examples are listed in PRECAUTIONS: Drug Interactions.

SPORANOX<sup>®</sup> is contraindicated for patients who have shown hypersensitivity to itraconazole. There is limited information regarding cross-hypersensitivity between itraconazole and other azole antifungal agents. Caution should be used when prescribing SPORANOX<sup>®</sup> to patients with hypersensitivity to other azoles.

### **WARNINGS**

#### **Hepatic Effects:**

**SPORANOX<sup>®</sup> has been associated with rare cases of serious hepatotoxicity, including liver failure and death. Some of these cases had neither pre-existing liver disease nor a serious underlying medical condition, and some of these cases developed within the first week of treatment. If clinical signs or symptoms develop that are consistent with liver disease, treatment should be discontinued and liver function testing performed. Continued SPORANOX<sup>®</sup> use or reinstatement of treatment with SPORANOX<sup>®</sup> is strongly discouraged unless there is a serious or life-threatening situation where the expected benefit exceeds the risk. (See PRECAUTIONS: Information for Patients and ADVERSE REACTIONS.)**

### **Cardiac Dysrhythmias:**

Life-threatening cardiac dysrhythmias and/or sudden death have occurred in patients using drugs such as cisapride, pimozide, methadone, or quinidine concomitantly with SPORANOX<sup>®</sup> and/or other CYP3A4 inhibitors. Concomitant administration of these drugs with SPORANOX<sup>®</sup> is contraindicated. (See BOXED WARNING, CONTRAINDICATIONS, and PRECAUTIONS: Drug Interactions.)

### **Cardiac Disease:**

SPORANOX<sup>®</sup> Oral Solution should not be used in patients with evidence of ventricular dysfunction unless the benefit clearly outweighs the risk. For patients with risk factors for congestive heart failure, physicians should carefully review the risks and benefits of SPORANOX<sup>®</sup> therapy. These risk factors include cardiac disease such as ischemic and valvular disease; significant pulmonary disease such as chronic obstructive pulmonary disease; and renal failure and other edematous disorders. Such patients should be informed of the signs and symptoms of CHF, should be treated with caution, and should be monitored for signs and symptoms of CHF during treatment. If signs or symptoms of CHF appear during administration of SPORANOX<sup>®</sup> Oral Solution, monitor carefully and consider other treatment alternatives which may include discontinuation of SPORANOX<sup>®</sup> Oral Solution administration.

Itraconazole has been shown to have a negative inotropic effect. When itraconazole was administered intravenously to anesthetized dogs, a dose-related negative inotropic effect was documented. In a healthy volunteer study of itraconazole intravenous infusion, transient, asymptomatic decreases in left ventricular ejection fraction were observed using gated SPECT imaging; these resolved before the next infusion, 12 hours later.

SPORANOX<sup>®</sup> has been associated with reports of congestive heart failure. In post-marketing experience, heart failure was more frequently reported in patients receiving a total daily dose of 400 mg although there were also cases reported among those receiving lower total daily doses.

Calcium channel blockers can have negative inotropic effects which may be additive to those of itraconazole. In addition, itraconazole can inhibit the metabolism of calcium channel blockers. Therefore, caution should be used when co-administering itraconazole and calcium channel blockers due to an increased risk of CHF. Concomitant administration of SPORANOX<sup>®</sup> and felodipine or nisoldipine is contraindicated.

Cases of CHF, peripheral edema, and pulmonary edema have been reported in the post-marketing period among patients being treated for onychomycosis and/or systemic fungal infections. (See CONTRAINDICATIONS, CLINICAL PHARMACOLOGY: Special

Populations, PRECAUTIONS: Drug Interactions, and ADVERSE REACTIONS: Post-marketing Experience for more information.)

**Interaction potential:**

SPORANOX<sup>®</sup> has a potential for clinically important drug interactions. Coadministration of specific drugs with itraconazole may result in changes in efficacy of itraconazole and/or the coadministered drug, life-threatening effects and/or sudden death. Drugs that are contraindicated, not recommended or recommended for use with caution in combination with itraconazole are listed in PRECAUTIONS: Drug Interactions.

**Interchangeability:**

SPORANOX<sup>®</sup> (itraconazole) Oral Solution and SPORANOX<sup>®</sup> Capsules should not be used interchangeably. This is because drug exposure is greater with the Oral Solution than with the Capsules when the same dose of drug is given. Only SPORANOX<sup>®</sup> Oral Solution has been demonstrated effective for oral and/or esophageal candidiasis.

**Hydroxypropyl- $\beta$ -cyclodextrin:**

SPORANOX<sup>®</sup> Oral Solution contains the excipient hydroxypropyl- $\beta$ -cyclodextrin which produced adenocarcinomas in the large intestine and exocrine pancreatic adenocarcinomas in a rat carcinogenicity study. These findings were not observed in a similar mouse carcinogenicity study. The clinical relevance of these adenocarcinomas is unknown. (See PRECAUTIONS: Carcinogenesis, Mutagenesis, and Impairment of Fertility.)

**Treatment of Severely Neutropenic Patients:**

SPORANOX<sup>®</sup> Oral Solution as treatment for oropharyngeal and/or esophageal candidiasis was not investigated in severely neutropenic patients. Due to its pharmacokinetic properties, SPORANOX<sup>®</sup> Oral Solution is not recommended for initiation of treatment in patients at immediate risk of systemic candidiasis.

**PRECAUTIONS**

**Hepatotoxicity:**

Rare cases of serious hepatotoxicity have been observed with SPORANOX<sup>®</sup> treatment, including some cases within the first week. It is recommended that liver function monitoring be considered in all patients receiving SPORANOX<sup>®</sup>. Treatment should be stopped immediately and liver function testing should be conducted in patients who develop signs and symptoms suggestive of liver dysfunction.

**Neuropathy:**

If neuropathy occurs that may be attributable to SPORANOX<sup>®</sup> Oral Solution, the treatment should be discontinued.

### **Cystic Fibrosis:**

If a patient with cystic fibrosis does not respond to SPORANOX<sup>®</sup> Oral Solution, consideration should be given to switching to alternative therapy (see CLINICAL PHARMACOLOGY: Special Populations).

### **Hearing Loss:**

Transient or permanent hearing loss has been reported in patients receiving treatment with itraconazole. Several of these reports included concurrent administration of quinidine which is contraindicated (see BOXED WARNING: Drug Interactions, CONTRAINDICATIONS: Drug Interactions and PRECAUTIONS: Drug Interactions). The hearing loss usually resolves when treatment is stopped, but can persist in some patients.

### **Information for Patients:**

- Only SPORANOX<sup>®</sup> Oral Solution has been demonstrated effective for oral and/or esophageal candidiasis.
- SPORANOX<sup>®</sup> Oral Solution contains the excipient hydroxypropyl- $\beta$ -cyclodextrin which produced adenocarcinomas in the large intestine and exocrine pancreatic adenocarcinomas in a rat carcinogenicity study. These findings were not observed in a similar mouse carcinogenicity study. The clinical relevance of these adenocarcinomas is unknown. (See Carcinogenesis, Mutagenesis, and Impairment of Fertility.)
- Taking SPORANOX<sup>®</sup> Oral Solution under fasted conditions improves the systemic availability of itraconazole. Instruct patients to take SPORANOX<sup>®</sup> Oral Solution without food, if possible.
- SPORANOX<sup>®</sup> Oral Solution should not be used interchangeably with SPORANOX<sup>®</sup> Capsules.
- Instruct patients about the signs and symptoms of congestive heart failure, and if these signs or symptoms occur during SPORANOX<sup>®</sup> administration, they should discontinue SPORANOX<sup>®</sup> and contact their healthcare provider immediately.
- Instruct patients to stop SPORANOX<sup>®</sup> treatment immediately and contact their healthcare provider if any signs and symptoms suggestive of liver dysfunction develop. Such signs and symptoms may include unusual fatigue, anorexia, nausea and/or vomiting, jaundice, dark urine, or pale stools.
- Instruct patients to contact their physician before taking any concomitant medications with itraconazole to ensure there are no potential drug interactions.
- Instruct patients that hearing loss can occur with the use of itraconazole. The hearing loss usually resolves when treatment is stopped, but can persist in some patients. Advise patients to discontinue therapy and inform their physicians if any hearing loss symptoms occur.

- Instruct patients that dizziness or blurred/double vision can sometimes occur with itraconazole. Advise patients that if they experience these events, they should not drive or use machines.

### **Drug Interactions:**

Itraconazole is mainly metabolized through CYP3A4. Other drugs that either share this metabolic pathway or modify CYP3A4 activity may influence the pharmacokinetics of itraconazole. Similarly, itraconazole may modify the pharmacokinetics of other drugs that share this metabolic pathway. Itraconazole is a potent CYP3A4 inhibitor and a P-glycoprotein inhibitor. When using concomitant medication, it is recommended that the corresponding label be consulted for information on the route of metabolism and the possible need to adjust dosages.

#### **Drugs that may decrease itraconazole plasma concentrations**

Coadministration of itraconazole with potent enzyme inducers of CYP3A4 may decrease the bioavailability of itraconazole and hydroxy-itraconazole to such an extent that efficacy may be reduced. Examples include:

- Antibacterials: isoniazid, rifabutin (see also under ‘Drugs that may have their plasma concentrations increased by itraconazole’), rifampicin
- Anticonvulsants: carbamazepine (see also under ‘Drugs that may have their plasma concentrations increased by itraconazole’), phenobarbital, phenytoin
- Antivirals: efavirenz, nevirapine

Therefore, administration of potent enzyme inducers of CYP3A4 with itraconazole is not recommended. It is recommended that the use of these drugs be avoided from 2 weeks before and during treatment with itraconazole, unless the benefits outweigh the risk of potentially reduced itraconazole efficacy. Upon coadministration, it is recommended that the antifungal activity be monitored and the itraconazole dose increased as deemed necessary.

#### **Drugs that may increase itraconazole plasma concentrations**

Potent inhibitors of CYP3A4 may increase the bioavailability of itraconazole. Examples include:

- Antibacterials: ciprofloxacin, clarithromycin, erythromycin
- Antivirals: ritonavir-boosted darunavir, ritonavir-boosted fosamprenavir, indinavir (see also under ‘Drugs that may have their plasma concentrations increased by itraconazole’), ritonavir (see also under ‘Drugs that may have their plasma concentrations increased by itraconazole’) and telaprevir.

It is recommended that these drugs be used with caution when coadministered with itraconazole oral solution. It is recommended that patients who must take itraconazole

concomitantly with potent inhibitors of CYP3A4 be monitored closely for signs or symptoms of increased or prolonged pharmacologic effects of itraconazole, and the itraconazole dose be decreased as deemed necessary.

#### Drugs that may have their plasma concentrations increased by itraconazole

Itraconazole and its major metabolite, hydroxy-itraconazole, can inhibit the metabolism of drugs metabolized by CYP3A4 and can inhibit the drug transport by P-glycoprotein, which may result in increased plasma concentrations of these drugs and/or their active metabolite(s) when they are administered with itraconazole. These elevated plasma concentrations may increase or prolong both therapeutic and adverse effects of these drugs. CYP3A4-metabolized drugs known to prolong the QT interval may be contraindicated with itraconazole, since the combination may lead to ventricular tachyarrhythmias including occurrences of torsade de pointes, a potentially fatal arrhythmia. Once treatment is stopped, itraconazole plasma concentrations decrease to an almost undetectable concentration within 7 to 14 days, depending on the dose and duration of treatment. In patients with hepatic cirrhosis or in subjects receiving CYP3A4 inhibitors, the decline in plasma concentrations may be even more gradual. This is particularly important when initiating therapy with drugs whose metabolism is affected by itraconazole.

Examples of drugs that may have their plasma concentrations increased by itraconazole presented by drug class with advice regarding coadministration with itraconazole:

<b>Table 1: Drugs that may have their plasma concentrations increased by itraconazole</b>				
<b>Drug Class</b>	<b>Contraindicated</b>	<b>Not Recommended</b>	<b>Use with Caution</b>	<b>Comments</b>
	<i>Under no circumstances is the drug to be coadministered with itraconazole, and up to two weeks after discontinuation of treatment with itraconazole.</i>	<i>It is recommended that the use of the drug be avoided during and up to two weeks after discontinuation of treatment with itraconazole, unless the benefits outweigh the potentially increased risks of side effects. If coadministration cannot be avoided, clinical monitoring for signs or symptoms of increased or prolonged effects or side effects of the interacting drug is recommended, and its dosage be reduced or interrupted as deemed necessary. When appropriate, it is recommended that plasma concentrations be measured. The label of the coadministered drug should be consulted for information on dose adjustment and adverse effects.</i>	<i>Careful monitoring is recommended when the drug is coadministered with itraconazole. Upon coadministration, it is recommended that patients be monitored closely for signs or symptoms of increased or prolonged effects or side effects of the interacting drug, and its dosage be reduced as deemed necessary. When appropriate, it is recommended that plasma concentrations be measured. The label of the coadministered drug should be consulted for information on dose adjustment and adverse effects.</i>	
Alpha Blockers		tamsulosin		

<b>Table 1: Drugs that may have their plasma concentrations increased by itraconazole</b>				
<b>Drug Class</b>	<b>Contraindicated</b>	<b>Not Recommended</b>	<b>Use with Caution</b>	<b>Comments</b>
Analgesics	methadone		alfentanil, buprenorphine IV and sublingual, fentanyl, oxycodone, sufentanil	<p><b>Methadone:</b> The potential increase in plasma concentrations of methadone when coadministered with SPORANOX<sup>®</sup> may increase the risk of serious cardiovascular events including QTc prolongation and <i>torsade de pointes</i>.</p> <p><b>Fentanyl:</b> The potential increase in plasma concentrations of fentanyl when coadministered with SPORANOX<sup>®</sup> may increase the risk of potentially fatal respiratory depression.</p> <p><b>Sufentanil:</b> No human pharmacokinetic data of an interaction with itraconazole are available. In vitro data suggest that sufentanil is metabolized by CYP3A4 and so potentially increased sufentanil plasma concentrations would be expected when coadministered with SPORANOX<sup>®</sup>.</p>
Antiarrhythmics	disopyramide, dofetilide, dronedaronone, quinidine		digoxin	<p><b>Disopyramide, dofetilide, dronedaronone, quinidine:</b> The potential increase in plasma concentrations of these drugs when coadministered with SPORANOX<sup>®</sup> may increase the risk of serious cardiovascular events including QTc prolongation.</p>

<b>Table 1: Drugs that may have their plasma concentrations increased by itraconazole</b>				
<b>Drug Class</b>	<b>Contraindicated</b>	<b>Not Recommended</b>	<b>Use with Caution</b>	<b>Comments</b>
Antibacterials	telithromycin, in subjects with severe renal impairment or severe hepatic impairment	rifabutin	telithromycin	<p><b>Telithromycin:</b> The potential increase in plasma concentrations of telithromycin in subjects with severe renal impairment or severe hepatic impairment, when coadministered with SPORANOX<sup>®</sup> may increase the risk of serious cardiovascular events including QT prolongation and torsade de pointes.</p> <p><b>Rifabutin:</b> See also under ‘Drugs that may decrease itraconazole plasma concentrations’.</p>
Anticoagulants and Antiplatelet Drugs	ticagrelor	apixaban, rivaroxaban	coumarins, cilostazol, dabigatran	<p><b>Ticagrelor:</b> The potential increase in plasma concentrations of ticagrelor may increase the risk of bleeding.</p> <p><b>Coumarins:</b> SPORANOX<sup>®</sup> may enhance the anticoagulant effect of coumarin-like drugs, such as warfarin.</p>

<b>Table 1: Drugs that may have their plasma concentrations increased by itraconazole</b>				
<b>Drug Class</b>	<b>Contraindicated</b>	<b>Not Recommended</b>	<b>Use with Caution</b>	<b>Comments</b>
Anticonvulsants		carbamazepine		<b>Carbamazepine:</b> <i>In vivo</i> studies have demonstrated an increase in plasma carbamazepine concentrations in subjects concomitantly receiving ketoconazole. Although there are no data regarding the effect of itraconazole on carbamazepine metabolism, because of the similarities between ketoconazole and itraconazole, concomitant administration of SPORANOX <sup>®</sup> and carbamazepine may inhibit the metabolism of carbamazepine. See also under 'Drugs that may decrease itraconazole plasma concentrations'.
Antidiabetics			repaglinide, saxagliptin	
Anthelmintics and Antiprotozoals			praziquantel	
Antimigraine Drugs	ergot alkaloids, such as dihydroergotamine, ergometrine (ergonovine), ergotamine, methylergometrine (methylergonovine)		eletriptan	<b>Ergot Alkaloids:</b> The potential increase in plasma concentrations of ergot alkaloids when coadministered with SPORANOX <sup>®</sup> may increase the risk of ergotism, ie. a risk for vasospasm potentially leading to cerebral ischemia and/or ischemia of the extremities.

<b>Table 1: Drugs that may have their plasma concentrations increased by itraconazole</b>				
<b>Drug Class</b>	<b>Contraindicated</b>	<b>Not Recommended</b>	<b>Use with Caution</b>	<b>Comments</b>
Antineoplastics	irinotecan	axitinib, dabrafenib, dasatinib, ibrutinib, nilotinib, sunitinib	bortezomib, busulphan, docetaxel, erlotinib, imatinib, ixabepilone, ponatinib, lapatinib, trimetrexate, vinca alkaloids	<b>Irinotecan:</b> The potential increase in plasma concentrations of irinotecan when coadministered with SPORANOX <sup>®</sup> may increase the risk of potentially fatal adverse events.
Antipsychotics, Anxiolytics and Hypnotics	lurasidone, oral midazolam, pimozide, triazolam		alprazolam, aripiprazole, buspirone, diazepam, haloperidol, midazolam IV, perospirone, quetiapine, ramelteon, risperidone	<b>Midazolam, triazolam:</b> Coadministration of SPORANOX <sup>®</sup> and oral midazolam, or triazolam may cause several-fold increases in plasma concentrations of these drugs. This may potentiate and prolong hypnotic and sedative effects, especially with repeated dosing or chronic administration of these agents.  <b>Pimozide:</b> The potential increase in plasma concentrations of pimozide when coadministered with SPORANOX <sup>®</sup> may increase the risk of serious cardiovascular events including QTc prolongation and <i>torsade de pointes</i> .
Antivirals		simprevir	maraviroc, indinavir, ritonavir, saquinavir	<b>Indinavir, ritonavir:</b> See also under 'Drugs that may increase itraconazole plasma concentrations'.
Beta Blockers			nadolol	

<b>Table 1: Drugs that may have their plasma concentrations increased by itraconazole</b>				
<b>Drug Class</b>	<b>Contraindicated</b>	<b>Not Recommended</b>	<b>Use with Caution</b>	<b>Comments</b>
Calcium Channel Blockers	felodipine, nisoldipine		other dihydropyridines, verapamil	<b>Calcium channel blockers</b> can have a negative inotropic effect which may be additive to those of itraconazole. The potential increase in plasma concentrations of calcium channel blockers when co-administered with SPORANOX <sup>®</sup> may increase the risk of congestive heart failure. <b>Dihydropyridines:</b> Concomitant administration of SPORANOX <sup>®</sup> may cause several-fold increases in plasma concentrations of dihydropyridines. Edema has been reported in patients concomitantly receiving SPORANOX <sup>®</sup> and dihydropyridine calcium channel blockers.
Cardiovascular Drugs, Miscellaneous	ranolazine	aliskiren, sildenafil, for the treatment of pulmonary hypertension	bosentan, riociguat	<b>Ranolazine:</b> The potential increase in plasma concentrations of ranolazine when coadministered with SPORANOX <sup>®</sup> may increase the risk of serious cardiovascular events including QTc prolongation.
Diuretics	eplerenone			<b>Eplerenone:</b> The potential increase in plasma concentrations of eplerenone when coadministered with SPORANOX <sup>®</sup> may increase the risk of hyperkalemia and hypotension.
Gastrointestinal Drugs	cisapride		aprepitant	<b>Cisapride:</b> The potential increase in plasma concentrations of cisapride when coadministered with SPORANOX <sup>®</sup> may increase the risk of serious cardiovascular events including QTc prolongation.

<b>Table 1: Drugs that may have their plasma concentrations increased by itraconazole</b>				
<b>Drug Class</b>	<b>Contraindicated</b>	<b>Not Recommended</b>	<b>Use with Caution</b>	<b>Comments</b>
Immunosuppressants		everolimus, temsirolimus	budesonide, ciclesonide, cyclosporine, dexamethasone, fluticasone, methylprednisolone, rapamycin (also known as sirolimus), tacrolimus	
Lipid Regulating Drugs	lovastatin, simvastatin		atorvastatin	The potential increase in plasma concentrations of atorvastatin, lovastatin, and simvastatin when coadministered with SPORANOX <sup>®</sup> may increase the risk of skeletal muscle toxicity, including rhabdomyolysis.
Respiratory Drugs		salmeterol		
Urological Drugs	fesoterodine, in subjects with moderate to severe renal impairment, or moderate to severe hepatic impairment, solifenacin, in subjects with severe renal impairment or moderate to severe hepatic impairment	darifenacin, vardenafil	fesoterodine, oxybutynin sildenafil, for the treatment of erectile dysfunction, solifenacin, tadalafil, tolterodine	<b>Fesoterodine:</b> The potential increase in plasma concentrations of the fesoterodine active metabolite may be greater in subjects with moderate to severe renal impairment, or moderate to severe hepatic impairment, which may lead to an increased risk of adverse reactions. <b>Solifenacin:</b> The potential increase in plasma concentrations of solifenacin in subjects with severe renal impairment or moderate to severe hepatic impairment, when coadministered with SPORANOX <sup>®</sup> may increase the risk of serious cardiovascular events including QT prolongation.

<b>Table 1: Drugs that may have their plasma concentrations increased by itraconazole</b>				
<b>Drug Class</b>	<b>Contraindicated</b>	<b>Not Recommended</b>	<b>Use with Caution</b>	<b>Comments</b>
Other	colchicine, in subjects with renal or hepatic impairment	colchicine, conivaptan, tolvaptan	cinacalcet	<p><b>Colchicine:</b> The potential increase in plasma concentrations of colchicine when coadministered with SPORANOX<sup>®</sup> may increase the risk of potentially fatal adverse events.</p> <p><b>Conivaptan and Tolvaptan:</b> A safe and effective dose of either conivaptan or tolvaptan has not been established when coadministered with SPORANOX<sup>®</sup>.</p>

## Drugs that may have their plasma concentrations decreased by itraconazole

Coadministration of itraconazole with the NSAID meloxicam may decrease the plasma concentration of meloxicam. It is recommended that meloxicam be used with caution when coadministered with itraconazole, and its effects or side effects be monitored. It is recommended that the dosage of meloxicam, if coadministered with itraconazole, be adjusted if necessary.

## Pediatric Population

Interaction studies have only been performed in adults.

## **Carcinogenesis, Mutagenesis, and Impairment of Fertility:**

### Itraconazole

Itraconazole showed no evidence of carcinogenicity potential in mice treated orally for 23 months at dosage levels up to 80 mg/kg/day (approximately 10x the maximum recommended human dose [MRHD]). Male rats treated with 25 mg/kg/day (3.1x MRHD) had a slightly increased incidence of soft tissue sarcoma. These sarcomas may have been a consequence of hypercholesterolemia, which is a response of rats, but not dogs or humans, to chronic itraconazole administration. Female rats treated with 50 mg/kg/day (6.25x MRHD) had an increased incidence of squamous cell carcinoma of the lung (2/50) as compared to the untreated group. Although the occurrence of squamous cell carcinoma in the lung is extremely uncommon in untreated rats, the increase in this study was not statistically significant.

Itraconazole produced no mutagenic effects when assayed in DNA repair test (unscheduled DNA synthesis) in primary rat hepatocytes, in Ames tests with *Salmonella typhimurium* (6 strains) and *Escherichia coli*, in the mouse lymphoma gene mutation tests, in a sex-linked recessive lethal mutation (*Drosophila melanogaster*) test, in chromosome aberration tests in human lymphocytes, in a cell transformation test with C3H/10T $\frac{1}{2}$  C18 mouse embryo fibroblasts cells, in a dominant lethal mutation test in male and female mice, and in micronucleus tests in mice and rats.

Itraconazole did not affect the fertility of male or female rats treated orally with dosage levels of up to 40 mg/kg/day (5x MRHD), even though parental toxicity was present at this dosage level. More severe signs of parental toxicity, including death, were present in the next higher dosage level, 160 mg/kg/day (20x MRHD).

### Hydroxypropyl- $\beta$ -cyclodextrin (HP- $\beta$ -CD)

Hydroxypropyl- $\beta$ -cyclodextrin (HP- $\beta$ -CD) is the solubilizing excipient used in SPORANOX<sup>®</sup> Oral Solution.

Hydroxypropyl- $\beta$ -cyclodextrin (HP- $\beta$ -CD) was found to produce neoplasms in the large intestine at 5000 mg/kg/day in rat carcinogenicity study. This dose was about 6 times amount contained in the recommended clinical dose of SPORANOX<sup>®</sup> Oral Solution based on body surface area comparisons. The clinical relevance of this finding is unknown. The slightly higher incidence of adenocarcinomas in the large intestines was linked to the hypertrophic/hyperplastic and inflammatory changes in the colonic mucosa brought about by HP- $\beta$ -CD-induced increased osmotic forces.

In addition, HP- $\beta$ -CD was found to produce pancreatic exocrine hyperplasia and neoplasia when administered orally to rats at doses of 500, 2000 or 5000 mg/kg/day for 25 months. Adenocarcinomas of the exocrine pancreas produced in the treated animals were not seen in the untreated group and are not reported in the historical controls. The recommended clinical dose of SPORANOX<sup>®</sup> Oral Solution contains approximately 1.7 times the amount of HP- $\beta$ -CD as was in the 500 mg/kg/day dose, based on body surface area comparisons. This finding was not observed in the mouse carcinogenicity study at doses of 500, 2000 or 5000 mg/kg/day for 22-23 months. This finding was also not observed in a 12-month toxicity study in dogs or in a 2-year toxicity study in female cynomolgus monkeys.

Since the development of the pancreatic tumors may be related to a mitogenic action of cholecystokinin and since there is no evidence that cholecystokinin has a mitogenic action in man, the clinical relevance of these findings is unknown.

HP- $\beta$ -CD has no antifertile effect, and is not mutagenic.

### **Pregnancy: Teratogenic effects. Pregnancy Category C:**

Itraconazole was found to cause a dose-related increase in maternal toxicity, embryotoxicity, and teratogenicity in rats at dosage levels of approximately 40-160 mg/kg/day (5-20x MRHD), and in mice at dosage levels of approximately 80 mg/kg/day (10x MRHD). Itraconazole has been shown to cross the placenta in a rat model. In rats, the teratogenicity consisted of major skeletal defects; in mice, it consisted of encephaloceles and/or macroglossia.

SPORANOX<sup>®</sup> Oral Solution contains the excipient hydroxypropyl- $\beta$ -cyclodextrin (HP- $\beta$ -CD). HP- $\beta$ -CD has no direct embryotoxic and no teratogenic effect.

There are no studies in pregnant women. SPORANOX<sup>®</sup> should be used in pregnancy only if the benefit outweighs the potential risk.

During post-marketing experience, cases of congenital abnormalities have been reported. (See ADVERSE REACTIONS: Post-marketing Experience.)

### **Nursing Mothers:**

Itraconazole is excreted in human milk; therefore, the expected benefits of SPORANOX<sup>®</sup> therapy for the mother should be weighed against the potential risk from exposure of itraconazole to the infant. The U.S. Public Health Service Centers for Disease Control and Prevention advises HIV-infected women not to breast-feed to avoid potential transmission of HIV to uninfected infants.

### **Pediatric Use:**

The efficacy and safety of SPORANOX<sup>®</sup> have not been established in pediatric patients.

The long-term effects of itraconazole on bone growth in children are unknown. In three toxicology studies using rats, itraconazole induced bone defects at dosage levels as low as 20 mg/kg/day (2.5x MRHD). The induced defects included reduced bone plate activity, thinning of the zona compacta of the large bones, and increased bone fragility. At a dosage level of 80 mg/kg/day (10x MRHD) over 1 year or 160 mg/kg/day (20x MRHD) for 6 months, itraconazole induced small tooth pulp with hypocellular appearance in some rats.

### **Geriatric Use:**

Clinical studies of SPORANOX<sup>®</sup> Oral Solution did not include sufficient numbers of subjects aged 65 years and over to determine whether they respond differently from younger subjects. It is advised to use SPORANOX<sup>®</sup> Oral Solution in these patients only if it is determined that the potential benefit outweighs the potential risks. In general, it is recommended that the dose selection for an elderly patient should be taken into consideration, reflecting the greater frequency of decreased hepatic, renal, or cardiac function, and of concomitant disease or other drug therapy.

Transient or permanent hearing loss has been reported in elderly patients receiving treatment with itraconazole. Several of these reports included concurrent administration of quinidine which is contraindicated (see **BOXED WARNING: Drug Interactions**, **CONTRAINDICATIONS: Drug Interactions** and **PRECAUTIONS: Drug Interactions**).

### **Renal Impairment:**

Limited data are available on the use of oral itraconazole in patients with renal impairment. The exposure of itraconazole may be lower in some patients with renal impairment. Caution should be exercised when itraconazole is administered in this patient population and dose adjustment may be needed. (See **CLINICAL PHARMACOLOGY: Special Populations** and **DOSAGE AND ADMINISTRATION**.)

### Hepatic Impairment:

Limited data are available on the use of oral itraconazole in patients with hepatic impairment. Caution should be exercised when this drug is administered in this patient population. It is recommended that patients with impaired hepatic function be carefully monitored when taking SPORANOX<sup>®</sup>. It is recommended that the prolonged elimination half-life of itraconazole observed in the single oral dose clinical trial with itraconazole capsules in cirrhotic patients be considered when deciding to initiate therapy with other medications metabolized by CYP3A4.

In patients with elevated or abnormal liver enzymes or active liver disease, or who have experienced liver toxicity with other drugs, treatment with SPORANOX<sup>®</sup> is strongly discouraged unless there is a serious or life-threatening situation where the expected benefit exceeds the risk. It is recommended that liver function monitoring be done in patients with pre-existing hepatic function abnormalities or those who have experienced liver toxicity with other medications. (See CLINICAL PHARMACOLOGY: Special Populations and DOSAGE AND ADMINISTRATION.)

### ADVERSE REACTIONS

Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a drug cannot be directly compared to rates in the clinical trials of another drug and may not reflect the rates observed in clinical practice.

SPORANOX<sup>®</sup> has been associated with rare cases of serious hepatotoxicity, including liver failure and death. Some of these cases had neither pre-existing liver disease nor a serious underlying medical condition. If clinical signs or symptoms develop that are consistent with liver disease, treatment should be discontinued and liver function testing performed. The risks and benefits of SPORANOX<sup>®</sup> use should be reassessed. (See WARNINGS: Hepatic Effects and PRECAUTIONS: Hepatotoxicity and Information for Patients.)

### Adverse Events Reported in Oropharyngeal or Esophageal Candidiasis Trials

U.S. adverse experience data are derived from 350 immunocompromised patients (332 HIV seropositive/AIDS) treated for oropharyngeal or esophageal candidiasis. Table 2 below lists adverse events reported by at least 2% of patients treated with SPORANOX<sup>®</sup> Oral Solution in U.S. clinical trials. Data on patients receiving comparator agents in these trials are included for comparison.

**Table 2: Summary of Adverse Events Reported by  $\geq 2\%$  of SPORANOX<sup>®</sup> Treated Patients in U.S. Clinical Trials (Total)**

Body System/ Adverse	Itraconazole		Fluconazole	Clotrimazole (n = 81 <sup>†</sup> ) %
	Total	All		

Event	(n = 350*) %	controlled studies (n = 272) %	(n = 125†) %	
<b>Gastrointestinal disorders</b>				
Nausea	11	10	11	5
Diarrhea	11	10	10	4
Vomiting	7	6	8	1
Abdominal pain	6	4	7	7
Constipation	2	2	1	0
<b>Body as a whole</b>				
Fever	7	6	8	5
Chest pain	3	3	2	0
Pain	2	2	4	0
Fatigue	2	1	2	0
<b>Respiratory disorders</b>				
Coughing	4	4	10	0
Dyspnea	2	3	5	1
Pneumonia	2	2	0	0
Sinusitis	2	2	4	0
Sputum increased	2	3	3	1
<b>Skin and appendages disorders</b>				
Rash	4	5	4	6
Increased sweating	3	4	6	1
Skin disorder unspecified	2	2	2	1
<b>Central/peripheral nervous system</b>				
Headache	4	4	6	6
Dizziness	2	2	4	1
<b>Resistance mechanism disorders</b>				
Pneumocystis carinii infection	2	2	2	0
<b>Psychiatric disorders</b>				
Depression	2	1	0	1

\* Of the 350 patients, 209 were treated for oropharyngeal candidiasis in controlled studies, 63 were treated for esophageal candidiasis in controlled studies and 78 were treated for oropharyngeal candidiasis in an open study.

† Of the 125 patients, 62 were treated for oropharyngeal candidiasis and 63 were treated for esophageal candidiasis.

‡ All 81 patients were treated for oropharyngeal candidiasis.

Adverse events reported by less than 2% of patients in U.S. clinical trials with SPORANOX<sup>®</sup> included: adrenal insufficiency, asthenia, back pain, dehydration, dyspepsia, dysphagia, flatulence, gynecomastia, hematuria, hemorrhoids, hot flushes, implantation complication, infection unspecified, injury, insomnia, male breast pain, myalgia, pharyngitis, pruritus, rhinitis, rigors, stomatitis ulcerative, taste perversion, tinnitus, upper respiratory tract infection, vision abnormal, and weight decrease. Edema, hypokalemia and menstrual disorders have been reported in clinical trials with itraconazole capsules.

### **Adverse Events Reported from Other Clinical Trials**

A comparative clinical trial in patients who received intravenous itraconazole followed by SPORANOX<sup>®</sup> Oral Solution or received Amphotericin B reported the following adverse events in the itraconazole intravenous/SPORANOX<sup>®</sup> Oral Solution treatment arm which are not listed above in the subsection “Adverse Events Reported in Oropharyngeal or Esophageal Candidiasis Trials” or listed below as postmarketing reports of adverse drug reactions: serum creatinine increased, blood urea nitrogen increased, renal function abnormal, hypocalcemia, hypomagnesemia, hypophosphatemia, hypotension, tachycardia and pulmonary infiltration.

In addition, the following adverse drug reactions were reported in patients who participated in SPORANOX<sup>®</sup> Oral Solution clinical trials:

*Cardiac Disorders:* cardiac failure;

*General Disorders and Administration Site Conditions:* edema;

*Hepatobiliary Disorders:* hepatic failure, hyperbilirubinemia;

*Metabolism and Nutrition Disorders:* hypokalemia;

*Reproductive System and Breast Disorders:* menstrual disorder

The following is a list of additional adverse drug reactions associated with itraconazole that have been reported in clinical trials of SPORANOX<sup>®</sup> Capsules and itraconazole IV excluding the adverse reaction term “Injection site inflammation” which is specific to the injection route of administration:

*Cardiac Disorders:* left ventricular failure;

*Gastrointestinal Disorders:* gastrointestinal disorder;

*General Disorders and Administration Site Conditions:* face edema;

*Hepatobiliary Disorders:* jaundice, hepatic function abnormal;

*Investigations:* alanine aminotransferase increased, aspartate aminotransferase increased, blood alkaline phosphatase increased, blood lactate dehydrogenase increased, gamma-glutamyltransferase increased, urine analysis abnormal;

*Metabolism and Nutrition Disorders:* hyperglycemia, hyperkalemia;

*Nervous System Disorders:* somnolence;

*Psychiatric Disorders:* confusional state;

*Renal and Urinary Disorders:* renal impairment;

*Respiratory, Thoracic and Mediastinal Disorders:* dysphonia;

*Skin and Subcutaneous Tissue Disorders:* rash erythematous;

*Vascular Disorders:* hypertension

In addition, the following adverse drug reaction was reported in children only who participated in SPORANOX<sup>®</sup> Oral Solution clinical trials: mucosal inflammation.

### **Post-marketing Experience**

Adverse drug reactions that have been first identified during post-marketing experience with SPORANOX<sup>®</sup> (all formulations) are listed in the table below. Because these reactions are reported voluntarily from a population of uncertain size, reliably estimating their frequency or establishing a causal relationship to drug exposure is not always possible.

**Table 3: Postmarketing Reports of Adverse Drug Reactions**

---

<b>Blood and Lymphatic System Disorders:</b>	Leukopenia, neutropenia, thrombocytopenia
<b>Immune System Disorders:</b>	Anaphylaxis; anaphylactic, anaphylactoid and allergic reactions; serum sickness; angioneurotic edema
<b>Metabolism and Nutrition Disorders:</b>	Hypertriglyceridemia
<b>Nervous System Disorders:</b>	Peripheral neuropathy, paresthesia, hypoesthesia, tremor
<b>Eye Disorders:</b>	Visual disturbances, including vision blurred and diplopia
<b>Ear and Labyrinth Disorders:</b>	Transient or permanent hearing loss
<b>Cardiac Disorders:</b>	Congestive heart failure
<b>Respiratory, Thoracic and Mediastinal Disorders:</b>	Pulmonary edema
<b>Gastrointestinal Disorders:</b>	Pancreatitis
<b>Hepatobiliary Disorders:</b>	Serious hepatotoxicity (including some cases of fatal acute liver failure), hepatitis, reversible increases in hepatic enzymes

<b>Skin and Subcutaneous Tissue Disorders:</b>	Toxic epidermal necrolysis, Stevens-Johnson syndrome, acute generalized exanthematous pustulosis, erythema multiforme, exfoliative dermatitis, leukocytoclastic vasculitis, alopecia, photosensitivity, urticaria
<b>Musculoskeletal and Connective Tissue Disorders:</b>	Arthralgia
<b>Renal and Urinary Disorders:</b>	Urinary incontinence, pollakiuria
<b>Reproductive System and Breast Disorders:</b>	Erectile dysfunction
<b>General Disorders and Administration Site Conditions:</b>	Peripheral edema
<b>Investigations:</b>	Blood creatine phosphokinase increased

---

There is limited information on the use of SPORANOX<sup>®</sup> during pregnancy. Cases of congenital abnormalities including skeletal, genitourinary tract, cardiovascular and ophthalmic malformations as well as chromosomal and multiple malformations have been reported during post-marketing experience. A causal relationship with SPORANOX<sup>®</sup> has not been established. (See CLINICAL PHARMACOLOGY: Special Populations, CONTRAINDICATIONS, WARNINGS, and PRECAUTIONS: Drug Interactions for more information.)

## **OVERDOSAGE**

Itraconazole is not removed by dialysis. In the event of accidental overdosage, supportive measures should be employed. Activated charcoal may be given if considered appropriate.

In general, adverse events reported with overdose have been consistent with adverse drug reactions already listed in this package insert for itraconazole. (See ADVERSE REACTIONS.)

## **DOSAGE AND ADMINISTRATION**

### **Treatment of Oropharyngeal and Esophageal Candidiasis:**

The solution should be vigorously swished in the mouth (10 mL at a time) for several seconds and swallowed.

The recommended dosage of SPORANOX<sup>®</sup> (itraconazole) Oral Solution for oropharyngeal candidiasis is 200 mg (20 mL) daily for 1 to 2 weeks. Clinical signs and symptoms of oropharyngeal candidiasis generally resolve within several days.

For patients with oropharyngeal candidiasis unresponsive/refractory to treatment with fluconazole tablets, the recommended dose is 100 mg (10 mL) b.i.d. For patients responding to therapy, clinical response will be seen in 2 to 4 weeks. Patients may be expected to relapse shortly after discontinuing therapy. Limited data on the safety of long-term use (>6 months) of SPORANOX<sup>®</sup> Oral Solution are available at this time.

The recommended dosage of SPORANOX<sup>®</sup> Oral Solution for esophageal candidiasis is 100 mg (10 mL) daily for a minimum treatment of three weeks. Treatment should continue for 2 weeks following resolution of symptoms. Doses up to 200 mg (20 mL) per day may be used based on medical judgment of the patient's response to therapy.

SPORANOX<sup>®</sup> Oral Solution and SPORANOX<sup>®</sup> Capsules should not be used interchangeably. Patients should be instructed to take SPORANOX<sup>®</sup> Oral Solution without food, if possible. Only SPORANOX<sup>®</sup> Oral Solution has been demonstrated effective for oral and/or esophageal candidiasis.

**Use in Patients with Renal Impairment:**

Limited data are available on the use of oral itraconazole in patients with renal impairment. Caution should be exercised when this drug is administered in this patient population. (See CLINICAL PHARMACOLOGY: Special Populations and PRECAUTIONS.)

**Use in Patients with Hepatic Impairment:**

Limited data are available on the use of oral itraconazole in patients with hepatic impairment. Caution should be exercised when this drug is administered in this patient population. (See CLINICAL PHARMACOLOGY: Special Populations, WARNINGS, and PRECAUTIONS.)

**HOW SUPPLIED**

SPORANOX<sup>®</sup> (itraconazole) Oral Solution is available in 150 mL amber glass bottles (NDC 50458-295-15) containing 10 mg of itraconazole per mL.

Store at or below 25°C (77°F). Do not freeze.

Keep out of reach of children.

Revised: 04/2015

©Janssen Pharmaceuticals, Inc 2003

Product of Ireland

Manufactured by:

Janssen Pharmaceutica NV..

Beerse, Belgium

Manufactured for:

Janssen Pharmaceuticals, Inc.

Titusville, NJ 08560

## REFERENCES

1. Clinical and Laboratory Standards Institute (CLSI). *Reference Method for Broth Dilution Antifungal Susceptibility Testing of Yeasts; Approved Standard-Third Edition*. CLSI document M27-A3. Clinical and Laboratory Standards Institute, 940 West Valley Road, Suite 1400, Wayne, Pennsylvania 19087-1898, USA, 2008.
2. Clinical and Laboratory Standards Institute (CLSI). *Reference Method for Broth Dilution Antifungal Susceptibility Testing of Yeasts; Fourth Informational Supplement*. CLSI document M27-S4. Clinical and Laboratory Standards Institute, 940 West Valley Road, Suite 2500, Wayne, Pennsylvania 19087 USA, 2012.