

200%




100%



Variable Data Coding Area: use a color that provides the greatest contrast for vision inspection system and human readable.


Variable Data Coding Color: (b) (4)



The Bayer Cross is Equal to the the "L" in Aleve
Current Size: .292"
Current "L" Size: .292"
Cross Meets Ratio Requirements of 7mm (.2755") or larger
 1/3rd of Cross required Clear Space (.292) = .097"
 1/6th of PDP Width (1.656") = .276"



(b) (4)



Actual Size 100%

(b) (4)

CATEGORY	PRIMARY PACKAGING REQUIRED	SECONDARY PACKAGING REQUIRED	ACTUAL SIZE IN FILE
Drug Facts Title:	N/A	>8.0 pt	-
Drug Facts	N/A	8 pt	-
(continued):	N/A	8 pt	-
Headings:	N/A	8 pt	4.5 pt
Subheadings:	N/A	6 pt	4.5 pt
Text:	N/A	6 pt	4.5 pt
Bullets:	N/A	5 pt (no larger)	4.5 pt
Leading:	N/A	+0.5 pt	4.75 pt
Characters Per Inch:	N/A	No more than 39	>39

Drug Facts (continued)

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use naproxen sodium during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not take more than directed
- the smallest effective dose should be used
- drink a full glass of water with each dose

Adults and children 12 years and older

- take 1 tablet every 8 to 12 hours while symptoms last
- for the first dose you may take 2 tablets within the first hour
- do not exceed 2 tablets in any 8- to 12-hour period
- do not exceed 3 tablets in a 24-hour period

Children under 12 years

- ask a doctor

Other information

- each tablet contains sodium 20 mg
- store at 20-25 °C (68-77°F). Avoid high humidity and excessive heat above 40°C (104°F).

Inactive ingredients FD&C blue #2 lake, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, povidone, talc, titanium dioxide

Questions or comments?
1-800-385-0688 (Mon-Fri 9AM-5PM EST)

3
25866151256
7

Bayer, the Bayer Cross, Aleve and All Day Strong are registered trademarks of Bayer.

Drug Facts (continued)

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- the stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have had high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic
- you have problems or serious side effects from taking pain relievers or fever reducers

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs
- you experience any of the following signs of stomach bleeding:

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
- red taint
- vomit blood
- have bloody or black stools
- you have stomach pain that does not get better
- chest pain
- trouble breathing
- weakness in one part or side of body
- slurred speech
- big swelling
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- you have difficulty swallowing
- it feels like the pill is stuck in your throat
- redness or swelling is present in the painful area
- any new symptoms appear

Do not use if carton is open or if foil seal imprinted with "SAFETY SQUEASE" on bottle opening is missing or broken.

Dist. by: Bayer HealthCare LLC
Whisperpany, NJ 07981

Lot _____
Exp _____

Made in _____

Bayer

XXXXXXX
0000

Drug Facts

Active Ingredient (in each tablet)
Naproxen sodium 220 mg (naproxen 200 mg) (NSAID)* ----Pain reliever/fever reducer
*nonsteroidal anti-inflammatory drug

Purposes

- temporarily relieves minor aches and pains due to:
- minor pain of arthritis
- muscular aches
- menstrual cramps
- toothache
- the common cold
- temporarily reduces fever

Warnings

Allergy alert: Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- asthma (wheezing)
- skin redness/itching
- swelling
- acid swelling
- stomach pain
- stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

50 FREE TABLETS WITH THIS PURCHASE

ALL DAY STRONG®

ALEVE®


naproxen sodium tablets, 220 mg (NSAID)
Pain reliever/fever reducer

150 TABLETS

Actual Size 100%


(b) (4)

CATEGORY	PRIMARY PACKAGING REQUIRED	SECONDARY PACKAGING REQUIRED	ACTUAL SIZE IN FILE
Drug Facts Title:	N/A	>8.0 pt	8.1 pt
Drug Facts	N/A	8 pt	8 pt
(continued):	N/A	8 pt	8 pt
Headings:	N/A	8 pt	8 pt
Subheadings:	N/A	6 pt	6 pt
Text:	N/A	6 pt	6 pt
Bullets:	N/A	5 pt (no larger)	5 pt
Leading:	N/A	+0.5 pt	6.5-6.9 pt
Characters Per Inch:	N/A	No more than 39	≤ 39



Actual Size 100%


This label may not be the latest approved by FDA. For labeling information, please visit <https://www.fda.gov>



Actual Size 100%

(b) (4)

CATEGORY	PRIMARY PACKAGING REQUIRED	SECONDARY PACKAGING REQUIRED	ACTUAL SIZE IN FILE
Drug Facts Title:	N/A	>8.0 pt	-
Drug Facts	N/A	8 pt	-
(continued):	N/A	8 pt	-
Headings:	N/A	8 pt	4.5 pt
Subheadings:	N/A	6 pt	4.5 pt
Text:	N/A	6 pt	4.5 pt
Bullets:	N/A	5 pt (no larger)	4.5 pt
Leading:	N/A	+0.5 pt	5.0 pt
Characters Per Inch:	N/A	No more than 39	>39



Actual Size 100%

(b) (4)

CATEGORY	PRIMARY PACKAGING REQUIRED	SECONDARY PACKAGING REQUIRED	ACTUAL SIZE IN FILE
Drug Facts Title:	N/A	>8.0 pt	8.1 pt
Drug Facts (continued):	N/A	8 pt	8 pt
Headings:	N/A	8 pt	8 pt
Subheadings:	N/A	6 pt	6 pt
Text:	N/A	6 pt	6 pt
Bullets:	N/A	5 pt (no larger)	5 pt
Leading:	N/A	+0.5 pt	6.5-6.9 pt
Characters Per Inch:	N/A	No more than 39	≤ 39



0000
XXXXXXX

Drug Facts

Active ingredient (in each tablet)
Naproxen sodium 220 mg (naproxen 200 mg) (NSAID)* Pain reliever/fever reducer
*nonsteroidal anti-inflammatory drug

Uses
temporarily relieves minor aches and pains due to:

- minor pain of arthritis
- muscular aches
- menstrual cramps
- toothache
- temporarily reduces fever

Warnings
Allergy alert: Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- asthma (wheezing)
- skin reddening
- blisters
- facial swelling
- shock
- rash

 If an allergic reaction occurs, stop use and seek medical help right away.
Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.
Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Drug Facts (continued)

Ask a doctor before use if

- the stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have had high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic
- you have problems or serious side effects from taking pain relievers or fever reducers

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
- have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
 - chest pain
 - trouble breathing
 - weakness in one part or side of body
 - slurred speech
 - leg swelling
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- you have difficulty swallowing
- it feels like the pill is stuck in your throat
- redness or swelling is present in the painful area
- any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use naproxen sodium during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.
Directions

- do not take more than directed
- the smallest effective dose should be used
- drink a full glass of water with each dose

Drug Facts (continued)

Adults and children 12 years and older

- take 1 tablet every 8 to 12 hours while symptoms last
- for the first dose you may take 2 tablets within the first hour
- do not exceed 2 tablets in any 8- to 12-hour period
- do not exceed 3 tablets in a 24-hour period
- ask a doctor

Children under 12 years

- ask a doctor

Other information

- each tablet contains: sodium 20 mg
- store at 20-25°C (68-77°F). Avoid high humidity and excessive heat above 40°C (104°F).

Inactive ingredients FD&C blue #2 lake, hydroxypropyl methylcellulose, polyethylene glycol, povidone, talc, titanium dioxide
Questions or comments?
 1-800-395-0689 (Mon - Fri 9AM - 5PM EST)

Bayer, the Bayer Cross, Aleve and All Day Strong are registered trademarks of Bayer.
 Dist. by:
 Bayer HealthCare LLC
 Whippany, NJ 07981
 Bayer

ALL DAY STRONG®
ALEVE®

BUY THIS TWIN PACK & SAVE \$5
 OFF MSRP OF TWO INDIVIDUAL 100-COUNT SIZE BOTTLES

ALL DAY STRONG®
ALEVE®

naproxen sodium tablets, 220 mg (NSAID)
 Pain reliever/fever reducer

STRENGTH TO LAST 12 HOURS

200 TABLETS TOTAL
 (2 BOTTLES X 100 TABLETS EACH)

STRENGTH TO LAST 12 HOURS



0 00000 00000 0

LOT/EXP MADE IN

ALL DAY STRONG®
ALEVE®

STRENGTH TO LAST 12 HOURS



0000
XXXXXXX

Do not use if carton is open or if foil seal imprinted with "SAFETY SQUEASE" on bottle opening is missing or broken.

Safety Squeeze is a registered trademark of The Procter & Gamble Company. Used with permission.

(b) (4)

CATEGORY	PRIMARY PACKAGING REQUIRED	SECONDARY PACKAGING REQUIRED	ACTUAL SIZE IN FILE
Drug Facts Title:	N/A	>8.0 pt	9 pt
Drug Facts	N/A	8 pt	8 pt
(continued):	N/A	8 pt	8 pt
Headings:	N/A	8 pt	8 pt
Subheadings:	N/A	6 pt	6 pt
Text:	N/A	6 pt	6 pt
Bullets:	N/A	5 pt (no larger)	5 pt
Leading:	N/A	+0.5 pt	6.5-6.9 pt
Characters Per Inch:	N/A	No more than 39	≤ 39

200%




100%



Variable Data Coding Area: use a color that provides the greatest contrast for vision inspection system and human readable.


Variable Data Coding Color: (b) (4)



The Bayer Cross is Equal to the the "L" in Aleve
 Current Size: .292"
 Current "L" Size: .292"
 Cross Meets Ratio Requirements of 7mm (.2755") or larger
 1/3rd of Cross required Clear Space (.292) = .097"
 1/6th of PDP Width (1.656") = .276"



(b) (4)



Actual Size 100%

(b) (4)

CATEGORY	PRIMARY PACKAGING REQUIRED	SECONDARY PACKAGING REQUIRED	ACTUAL SIZE IN FILE
Drug Facts Title:	N/A	>8.0 pt	-
Drug Facts	N/A	8 pt	-
(continued):	N/A	8 pt	-
Headings:	N/A	8 pt	4.5 pt
Subheadings:	N/A	6 pt	4.5 pt
Text:	N/A	6 pt	4.5 pt
Bullets:	N/A	5 pt (no larger)	4.5 pt
Leading:	N/A	+0.5 pt	4.75 pt
Characters Per Inch:	N/A	No more than 39	>39

Bayer, the Bayer Cross, Aleve and All Day Strong are registered trademarks of Bayer.

Drug Facts (continued)

Other information

- each caplet contains sodium 20 mg
- store at 20-25°C (68-77°F). Avoid high humidity and excessive heat above 40°C (104°F).

Inactive ingredients: FD&C Blue #2 lake, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, povidone, talc, titanium dioxide

Questions or comments?
1-800-985-0689 (Mon-Fri 9AM-5PM EST)

Bayer, the Bayer Cross, Aleve and All Day Strong are registered trademarks of Bayer.

Drug Facts (continued)

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- the stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver disease, kidney disease, asthma, or had a stroke
- you are taking a diuretic
- you have problems or serious side effects from taking pain relievers or fever reducers

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
- have stomach pain that does not get better
- chest pain
- trouble breathing
- weakness in one part or side of body
- slurred speech
- itchy swelling
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- you have difficulty swallowing
- it feels like the pill is stuck in your throat
- redness or swelling is present in the painful area
- any new symptoms appear

Bayer, the Bayer Cross, Aleve and All Day Strong are registered trademarks of Bayer.

Drug Facts

Active ingredient *Purposes*

Naproxen sodium 220 mg (naproxen 200 mg (NSAID)*, ----Pain reliever/fever reducer/nonsteroidal anti-inflammatory drug)

Uses

- temporarily relieves minor aches and pains due to:
 - minor pain of arthritis
 - muscular aches
 - menstrual cramps
 - toothache
- temporarily reduces fever

Warnings

Allergy alert: Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- asthma (wheezing)
- skin redness
- blisters
- rash

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

XXXXXXXX
0000

Bayer, the Bayer Cross, Aleve and All Day Strong are registered trademarks of Bayer.

Do not use if carton is open or if foil seal imprinted with "SAFETY-SQUEASE" on bottle opening is missing or broken.

Dist. by: Bayer HealthCare LLC
Whispery, NJ 07881

Lot
Exp

Bayer
Made in

50 FREE CAPLETS WITH THIS PURCHASE

ALL DAY STRONG®

ALEVE®

naproxen sodium tablets, 220 mg (NSAID)
Pain reliever/fever reducer


150 CAPLETS
100 CAPLETS
CAPSULE-SHAPED TABLETS

STRENGTH TO LAST 12 HOURS

Actual Size 100%


CATEGORY	PRIMARY PACKAGING REQUIRED	SECONDARY PACKAGING REQUIRED	ACTUAL SIZE IN FILE
Drug Facts Title:	N/A	>8.0 pt	8.1 pt
Drug Facts (continued):	N/A	8 pt	8 pt
Headings:	N/A	8 pt	8 pt
Subheadings:	N/A	6 pt	6 pt
Text:	N/A	6 pt	6 pt
Bullets:	N/A	5 pt (no larger)	5 pt
Leading:	N/A	+0.5 pt	6.5-6.9 pt
Characters Per Inch:	N/A	No more than 39	≤ 39

200%



The Bayer Cross is Equal to the "A" in Aleve
 Current Size: .355"
 Current "A" Size: .355"
 Cross Meets Ratio Requirements of 7mm (.2755") or larger
 1/3rd of Cross required Clear Space (.355) = .118"
 1/6th of PDP Height (2.134") = .355"





Actual Size 100%

DRUG FACTS INFO.			
CATEGORY	PRIMARY PACKAGING REQUIRED	SECONDARY PACKAGING REQUIRED	ACTUAL SIZE IN FILE
Drug Facts Title:	N/A	>8.0 pt	8.1 pt
Drug Facts	N/A	8 pt	8 pt
(continued):	N/A	8 pt	8 pt
Headings:	N/A	8 pt	8 pt
Subheadings:	N/A	6 pt	6 pt
Text:	N/A	6 pt	6 pt
Bullets:	N/A	5 pt (no larger)	5 pt
Leading:	N/A	+0.5 pt	6.5-6.7 pt
Characters Per Inch:	N/A	No more than 39	≤ 39

Drug Facts (continued)

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use naproxen sodium during the last 3 months of pregnancy, unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not take more than directed
- the smallest effective dose should be used
- drink a full glass of water with each dose

Adults and children 12 years and older

- take 1 caplet every 8 to 12 hours while symptoms last
- for the first dose you may take 2 caplets within the first hour
- do not exceed 2 caplets in any 8-to-12-hour period
- do not exceed 3 caplets in a 24-hour period
- ask a doctor

Children under 12 years

Other Information

- each caplet contains: sodium 20 mg
- store at 20-25°C (68-77°F). Avoid high humidity and excessive heat above 40°C (104°F).

Inactive Ingredients FDAC blue #2 lake, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polydioxane, talc, titanium dioxide

Questions or comments? 1-800-368-6999 (Non-PHARM - RPM EST)

Bayer

Drug Facts

Active Ingredient (in each caplet) Purposes

Naproxen sodium 220 mg (naproxen 200 mg) (NSAID)*.....Pain reliever/fever reducer

* Nonsteroidal anti-inflammatory drug

Division Naprosyn®
Manufactured by
Bayer HealthCare under
the following: 3-25866-53647 1

LOT: 12345678
NDC: 31131-01-01

Drug Facts (continued)

Uses

- temporarily relieves minor aches and pains due to:
 - minor pain of arthritis
 - muscular aches
 - backache
 - menstrual cramps
 - headache
 - toothache
 - the common cold
- temporarily reduces fever

Warnings

Allergy alert: Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin.

Symptoms may include:

- hives
- swelling of the face, lips, tongue, or throat
- skin redness or rash
- blistering
- difficulty breathing

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- the stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver disease, kidney disease, asthma, or had a stroke
- you are taking a diuretic
- you have problems or serious side effects from taking pain relievers or fever reducers

Ask a doctor or pharmacist before use if you are


- under a doctor's care for any serious condition
- taking any other drug

When using this product


- take with food or milk; if stomach upset occurs

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - red, black, or tarry stools
 - vomiting blood
 - have bloody or black stools
 - have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
 - chest pain
 - trouble breathing
 - weakness in one part or side of body
 - sudden speech
 - leg swelling
 - pain gets worse or lasts more than 10 days
 - fever gets worse or lasts more than 3 days
 - you have difficulty swallowing
 - it feels like the pill is stuck in your throat
 - redness or swelling is present in the painful area
 - any new symptoms appear



Actual Size 100%



Actual Size 100%

DRUG FACTS INFO.			
CATEGORY	PRIMARY PACKAGING REQUIRED	SECONDARY PACKAGING REQUIRED	ACTUAL SIZE IN FILE
Drug Facts Title:	N/A	>8.0 pt	8.1 pt
Drug Facts	N/A	8 pt	8 pt
(continued):	N/A	8 pt	8 pt
Headings:	N/A	8 pt	8 pt
Subheadings:	N/A	6 pt	6 pt
Text:	N/A	6 pt	6 pt
Bullets:	N/A	5 pt (no larger)	5 pt
Leading:	N/A	+0.5 pt	6.5-6.7 pt
Characters Per Inch:	N/A	No more than 39	≤ 39

Drug Facts (continued)

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use naproxen sodium during the last 3 months of pregnancy, unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not take more than directed
- the smallest effective dose should be used
- drink a full glass of water with each dose

Adults and children 12 years and older

- take 1 caplet every 8 to 12 hours while symptoms last
- for the first dose you may take 2 caplets within the first hour
- do not exceed 2 caplets in any 8-to-12-hour period
- do not exceed 3 caplets in a 24-hour period
- ask a doctor

Children under 12 years

Other Information

- each caplet contains: sodium 20 mg
- store at 20-25°C (68-77°F). Avoid high humidity and excessive heat above 40°C (104°F).

Inactive Ingredients FDAC blue #2 lake, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polydioxane, talc, titanium dioxide

Questions or comments? 1-800-368-6999 (Non-PHARM - RPM EST)

Bayer

Drug Facts

Active Ingredient (in each caplet) Purposes

Naproxen sodium 220 mg (naproxen 200 mg) (NSAID)*.....Pain reliever/fever reducer

* Nonsteroidal anti-inflammatory drug

Division Naproxen Sodium 220 mg Caplets (NSAID) (Pain Reliever/ Fever Reducer) is a registered trademark of Bayer HealthCare under the following information: 3-25866-53647 1

LOT: N/A

Drug Facts (continued)

Uses

- temporarily relieves minor aches and pains due to:
 - minor pain of arthritis
 - muscular aches
 - backache
 - menstrual cramps
 - headache
 - toothache
 - the common cold
- temporarily reduces fever

Warnings

Allergy alert: Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin.

Symptoms may include:

- hives
- swelling of the face, lips, tongue, or throat
- skin redness or rash
- blistering
- difficulty breathing

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- the stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver disease, kidney disease, asthma, or had a stroke
- you are taking a diuretic
- you have problems or serious side effects from taking pain relievers or fever reducers

Ask a doctor or pharmacist before use if you are


- under a doctor's care for any serious condition
- taking any other drug

When using this product

- take with food or milk; if stomach upset occurs

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - red, black, or tarry stools
 - vomiting blood
 - have bloody or black stools
 - have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
 - chest pain
 - trouble breathing
 - weakness in one part or side of body
 - sudden speech
 - leg swelling
 - pain gets worse or lasts more than 10 days
 - fever gets worse or lasts more than 3 days
 - you have difficulty swallowing
 - it feels like the pill is stuck in your throat
 - redness or swelling is present in the painful area
 - any new symptoms appear



Actual Size 100%

(b) (4)

CATEGORY	PRIMARY PACKAGING REQUIRED	SECONDARY PACKAGING REQUIRED	ACTUAL SIZE IN FILE
Drug Facts Title:	N/A	>8.0 pt	8.1 pt
Drug Facts (continued):	N/A	8 pt	8 pt
Headings:	N/A	8 pt	8 pt
Subheadings:	N/A	6 pt	6 pt
Text:	N/A	6 pt	6 pt
Bullets:	N/A	5 pt (no larger)	5 pt
Leading:	N/A	+0.5 pt	6.5-7 pt
Characters Per Inch:	N/A	No more than 39	≤ 39



0000
XXXXXXX

Drug Facts

Active ingredient (in each caplet)
Naproxen sodium 220 mg (naproxen 200 mg) (NSAID)* Pain reliever/fever reducer
*nonsteroidal anti-inflammatory drug

Uses

- temporarily relieves minor aches and pains due to:
 - minor pain of arthritis
 - muscular aches
 - menstrual cramps
 - toothache
- temporarily reduces fever

Warnings

Allergy alert: Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- asthma (wheezing)
- skin reddening
- blisters
- facial swelling
- shock
- rash

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Drug Facts (continued)

Ask a doctor before use if

- the stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have had high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic
- you have problems or serious side effects from taking pain relievers or fever reducers

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
 - chest pain
 - trouble breathing
 - weakness in one part or side of body
 - slurred speech
 - leg swelling
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- you have difficulty swallowing
- it feels like the pill is stuck in your throat
- redness or swelling is present in the painful area
- any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use naproxen sodium during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not take more than directed
- the smallest effective dose should be used
- drink a full glass of water with each dose

Drug Facts (continued)

Adults and children 12 years and older

- take 1 caplet every 8 to 12 hours while symptoms last
- for the first dose you may take 2 caplets within the first hour
- do not exceed 2 caplets in any 8- to 12-hour period
- do not exceed 3 caplets in a 24-hour period
- ask a doctor

Children under 12 years

- ask a doctor

Other information

- each caplet contains: sodium 20 mg
- store at 20-25°C (68-77°F). Avoid high humidity and excessive heat above 40°C (104°F).

Inactive ingredients FD&C blue #2 lake, hydroxypropyl methylcellulose, polyethylene glycol, povidone, talc, titanium dioxide

Questions or comments?
1-800-395-0689 (Mon - Fri 9AM - 5PM EST)

Bayer, the Bayer Cross, Aleve and All Day Strong are registered trademarks of Bayer.

Dist. by:
Bayer HealthCare LLC
Whippany, NJ 07981

Bayer



0000
XXXXXXX



0 000000 000000 0

LOT/EXP MADE IN

ALL DAY STRONG®
ALEVE

BUY THIS TWIN PACK & SAVE \$5
OFF MSRP OF TWO INDIVIDUAL 100-COUNT SIZE BOTTLES

ALL DAY STRONG®
ALEVE

naproxen sodium tablets, 220 mg (NSAID)
Pain reliever/fever reducer

STRENGTH TO LAST 12 HOURS

200 CAPLETS* TOTAL
(2 BOTTLES x 100 CAPLETS* EACH)
*CAPSULE-SHAPED TABLETS

STRENGTH TO LAST 12 HOURS

Do not use if carton is open or if foil seal imprinted with "SAFETY SQUEASE" on bottle opening is missing or broken.

Safety Squeeze is a registered trademark of The Procter & Gamble Company. Used with permission.

STRENGTH TO LAST 12 HOURS

(b) (4)

CATEGORY	PRIMARY PACKAGING REQUIRED	SECONDARY PACKAGING REQUIRED	ACTUAL SIZE IN FILE
Drug Facts Title:	N/A	>8.0 pt	9 pt
Drug Facts	N/A	8 pt	8 pt
(continued):	N/A	8 pt	8 pt
Headings:	N/A	8 pt	8 pt
Subheadings:	N/A	6 pt	6 pt
Text:	N/A	6 pt	6 pt
Bullets:	N/A	5 pt (no larger)	5 pt
Leading:	N/A	+0.5 pt	6.5-6.9 pt
Characters Per Inch:	N/A	No more than 39	≤ 39

This is a representation of an electronic record that was signed electronically and this page is the manifestation of the electronic signature.

/s/

KAREN M MAHONEY
03/29/2018