

20 FREE TABLETS WITH THIS PURCHASE

ALL DAY STRONG

ALEVE

naproxen sodium tablets, 220 mg (NSAID)
Pain reliever/fever reducer

STRENGTH TO LAST 12 HOURS

ACTUAL SIZE 110 TABLETS

LOT EXP MADE IN

LOT & EXP CODE AREA

Drug Facts (continued)

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- the stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic
- you have problems or serious side effects from taking pain relievers or fever reducers

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because naproxen may decrease this benefit of aspirin
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - have bloody or black stools
 - have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
 - chest pain
 - trouble breathing
 - weakness in one part or side of body
 - slurred speech
 - leg swelling
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- you have difficulty swallowing
- it feels like the pill is stuck in your throat
- redness or swelling is present in the painful area
- any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use naproxen sodium during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Drug Facts (continued)

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not take more than directed
- the smallest effective dose should be used
- drink a full glass of water with each dose

Adults and children 12 years and older

- take 1 tablet every 8 to 12 hours while symptoms last
- for the first dose you may take 2 tablets within the first hour
- do not exceed 2 tablets in any 8- to 12-hour period
- do not exceed 3 tablets in a 24-hour period

Children under 12 years

- ask a doctor

Other information

- each tablet contains: sodium 20 mg
- store at 20-25°C (68-77°F). Avoid high humidity and excessive heat above 40°C (104°F).

Inactive ingredients: FD&C blue #2 lake, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, povidone, talc, titanium dioxide

Questions or comments?
1-800-995-0689 (Mon - Fri 9AM - 5PM EST)

Bayer, the Bayer Cross, Aleve and All Day Strong are registered trademarks of Bayer.
Dist. by: Bayer HealthCare LLC
Whispery, NJ 07981

Drug Facts

Active ingredient (in each tablet)

Naproxen sodium 220 mg (naproxen 200 mg) (NSAID)¹..... Pain reliever/fever reducer
¹nonsteroidal anti-inflammatory drug

Purposes

- temporarily relieves minor aches and pains due to:
 - muscular aches
 - backache
 - headache
 - the common cold
 - temporarily reduces fever
- minor pain of arthritis
- menstrual cramps
- toothache

Uses

- temporarily relieves minor aches and pains due to:
 - muscular aches
 - backache
 - headache
 - the common cold
 - temporarily reduces fever

Warnings

Allergy alert: Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- swelling (wheezing)
- skin redness
- itching
- facial swelling
- choke
- rash
- bleisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:


- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Actual Size 100%


FDA00379-1

DRUG FACTS INFO (updated 5/18/18)		
CATEGORY	REQUIRED	ACTUAL IN FILE
Drug Facts Title	>8 pt.**	8.1 pt.
Drug Facts Continued	8 pt.	8 pt.
Headings [†]	8 pt bold italic [†]	8 pt.
Subheadings	≥6 pt.	6 pt.
Body Text	≥6 pt.	6 pt.
Bullets	5 pt.	12 pt.
Loading	≥0.5 pt.	0.5 pt.
Headlines	0.5 pt. rule	0.5 pt.
Box and Barlines [†]	>0.5 pt. rule	2.5 pt.
Characters Per Inch	No more than 30	≤30

[†] Headings are to be at least 2 pts larger than Body Text.
^{**} Drug Facts Title to be larger than all other type size used in DFP.
^{††} FDA Recommends 2.5 pt. rule.



200%



Actual Size 100%

FDA00380-1



DRUG FACTS INFO: (Modified Format)		
CATEGORY	REQUIRED	ACTUAL ON FILE
Drug Facts Title	>7.0 pt.	-
Drug Facts Continued	7 pt.	-
Headings*	7 pt- bold italic *	4.5 pt.
Subheadings	≥6 pt.	4.5 pt.
Body Text	≥6 pt.	4.5 pt.
Bullets	5 pt.	6 pt.
Leading	≥0 pt.	4.5 pt.
Hairlines	0.5 pt rule	-
Box and Barline†	>0.5 pt rule	-
Characters Per Inch:	No more than 99	> 99

* Headings are to be at least 2pts larger than Body Text
 ** Drug Facts Title to be larger than all other type size used in DFP
 † Box Barline may be omitted and color contrast used to highlight Drug Facts Info

20 FREE CAPLETS WITH THIS PURCHASE

ALL DAY STRONG[®]
ALEVE
naproxen sodium tablets, 220 mg (NSAID)
Pain reliever/fever reducer

STRENGTH TO LAST 12 HOURS

110 ACTUAL SIZE CAPLETS

LOT EXP MADE IN

Drug Facts (continued)
Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.
Do not use:
• if you have ever had an allergic reaction to any other pain reliever/fever reducer
• right before or after heart surgery
Ask a doctor before use if:
• the stomach bleeding warning applies to you
• you have a history of stomach problems, such as heartburn
• you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
• you are taking a diuretic
• you have problems or serious side effects from taking pain relievers or fever reducers
Ask a doctor or pharmacist before use if you are:
• under a doctor's care for any serious condition
• taking aspirin for heart attack or stroke, because naproxen may decrease this benefit of aspirin
• taking any other drug
When using this product:
• take with food or milk if stomach upset occurs
Stop use and ask a doctor if:
• you experience any of the following signs of stomach bleeding:
• feel faint • vomit blood
• have bloody or black stools
• have stomach pain that does not get better
• you have symptoms of heart problems or stroke:
• chest pain • trouble breathing
• weakness in one part or side of body
• slurred speech • leg swelling
• pain gets worse or lasts more than 10 days
• fever gets worse or lasts more than 3 days
• you have difficulty swallowing
• it feels like the pill is stuck in your throat
• redness or swelling is present in the painful area
• any new symptoms appear
If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use naproxen sodium during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Drug Facts (continued)
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.
Directions
• do not take more than directed
• the smallest effective dose should be used
• drink a full glass of water with each dose
Adults and children 12 years and older
• take 1 caplet every 8 to 12 hours while symptoms last
• for the first dose you may take 2 caplets within the first hour
• do not exceed 2 caplets in any 8- to 12-hour period
• do not exceed 3 caplets in a 24-hour period
Children under 12 years
• ask a doctor
Other information
• each caplet contains: sodium 20 mg
• store at 20-25°C (68-77°F). Avoid high humidity and excessive heat above 40°C (104°F).
Inactive ingredients: FD&C blue #2 lake, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, povidone, talc, titanium dioxide
Questions or comments?
1-800-995-0689 (Mon-Fri 9AM-5PM EST)
Bayer, the Bayer Cross, Aleve and All Day Strong are registered trademarks of Bayer.
Dist. by:
Bayer HealthCare LLC
Whispering, NJ 07981


Drug Facts
Active ingredient (in each caplet)
Naproxen sodium 220 mg (NSAID)
Naproxen 200 mg (NSAID)
Purposes
• Pain reliever/fever reducer
• nonsteroidal anti-inflammatory drug
Uses
• temporarily relieves minor aches and pains due to:
• minor pain of arthritis
• muscular aches • backache
• menstrual cramps • headache
• toothache
• the common cold
• temporarily reduces fever
Warnings
Allergy alert: Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:
• hives • facial swelling
• asthma (wheezing) • shock
• skin redness • rash
• blisters
If an allergic reaction occurs, stop use and seek medical help right away.
Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:
• are age 60 or older
• have had stomach ulcers or bleeding problems
• take a blood thinning (anticoagulant) or steroid drug
• take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
• have 3 or more alcoholic drinks every day while using this product
• take more or for a longer time than directed

Actual Size 100%

FDA00375-2

DRUG FACTS INFO (updated 5/18/18)		
CATEGORY	REQUIRED	ACTUAL IN FILE
Drug Facts Title	>8 pt.**	9 pt.
Drug Facts Continued	8 pt.	8 pt.
Headings*	8 pt. bold italic*	8 pt.
Subheadings	≥6 pt.	6 pt.
Body Text	≥6 pt.	6 pt.
Bullets	6 pt.	12 pt.
Leading	≥0.5 pt.	0.5 pt.
Headlines	0.5 pt. rule	0.5 pt.
Box and Barlines†	>0.5 pt. rule	2.5 pt.
Characters Per Inch	No more than 29	≤30

* Headings are to be at least 20% larger than Body Text.
** Drug Facts Title to be larger than all other type size used in DFP.
† FDA Recommends 2.5pt. rule.



200%



Actual Size 100%

FDA00376-2



DRUG FACTS INFO: Modified Format - updated 5/18/18		
CATEGORY	REQUIRED	ACTUAL IN FILE
Drug Facts Title	>7.0 pt	-
Drug Facts Continued	7 pt	-
Headings*	7 pt- bold italic *	-
Subheadings	≥6 pt	4.5 pt
Body Text	≥6 pt	4.5 pt
Bullets	5 pt	8 pt
Leading	≥0 pt	4.75 pt
Headlines	0.5 pt rule	-
Box and Barlines†	>0.5 pt rule	-
Characters Per Inch:	No more than 39	≥ 39


* Headings are to be at least 2pts larger than Body Text.
** Drug Facts Title to be larger than all other type size used in DFP.
† Box Barline may be omitted and color contrast used to highlight Drug Facts Info.

200%



Actual Size 100%

FDA00373-2



Actual Size 100%

FDA00174-1

DRUG FACTS INFO (updated 5/18/10)		
CATEGORY	REQUIRED	ACTUAL IN FILE
Drug Facts Title	>8 pt.**	8.1 pt.
Drug Facts Continued	8 pt.	8 pt.
Headings*	8 pt.-bold italic*	8 pt.
Subheadings	>6 pt.	6 pt.
Body Text	>6 pt.	6 pt.
Bullets	5 pt.	12 pt.
Leading	≥0.5 pt.	0.5 - 0.7 pt.
Hairlines	0.5 pt. rule	0.5 pt.
Box and Barline†	>0.5 pt. rule	2.5 pt.
Characters Per Inch:	No more than 30	≤30

* Headings are to be at least 5 pts. larger than Body Text.
 ** Drug Facts Title to be larger than all other text used in DFP.
 † FDA Recommends 2.5pt. rule

Drug Facts (continued)

- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- you have difficulty swallowing
- it feels like the pill is stuck in your throat
- redness or swelling is present in the painful area
- any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use naproxen when during the last 6 months of pregnancy unless advised by your health care provider. Naproxen may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- **do not take more than directed**
- **do not exceed the dose listed below**
- **do not exceed 2 capsules in any 8- to 12-hour period**
- **do not exceed 3 capsules in a 24-hour period**
- ask a doctor

Children under 12 years

Other Information

- each capsule contains: sodium, 20 mg
- store at 20-25°C (68-77°F). Avoid high humidity and excessive heat above 40°C (104°F).

Inactive Ingredients: FDAC blue 42 lake, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, povidone, talc, titanium dioxide

Questions or comments? 1-800-858-0880 (ask for NA-0153)

Bayer

Drug Facts

Active Ingredient (in each capsule): Naproxen sodium 250 mg (NSAID)
Purpose: Pain reliever/fever reducer
Warnings: Naproxen sodium is a nonsteroidal anti-inflammatory drug.

Do not use if you have had an allergic reaction to any other NSAID or aspirin.
 Do not use if you have had an allergic reaction to any other NSAID or aspirin.
 Do not use if you have had an allergic reaction to any other NSAID or aspirin.

Drug Facts (continued)

Uses

- temporarily relieves minor aches and pains due to:
 - minor pain of arthritis
 - muscular aches
 - headache
 - menstrual cramps
 - toothache
- temporarily reduces fever

Warnings

Allergy alert: Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- skin redness/itching
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are aged 65 or older
- take aspirin or other drugs containing aspirin
- take a blood thinner (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- use more or for a longer time than directed

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These risks may be higher if you use NSAIDs for a long time. The longer you use NSAIDs, the more serious the risk. The longer you use NSAIDs, the more serious the risk. The longer you use NSAIDs, the more serious the risk.

Do not use

- if you have ever had an allergic reaction to any other NSAID or aspirin
- if you have ever had an allergic reaction to any other NSAID or aspirin
- if you have ever had an allergic reaction to any other NSAID or aspirin

Ask a doctor before use if

- the stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic
- you have problems or serious side effects from taking pain relievers or fever reducers

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because naproxen may decrease the benefit of aspirin
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
- have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
 - chest pain
 - trouble breathing
 - weakness in one part of your body
 - slurred speech
 - leg swelling



Actual Size 100%

FDA00358-2

DRUG FACTS INFO (Updated 5/18/19)		
CATEGORY	REQUIRED	ACTUAL IN FILE
Drug Facts Title	>8 pt.**	9 pt.
Drug Facts Continued	8 pt.	8 pt.
Headings*	8 pt bold italic*	8 pt.
Subheadings	≥6 pt.	6 pt.
Body Text	≥6 pt.	6 pt.
Bullets	5 pt.	12 pt.
Leading	≥0.5 pt.	0.5 pt.
Headlines	0.5 pt. rule	0.5 pt.
Box and Barlines†	>0.5 pt. rule	2.5 pt.
Characters Per Inch	No more than 30	≤30

* Headings are to be at least 2pts larger than Body Text.
** Drug Facts Title to be larger than all other type size used in DFP.
† FDA Recommendation: 2.5pt rule.



200%



Actual Size 100%

FDA00359-2

See Carton
for Full Information

Warnings: Allergy alert: Naproxen sodium may cause a severe allergic reaction which may include: hives, facial swelling, asthma (wheezing) or shock. **Stomach bleeding warning:** This product contains a nonsteroidal anti-inflammatory drug (NSAID), which may cause severe stomach bleeding. The chance is higher if you: ● are age 60 or older ● have had stomach ulcers or bleeding problems ● take a blood thinning (anticoagulant) or steroid drug ● take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ● have 3 or more alcoholic drinks every day while using this product ● take more or for a longer time than directed. **Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed. **Do not use** if you have ever had an allergic reaction to any other pain reliever/fever reducer. **Stop use and ask a doctor** if you experience any of the following signs of stomach bleeding (stomach pain, vomit blood, have bloody or black stools, have stomach pain that does not get better), have symptoms of heart problems or stroke (chest pain, trouble breathing, weakness in one part or side of body, slurred speech, leg swelling), or if pain or fever gets worse. **If pregnant or breast-feeding**, ask a health professional before use. **Directions** Adults and children 12 years and older: take 1 tablet with water every 8 to 12 hours while symptoms last. For the first dose you may take 2 tablets within the first hour. Do not exceed 3 tablets in a 24-hour period. Children under 12 years: ask a doctor. **Questions or comments?** 1-800-395-0899 (Mon-Fri 8AM – 5PM EST)

Dist. by:
Bayer HealthCare LLC
Whippany, NJ 07981

86601648

DRUG FACTS INFO: (Modified Format—updated 5/18/18)		
CATEGORY	REQUIRED	ACTUAL IN FILE
Drug Facts Title	>7.0 pt.	-
Drug Facts Continued	7 pt.	-
Headings*	7 pt- bold italic *	5 pt.
Subheadings	≥6 pt.	5 pt.
Body Text	≥6 pt.	5 pt.
Bullets	5 pt.	9 pt.
Leading	≥0 pt.	5.65 pt.
Hairlines	0.5 pt. rule	-
Box and Barlines†	>0.5 pt. rule	-
Characters Per Inch:	No more than 38	> 38

* Headings are to be at least 2pts larger than Body Text.
 ** Drug Facts Title to be larger than all other type size used in DPP.
 † Box Barline may be omitted and color contrast used to highlight Drug Facts Info.

This is a representation of an electronic record that was signed electronically. Following this are manifestations of any and all electronic signatures for this electronic record.

/s/

NUSHIN F TODD
12/31/2020 10:37:00 AM