This label may not be the latest approved by FDA.
For current labeling information, please visit https://www.fda.gov/drugsatfda

100%



Actual Size 100%

FDA00357-5

ICL (updated 2/27/19)

CATEGORY	Bayer Standard	ACTUAL IN FILE
Drug Facts Title	>7 pt*	-
Drug Facts Continued	7 pt*	-
Headings	7 pt-bold italic*	4.5 pt
Subheadings	6 pt*	4.5 pt
Body Text	6 pt*	4.5 pt
Bullets	5 pt	5.5 pt
Hairlines	0.5 pt rule	-
Barlines‡	> 0.5 pt rule	-

Characters per inch rule is not applicable to ICL.
*ICL minimum font size is 4.5 pt.
< 0.5 pt leading may be used as long as ascenders and descenders do not touch.
† FDA recommends size 2.5 pt type

This label may not be the latest approved by FDA.
For current labeling information, please visit https://www.fda.gov/drugsatfda



Actual Size 100%

FDA00223-2

ICL (updated 2/27/19)

CATEGORY	Bayer Standard	ACTUAL IN FILE
Drug Facts Title	>7 pt*	-
Drug Facts Continued	7 pt*	-
Headings	7 pt-bold talic*	4.5 pt
Subheadings	6 pt*	4.5 pt
Body Text	6 pt*	4.5 pt
Bullets	5 pt	5.5 pt
Hairlines	0.5 pt rule	-
Bar ines†	> 0.5 pt rule	-

Characters per inch rule is not applicable to ICL.
*ICL minimum font size is 4.5 pt.
< 0.5 pt leading may be used as long as ascenders and descenders do not touch.
† FDA recommends size 2.5 pt type

Reference ID: 4772147

This label may not be the latest approved by FDA.
For current labeling information, please visit https://www.fda.gov/drugsatfda

Actual Size 100%

FDA00381-1

DRUG FACTS INFO (Updated 5/18/18)		
CATEGORY	REQUIRED	ACTUAL IN FILE
Drug Facts Title	>8 pt. * *	8.1 pt.
Drug Facts Continued	8 pt.	8 pt.
Headings*	8 pt.bold/ italic*	8 pt.
Subheadings	≥6 pt.	6 pt.
Body Text	≥6 pt.	6 pt.
Bullets	5 pt.	12 pt.
Leading	≥0.5 pt.	0 - 0.5 pt.
Hair lines	0.5 pt rule	0.5 pt.
Box and Borlines*	>2.5 pt rule	2.5 pt.
Characters Per Inch	No more than 30	<29

* Headings are to be at least 2pts larger than Body Text.
** Drug Facts Title to be larger than all other type size used in OPP 1 FDA Recommends 2.7pt rule

Reference ID: 4772147

This label may not be the latest approved by FDA.
For current labeling information, please visit https://www.fda.gov/drugsatfda

100%

Actual Size 100%

FDA00345-5

ICL (updated 2/27/19)

CATEGORY	Bayer Standard	ACTUAL IN FILE
Drug Facts Title	>7 pt*	-
Drug Facts Continued	7 pt*	-
Headings	7 pt-bold italic*	4.5 pt
Subheadings	6 pt*	4.5 pt
Body Text	6 pt*	4.5 pt
Bullets	5 pt	5.5 pt
Hairlines	0.5 pt rule	-
Barlines†	> 0.5 pt rule	-

Characters per inch rule is not applicable to ICL.
*ICL minimum font size is 4.5 pt.
< 0.5 pt leading may be used as long as ascenders and descenders do not touch.
† FDA recommends size 2.5 pt type

This label may not be the latest approved by FDA.
For current labeling information, please visit https://www.fda.gov/drugsatfda

100%

Actual Size 100%

FDA00344-3

ICL (updated 2/27/19)

CATEGORY	Bayer Standard	ACTUAL IN FILE
Drug Facts Title	>7 pt*	-
Drug Facts Continued	7 pt*	-
Headings	7 pt-bold italic*	4.5 pt
Subheadings	6 pt*	4.5 pt
Body Text	6 pt*	4.5 pt
Bullets	5 pt	5.5 pt
Hairlines	0.5 pt rule	-
Barlines‡	> 0.5 pt rule	-

Characters per inch rule is not applicable to ICL.
*ICL minimum font size is 4.5 pt.
< 0.5 pt leading may be used as long as ascenders and descenders do not touch.
† FDA recommends size 2.5 pt type

Reference ID: 4772147

This label may not be the latest approved by FDA.
For current labeling information, please visit https://www.fda.gov/drugsatfda

All Day Strong®

Drug Facts

Active ingredient (in each caplet) — **Purposes**

Naproxen sodium 220 mg (naproxen 200 mg) (NSAID)* ..... Pain reliever/fever reducer

*nonsteroidal anti-inflammatory drug

Uses

- temporarily relieves minor aches and pains due to:
  - minor pain of arthritis
  - muscular aches
  - backache
  - menstrual cramps
  - headache
  - toothache
  - the common cold
- temporarily reduces fever

Warnings

Allergy alert: Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Drug Facts (continued)

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- the stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic
- you have problems or serious side effects from taking pain relievers or fever reducers

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because naproxen may decrease this benefit of aspirin
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
  - feel faint
  - vomit blood
  - have bloody or black stools
  - have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
  - chest pain
  - trouble breathing
  - weakness in one part or side of body
  - slurred speech
  - leg swelling
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- you have difficulty swallowing
- it feels like the pill is stuck in your throat
- redness or swelling is present in the painful area
- any new symptoms appear

Drug Facts (continued)

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use naproxen sodium during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- **do not take more than directed**
- **the smallest effective dose should be used**
- drink a full glass of water with each dose

Adults and children 12 years and older	• take 1 caplet every 8 to 12 hours while symptoms last • for the first dose you may take 2 caplets within the first hour • do not exceed 2 caplets in any 8- to 12-hour period • do not exceed 3 caplets in a 24-hour period
Children under 12 years	• ask a doctor

Other information

- **each caplet contains:** sodium 20 mg
- store at 20-25°C (68-77°F). Avoid high humidity and excessive heat above 40°C (104°F).

Inactive ingredients

FD&C blue #2 lake, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, povidone, talc, titanium dioxide

Questions or comments?

1-800-395-0689 (Mon – Fri 9AM – 5PM EST)

DO NOT USE IF POUCH IS TORN OR OPEN.

Bayer, the Bayer Cross, Aleve and All Day Strong are registered trademarks of Bayer.

Dist. by: Bayer HealthCare LLC, Whippany, NJ 07981

©2020 Bayer

00000000

Bayer

Actual Size 100%

FDA00413-3

DRUG FACTS INFO (updated 5/18/18)

CATEGORY	REQUIRED	ACTUAL IN FILE
Drug Facts Title	>8 pt **	8.1 pt
Drug Facts Continued	8 pt	8 pt
Headings†	8 pt-bold talic†	8 pt
Subheadings	≥6 pt	6 pt
Body Text	≥6 pt	6 pt
Bullets	5 pt	12 pt
Leading	≥0.5 pt	0.5 - 0.7 pt
Hairlines	0.5 pt rule	0.5 pt
Box and Barlines†	>0.5 pt rule	2.5 pt
Characters Per Inch:	No more than 39	≤39

† Headings are to be at least 2pts larger than Body Text.
** Drug Facts Title to be larger than all other type size used in DFP
† FDA Recommends 2.5pt rule

Reference ID: 4772147

This label may not be the latest approved by FDA.
For current labeling information, please visit https://www.fda.gov/drugsatfda

100%

Actual Size 100%

FDA00343-5

ICL (updated 2/27/19)

CATEGORY	Bayer Standard	ACTUAL IN FILE
Drug Facts Title	>7 pt*	-
Drug Facts Continued	7 pt*	-
Headings	7 pt-bold italic*	4.5 pt
Subheadings	6 pt*	4.5 pt
Body Text	6 pt*	4.5 pt
Bullets	5 pt	5.5 pt
Hairlines	0.5 pt rule	-
Barlines‡	> 0.5 pt rule	-

Characters per inch rule is not applicable to ICL.
*ICL minimum font size is 4.5 pt.
< 0.5 pt leading may be used as long as ascenders and descenders do not touch.
† FDA recommends size 2.5 pt type

This label may not be the latest approved by FDA.
For current labeling information, please visit https://www.fda.gov/drugsatfda

100%

Actual Size 100%

FDA00344-3

ICL (updated 2/27/19)

CATEGORY	Bayer Standard	ACTUAL IN FILE
Drug Facts Title	>7 pt*	-
Drug Facts Continued	7 pt*	-
Headings	7 pt-bold italic*	4.5 pt
Subheadings	6 pt*	4.5 pt
Body Text	6 pt*	4.5 pt
Bullets	5 pt	5.5 pt
Hairlines	0.5 pt rule	-
Barlines‡	> 0.5 pt rule	-

Characters per inch rule is not applicable to ICL.
*ICL minimum font size is 4.5 pt.
< 0.5 pt leading may be used as long as ascenders and descenders do not touch.
† FDA recommends size 2.5 pt type

Reference ID: 4772147

This label may not be the latest approved by FDA.
For current labeling information, please visit https://www.fda.gov/drugsatfda

All Day Strong®

MANUFACTURER'S COUPON EXPIRES 00/00/0000

SAVE $2.00

ON ANY **ONE (1) ALEVE®**
20CT. or LARGER
(EXCLUDES ALEVE-D®)

CONSUMER: LIMIT ONE COUPON PER PURCHASE on specified product(s), size(s) and quantity stated. LIMIT OF 4 EXACT SAME COUPONS PER HOUSEHOLD PER DAY. You pay any sales tax. Void if copied, sold, transferred, altered, or auctioned.

RETAILER: You are authorized to act as our agent and redeem this coupon at face value on the specified product(s). We will reimburse you for the face value of this coupon plus 8¢ if submitted in compliance with the Bayer HealthCare LLC Coupon Redemption Policy, incorporated herein by reference. **USE NOT CONSISTENT WITH THESE TERMS MAY CONSTITUTE FRAUD AND MAY VOID ALL COUPONS SUBMITTED.** Send coupons to: Bayer HealthCare LLC 1355, P.O. Box 880001, El Paso, TX 88588-0001. Cash Value, 1/20th of 1¢. Good only in the U.S.A.

0000000-000000

Actual Size 100%

FDA00414-5

Drug Facts

Active ingredient (in each caplet) — **Purposes**
Naproxen sodium 220 mg (naproxen 200 mg) (NSAID)* ..... Pain reliever/fever reducer
*nonsteroidal anti-inflammatory drug

Uses
- temporarily relieves minor aches and pains due to:
  - minor pain of arthritis
  - muscular aches
  - backache
  - menstrual cramps
  - headache
  - toothache
  - the common cold
- temporarily reduces fever

Warnings
Allergy alert: Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:
- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:
- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Drug Facts (continued)

Do not use
- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if
- the stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic
- you have problems or serious side effects from taking pain relievers or fever reducers

Ask a doctor or pharmacist before use if you are
- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because naproxen may decrease this benefit of aspirin
- taking any other drug

When using this product
- take with food or milk if stomach upset occurs

Stop use and ask a doctor if
- you experience any of the following signs of stomach bleeding:
  - feel faint
  - vomit blood
  - have bloody or black stools
  - have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
  - chest pain
  - trouble breathing
  - weakness in one part or side of body
  - slurred speech
  - leg swelling
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- you have difficulty swallowing
- it feels like the pill is stuck in your throat
- redness or swelling is present in the painful area
- any new symptoms appear

Drug Facts (continued)

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use naproxen sodium during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions
- **do not take more than directed**
- **the smallest effective dose should be used**
- drink a full glass of water with each dose

Adults and children 12 years and older	• take 1 caplet every 8 to 12 hours while symptoms last • for the first dose you may take 2 caplets within the first hour • do not exceed 2 caplets in any 8- to 12-hour period • do not exceed 3 caplets in a 24-hour period
Children under 12 years	• ask a doctor

Other information
- **each caplet contains:** sodium 20 mg
- store at 20-25°C (68-77°F). Avoid high humidity and excessive heat above 40°C (104°F).

Inactive ingredients FD&C blue #2 lake, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, povidone, talc, titanium dioxide

Questions or comments? 1-800-395-0689
(Mon – Fri 9AM – 5PM EST)

DO NOT USE IF POUCH IS TORN OR OPEN.

Bayer, the Bayer Cross, Aleve and All Day Strong are registered trademarks of Bayer.
Dist. by: Bayer HealthCare LLC, Whippany, NJ 07981
©2020 Bayer 00000000 Bayer

DRUG FACTS INFO (updated 5/18/18)

CATEGORY	REQUIRED	ACTUAL IN FILE
Drug Facts Title	>8 pt **	8.1 pt
Drug Facts Continued	8 pt	8 pt
Headings†	8 pt-bold italic*	8 pt
Subheadings	≥6 pt	6 pt
Body Text	≥6 pt	6 pt
Bullets	5 pt	12 pt.
Leading	≥0.5 pt	0.5 - 0.7 pt
Hairlines	0.5 pt rule	0.5 pt
Box and Barlines†	>0.5 pt rule	2.5 pt
Characters Per Inch:	No more than 39	≤39

† Headings are to be at least 2pts larger than Body Text.
** Drug Facts Title to be larger than all other type size used in DFP
† FDA Recommends 2.5pt rule

Reference ID: 4772147

This label may not be the latest approved by FDA.
For current labeling information, please visit https://www.fda.gov/drugsatfda

100%

Actual Size 100%

FDA00377-3

ICL (updated 2/27/19)		
CATEGORY	**Bayer Standard**	**ACTUAL IN FILE**
Drug Facts Title	>7 pt*	-
Drug Facts Continued	7 pt*	-
Headings	7 pt-bold talic*	4.5 pt
Subheadings	6 pt*	4.5 pt
Body Text	6 pt*	4.5 pt
Bullets	5 pt	5.5 pt
Hairlines	0.5 pt rule	-
Bar ines†	> 0.5 pt rule	-

Characters per inch rule is not applicable to ICL.
*ICL minimum font size is 4.5 pt.
< 0.5 pt leading may be used as long as ascenders and descenders do not touch.
† FDA recommends size 2.5 pt type

Reference ID: 4772147

This label may not be the latest approved by FDA.
For current labeling information, please visit https://www.fda.gov/drugsatfda

100%

Reference ID: 4772147

Actual Size 100%

FDA00378-1

ICL (updated 2/27/19)		
CATEGORY	**Bayer Standard**	**ACTUAL IN FILE**
Drug Facts Ttle	>7 pt*	-
Drug Facts Continued	7 pt*	-
Headings	7 pt-bold italic*	4.5 pt
Subheadings	6 pt*	4.5 pt
Body Text	6 pt*	4.5 pt
Bullets	5 pt	5.5 pt
Hairlines	0.5 pt rule	-
Barlines†	> 0.5 pt rule	-

Characters per inch rule is not applicable to ICL.
*ICL minimum font size is 4.5 pt.
< 0.5 pt leading may be used as long as ascenders and descenders do not touch.
† FDA recommends size 2.5 pt type

This label may not be the latest approved by FDA.
For current labeling information, please visit https://www.fda.gov/drugsatfda

All Day Strong®

Actual Size 100%

FDA00412-4

Drug Facts

Active ingredient (in each tablet) — **Purposes**

Naproxen sodium 220 mg (naproxen 200 mg) (NSAID)* ..... Pain reliever/fever reducer

*nonsteroidal anti-inflammatory drug

Uses

- temporarily relieves minor aches and pains due to:
  - headache
  - minor pain of arthritis
  - backache
  - toothache
  - muscular aches
  - menstrual cramps
  - the common cold
- temporarily reduces fever

Warnings

Allergy alert: Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Drug Facts (continued)

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- the stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic
- you have problems or serious side effects from taking pain relievers or fever reducers

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because naproxen may decrease this benefit of aspirin
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
  - feel faint
  - vomit blood
  - have bloody or black stools
  - have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
  - chest pain
  - trouble breathing
  - weakness in one part or side of body
  - slurred speech
  - leg swelling
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- you have difficulty swallowing
- it feels like the pill is stuck in your throat
- redness or swelling is present in the painful area
- any new symptoms appear

Drug Facts (continued)

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use naproxen sodium during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- **do not take more than directed**
- **the smallest effective dose should be used**
- drink a full glass of water with each dose

Adults and children 12 years and older	• take 1 tablet every 8 to 12 hours while symptoms last • for the first dose you may take 2 tablets within the first hour • do not exceed 2 tablets in any 8- to 12-hour period • do not exceed 3 tablets in a 24-hour period
Children under 12 years	• ask a doctor

Other information

- **each tablet contains:** sodium 20 mg
- store at 20-25°C (68-77°F). Avoid high humidity and excessive heat above 40°C (104°F).

Inactive ingredients

FD&C blue #2 lake, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, povidone, talc, titanium dioxide

Questions or comments?

1-800-395-0689 (Mon – Fri 9AM – 5PM EST)

DO NOT USE IF POUCH IS TORN OR OPEN.

Bayer, the Bayer Cross, Aleve and All Day Strong are registered trademarks of Bayer.

Dist. by: Bayer HealthCare LLC, Whippany, NJ 07981

©2020 Bayer

00000000

Bayer

DRUG FACTS INFO (updated 5/18/18)

CATEGORY	REQUIRED	ACTUAL IN FILE
Drug Facts Title	>8 pt [+§]	8.1 pt
Drug Facts Continued	8 pt.	8 pt.
Headings[§]	8 pt-bold tallc[§]	8 pt.
Subheadings	≥6 pt.	6 pt.
Body Text	≥6 pt.	6 pt.
Bullets	5 pt.	12 pt.
Leading	≥0.5 pt.	0.5 - 0.7 pt.
Hairlines	0.5 pt rule	0.5 pt.
Box and Barlines[†]	>0.5 pt rule	2.5 pt.
Characters Per Inch:	No more than 39	≤39

§ Headings are to be at least 2pts larger than Body Text.
+§ Drug Facts Title to be larger than all other type size used in DFP
† FDA Recommends 2.5pt rule

Reference ID: 4772147

This label may not be the latest approved by FDA.
For current labeling information, please visit https://www.fda.gov/drugsatfda

---

This is a representation of an electronic record that was signed electronically. Following this are manifestations of any and all electronic signatures for this electronic record.

/s/

NUSHIN F TODD
04/01/2021 11:09:16 AM