

Tagamet[®]
 HB
 200

TAKE ANYTIME YOU NEED IT:
before meal • during meal • after meal

Tagamet[®]
 HB
 200
**Cimetidine Tablets 200 mg
 Acid Reducer**

JUST ONE TABLET
RELIEVES and PREVENTS
Heartburn and Acid Indigestion

Actual size



6
TABLETS
 (6 DOSES)

Tagamet[®]
 HB
 200

Tamper Evident: Each tablet
 individually sealed in a
 blister pack. Do not use if
 blister or foil is open or torn.

Dist. by Mediatech Products, Inc.
 Tarrytown, NY 10591
 A Presidio Consumer Healthcare company
 ©2018 Trade dress is owned by
 Mediatech Products, Inc. All rights reserved.
 Made in Spain. TALS004101



LOT:
 EXP:

- Tips for Managing Heartburn**
- Do not lie flat or bend over soon after eating
 - Do not eat late at night, or just before bedtime
 - Certain foods or drinks are more likely to cause heartburn, such as rich, spicy, fatty, and fried foods, chocolate, caffeine, alcohol, even some fruits and vegetables
 - Eat slowly and do not eat big meals
 - If you are overweight, lose weight
 - If you smoke, quit smoking
 - Raise the head of your bed
 - Wear loose fitting clothing around your stomach

10629-01-17

Drug Facts

Active ingredient (in each tablet)
 Cimetidine 200 mg.....**Purpose**
 Acid reducer

Uses ■ relieves heartburn associated with acid indigestion and sour stomach
 ■ prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain foods and beverages

Warnings

Allergy alert: Do not use if you are allergic to cimetidine or other acid reducers
Do not use ■ if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition.
 See your doctor. ■ with other acid reducers

Ask a doctor before use if you have

- frequent chest pain
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain
- had heartburn over 3 months. This may be a sign of a more serious condition.
- heartburn with lightheadedness, sweating or dizziness
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
- kidney disease
- liver disease

Ask a doctor or pharmacist before use if you are taking ■ a prescription drug.
 Acid reducers may interact with certain prescription drugs.

Stop use and ask a doctor if ■ you need to take this product for more than 14 days
 ■ stomach pain continues

■ your heartburn continues or worsens
If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions

- adults and children 12 years and over:
- to relieve symptoms, swallow 1 tablet with a glass of water
- to prevent symptoms, swallow 1 tablet with a glass of water **right before or any time up to 30 minutes before** eating food or drinking beverages that cause heartburn
- do not take more than 2 tablets in 24 hours
- children under 12 years: ask a doctor

Other information store at 15°-30°C (59°-86°F)

Inactive ingredients cellulose, corn starch, hypromellose, magnesium stearate, polyethylene glycol, polysorbate 80, povidone, sodium lauryl sulfate, sodium starch glycolate, titanium dioxide

Questions? Call toll-free 1-800-482-4394 weekdays tagamet.com

This label may not be the latest approved by FDA.
 For current labeling information, please visit <https://www.fda.gov/drugsatfda>

Font size: 9 pt
 Leading: 6 pt
 Horizontal scale: 95%
 Alignment: Left

Tagamet
 HB
 200

Dist. by: Medtech Products Inc., Tarrytown, NY 10591
 A Prestige Consumer Healthcare company
 ©2019 Trade dress is owned by Medtech Products Inc.
 All rights reserved. Made in Spain. TGIS002102
 For more information, visit tagamet.com
 Tamper Evident: Each tablet individually sealed in a blister pack. Do not use if blister or foil is open or torn.

002^{HB}
Tagamet

TAKE ANYTIME YOU NEED IT:
before meal • during meal • after meal

Tagamet
 HB
200
 Cimetidine Tablets 200 mg
 Acid Reducer

JUST ONE TABLET
RELIEVES and PREVENTS
Heartburn and Acid Indigestion

Actual size



30
TABLETS
 (30 DOSES)



0 42037 10363 7

LOT:
 EXP:

- Tips for Managing Heartburn**
- Do not lie flat or bend over soon after eating.
 - Do not eat late at night, or just before bedtime.
 - Certain foods or drinks are more likely to cause heartburn, such as rich, spicy, fatty, and fried foods; chocolate; caffeine; alcohol; even some fruits and vegetables.
 - Eat slowly and do not eat big meals.
 - If you are overweight, lose weight.
 - If you smoke, quit smoking.
 - Raise the head of your bed.
 - Wear loose fitting clothing around your stomach.

Drug Facts

Active ingredient (in each tablet) Cimetidine 200 mg
Purpose Acid reducer

Uses ■ relieves heartburn associated with acid indigestion and sour stomach
 ■ prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain foods and beverages

Warnings
Allergy alert: Do not use if you are allergic to cimetidine or other acid reducers
Do not use ■ if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition.
 See your doctor. ■ with other acid reducers

Ask a doctor before use if you have
 ■ frequent chest pain ■ frequent wheezing, particularly with heartburn
 ■ unexplained weight loss ■ nausea or vomiting ■ stomach pain
 ■ had heartburn over 3 months. This may be a sign of a more serious condition.
 ■ heartburn with lightheadedness, sweating or dizziness
 ■ chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
 ■ kidney disease ■ liver disease

Ask a doctor or pharmacist before use if you are taking ■ a prescription drug.
 Acid reducers may interact with certain prescription drugs.

Stop use and ask a doctor if ■ you need to take this product for more than 14 days
 ■ stomach pain continues ■ your heartburn continues or worsens

If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions
 ■ adults and children 12 years and over:
 ■ to relieve symptoms, swallow 1 tablet with a glass of water
 ■ to prevent symptoms, swallow 1 tablet with a glass of water **right before or any time up to 30 minutes before** eating food or drinking beverages that cause heartburn
 ■ do not take more than 2 tablets in 24 hours
 ■ children under 12 years: ask a doctor

Other information store at 15°-30°C (59°-86°F)

Inactive ingredients cellulose, corn starch, hypromellose, magnesium stearate, polyethylene glycol, polysorbate 80, povidone, sodium lauryl sulfate, sodium starch glycolate, titanium dioxide

Questions? Call toll-free 1-800-482-4394 weekdays tagamet.com

10628-01-17

Quantity
 Font family: Myriad Pro
 Font size: 10.51517 pt

JOB #: 1G05001102
 TECH DRAWING #: J80940A / DL986
 UPC#: 0-42037-00077-6

This label may not be the latest approved by FDA.
 For current labeling information, please visit <https://www.fda.gov/drugsatfda>

002^{HB}
Tagamet

+20% MORE FREE
36 TABLETS FOR THE PRICE OF 30

TAKE ANYTIME YOU NEED IT:
 before meal • during meal • after meal

Tagamet
^{HB} **200**
 Cimetidine Tablets 200 mg
 Acid Reducer

JUST ONE TABLET
 RELIEVES and PREVENTS
 Heartburn and Acid Indigestion

Actual Size



30 36
TABLETS
 (30 36 DOSES)

Tagamet
^{HB} **200**

**Just ONE TABLET RELIEVES
 and PREVENTS Heartburn
 and Acid Indigestion**

Tamper-Evident: Each tablet individually sealed in a blister pack. Do not use if blister or foil is open or torn.

Dist. by Mediatech Products, Inc., Terroton, NY 10591
 A Prescribing Consumer Healthcare company
 ©2019. Trade dress is owned by Mediatech Products, Inc.
 All rights reserved. Made in Spain. 1G05001102
 For more information, visit tagamet.com

- Tips for Managing Heartburn**
- Do not lie flat or bend over soon after eating
 - Do not eat late at night, or just before bedtime
 - Certain foods or drinks are more likely to cause heartburn, such as rich, spicy, fatty, and fried foods, chocolate, caffeine, alcohol, even some fruits and vegetables
 - Eat slowly and do not eat big meals
 - If you are overweight, lose weight
 - If you smoke, quit smoking
 - Raise the head of your bed
 - Wear loose-fitting clothing around your stomach

Drug Facts

Active ingredient (in each tablet)	Purpose
Cimetidine 200 mg	Acid reducer

Uses ■ relieves heartburn associated with acid indigestion and sour stomach
 ■ prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain foods and beverages

Warnings
Allergy alert: Do not use if you are allergic to cimetidine or other acid reducers
Do not use ■ if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor. ■ with other acid reducers

Ask a doctor before use if you have

- frequent chest pain
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain
- had heartburn over 3 months. This may be a sign of a more serious condition.
- heartburn with lightheadedness, sweating or dizziness
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
- kidney disease
- liver disease

Ask a doctor or pharmacist before use if you are taking ■ a prescription drug. Acid reducers may interact with certain prescription drugs.

Stop use and ask a doctor if ■ you need to take this product for more than 14 days
 ■ stomach pain continues ■ your heartburn continues or worsens

If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions

- adults and children 12 years and over:
 - to relieve symptoms, swallow 1 tablet with a glass of water
 - to prevent symptoms, swallow 1 tablet with a glass of water right before or any time up to 30 minutes before eating food or drinking beverages that cause heartburn
 - do not take more than 2 tablets in 24 hours
 - children under 12 years: ask a doctor

Other information store at 15°-30°C (59°-86°F)

Inactive ingredients cellulose, corn starch, hypromellose, magnesium stearate, polyethylene glycol, polysorbate 80, povidone, sodium lauryl sulfate, sodium starch glycolate, titanium dioxide

Questions? Call toll-free 1-800-482-4394 weekdays tagamet.com



LOT:
 EXP:

Reference ID: 4590825
 Tablets (DOSES)
 Font family: Myriad Pro

Font family: Myriad Pro

TIPS BODY
 Font family: Helvetica
 Font style: Bold Condensed

TIPS HEADER
 Font family: Helvetica
 Font style: Bold Condensed

Tagamet[®]
HB
200

BETTER VALUE
40 PILLS for the PRICE OF 30

TAKE ANYTIME YOU NEED IT:
before meal • during meal • after meal

Tagamet[®]
HB
200
Cimetidine Tablets 200 mg
Acid Reducer

JUST ONE TABLET
RELIEVES and PREVENTS
Heartburn and Acid Indigestion

Actual size



40
TABLETS
(40 DOSES)

Temper-Evident: Each tablet, individually sealed, in a blister pack. Do not use if blister or foil is open or torn.

Tagamet[®]
HB
200

Just ONE TABLET RELIEVES
and PREVENTS Heartburn
and Acid Indigestion

Dist. by Mediatech Products Inc., Tarrytown, NY 10591
A Prestige Consumer Healthcare company
©2019 Trade dress is owned by Mediatech Products Inc.
All rights reserved. Made in Spain. TGIS007103
For more information, visit tagamet.com



0 42037 00078 3

LOT:
EXP:

- Tips for Managing Heartburn**
- Do not lie flat or bend over soon after eating
 - Do not eat late at night, or just before bedtime
 - Certain foods or drinks are more likely to cause heartburn, such as rich, spicy, fatty, and fried foods, chocolate, caffeine, alcohol, even some fruits and vegetables
 - Eat slowly and do not eat big meals
 - If you are overweight, lose weight
 - If you smoke, quit smoking
 - Raise the head of your bed
 - Wear loose fitting clothing around your stomach

Drug Facts

Active ingredient (in each tablet)

Cimetidine 200 mg

Purpose

Acid reducer

Uses

- relieves heartburn associated with acid indigestion and sour stomach
- prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain foods and beverages

Warnings

Allergy alert: Do not use if you are allergic to cimetidine or other acid reducers

Do not use ■ if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition.

See your doctor. ■ with other acid reducers

Ask a doctor before use if you have

- frequent chest pain
- unexplained weight loss
- had heartburn over 3 months. This may be a sign of a more serious condition.
- heartburn with lightheadedness, sweating or dizziness
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
- kidney disease
- frequent wheezing, particularly with heartburn
- nausea or vomiting
- stomach pain
- liver disease

Ask a doctor or pharmacist before use if you are taking ■ a prescription drug. Acid reducers may interact with certain prescription drugs.

Stop use and ask a doctor if ■ you need to take this product for more than 14 days ■ stomach pain continues ■ your heartburn continues or worsens

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions

- adults and children 12 years and over:
 - to relieve symptoms, swallow 1 tablet with a glass of water
 - to prevent symptoms, swallow 1 tablet with a glass of water **right before or any time up to 30 minutes before** eating food or drinking beverages that cause heartburn
- do not take more than 2 tablets in 24 hours
- children under 12 years: ask a doctor

Other information store at 15°-30°C (59°-86°F)

Inactive ingredients cellulose, corn starch, hypromellose, magnesium stearate, polyethylene glycol, polysorbate 80, povidone, sodium lauryl sulfate, sodium starch glycolate, titanium dioxide

Questions? Call toll-free 1-800-482-4394 weekdays

tagamet.com

13270-MM-YY

50 Tablets
 USA
 TGUS009102
 J61602A
 0-42037-10745-1
 PRINTS 6/0

Regulatory



Drug facts
 Font family: Prestige DF
 Font size: 9 pt
 Leading: 6 pt
 Horizontal scale: 95%
 Alignment: Left

evident
 Neue
 6 pt
 7 pt
 100%
 Left

200
 9 pt
 10%
 Left

Name
 Pro
 2 pt
 0%
 ght

bution
 T Std
 6 pt
 5 pt
 95%
 Right

Tamper Evident: Each tablet individually sealed in a blister pack. Do not use if blister or foil is open or torn.

Dist. by Medtech Products Inc., Tarrytown, NY 10591
 A Prestige Consumer Healthcare company
 ©2019 Trade dress is owned by Medtech Products Inc.
 All rights reserved. Made in Spain. TGUS009102

Just ONE TABLET RELIEVES and PREVENTS Heartburn and Acid Indigestion

- Tips for Managing Heartburn**
- Do not lie flat or bend over soon after eating
 - Do not eat late at night, or just before bedtime
 - Certain foods or drinks are more likely to cause heartburn, such as rich, spicy, fatty, and fried foods, chocolate, caffeine, alcohol, even some fruits and vegetables
 - Eat slowly and do not eat big meals
 - If you are overweight, lose weight
 - If you smoke, quit smoking
 - Raise the head of your bed
 - Wear loose fitting clothing around your stomach

Drug Facts

Active ingredient (in each tablet)	Purpose
Cimetidine 200 mg	Acid reducer

Uses ■ relieves heartburn associated with acid indigestion and sour stomach
 ■ prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain foods and beverages

Warnings
Allergy alert: Do not use if you are allergic to cimetidine or other acid reducers
Do not use ■ if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor. ■ with other acid reducers

Ask a doctor before use if you have

- frequent chest pain
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain
- had heartburn over 3 months. This may be a sign of a more serious condition.
- heartburn with lightheadedness, sweating or dizziness
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
- kidney disease
- liver disease

Ask a doctor or pharmacist before use if you are taking ■ a prescription drug. Acid reducers may interact with certain prescription drugs.

Stop use and ask a doctor if ■ you need to take this product for more than 14 days
 ■ stomach pain continues ■ your heartburn continues or worsens

If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions

- adults and children 12 years and over:
 - to relieve symptoms, swallow 1 tablet with a glass of water
 - to prevent symptoms, swallow 1 tablet with a glass of water right before or any time up to 30 minutes before eating food or drinking beverages that cause heartburn
 - do not take more than 2 tablets in 24 hours
 - children under 12 years: ask a doctor

Other information store at 15°-30°C (59°-86°F)

Inactive ingredients cellulose, corn starch, hypromellose, magnesium stearate, polyethylene glycol, polysorbate 80, povidone, sodium lauryl sulfate, sodium starch glycolate, titanium dioxide

Questions? Call toll-free 1-800-482-4394 weekdays tagamet.com



quantity
 Font family: Myriad Pro
 Font size: 12.46756 pt
 Leading: 11 pt
 Horizontal scale: 100%

TIPS BODY
 Font family: Helvetica
 Font style: Bold Condensed
 Font size: 7 pt
 Leading: 8 pt
 Vertical scale: 100%

TIPS HEADER
 Font family: Helvetica
 Font style: Bold Condensed
 Font size: 9 pt
 Leading: (10.8 pt)
 Vertical scale: 100%