

MM/YY  
 XXXXXX  
 FORMAT BELOW:  
 PRINTED OR DEBOSSED ONLINE  
 LOT/EXP

NO COATING

12789-MM-YY

Dist. by Medtech Products Inc.  
 Tarrytown, NY 10591  
 A Prestige Consumer Healthcare company  
 ©2020 Trade dress is owned by  
 Medtech Products Inc. All rights reserved.  
 Made in Spain. TGUS013101

GLUE - NO COATING

**Drug Facts**

Active ingredient	Purpose
Cimetidine 200 mg	Acid reducer

**Uses** ■ relieves heartburn associated with acid indigestion and sour stomach  
 ■ prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain foods and beverages

**Warnings**  
**Allergy alert:** Do not use if you are allergic to cimetidine or other acid reducers.

**Do not use** ■ if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor. ■ with other acid reducers

**Ask a doctor before use if you have** ■ frequent chest pain ■ frequent wheezing, particularly with heartburn ■ unexplained weight loss ■ nausea or vomiting ■ stomach pain ■ had heartburn over 3 months. This may be a sign of a more serious condition. ■ heartburn with lightheadedness, sweating or dizziness ■ chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness ■ kidney disease ■ liver disease

**Ask a doctor or pharmacist before use if you are taking** ■ a prescription drug.

Acid reducers may interact with certain prescription drugs.

**Stop use and ask a doctor if** ■ you need to take this product for more than 14 days ■ stomach pain continues ■ your heartburn continues or worsens

**If pregnant or breast-feeding**, ask a health professional before use.  
**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

**Directions** Adults and children 12 years and over: ■ to relieve symptoms, swallow 1 tablet with a glass of water ■ to prevent symptoms, swallow 1 tablet with a glass of water right before or any time up to 30 minutes before eating food or drinking beverages that cause heartburn ■ do not take more than 2 tablets in 24 hours. Children under 12 years: ask a doctor

**Other information**  
 store at 15°-30°C (59°-86°F)

**Inactive ingredients** cellulose, corn starch, hypromellose, magnesium stearate, polyethylene glycol, polysorbate 80, povidone, sodium lauryl sulfate, sodium starch glycolate, titanium dioxide

**Questions?**  
 Call toll-free 1-800-482-4394 weekdays.

Keep outer carton for full information.

**NEW VALUE SIZE!**

**TAKE ANYTIME YOU NEED IT:**  
 before meal · during meal · after meal

**Tagamet<sup>®</sup> HB 200**  
 Cimetidine Tablets 200 mg  
 Acid Reducer

**JUST ONE TABLET**  
 RELIEVES and PREVENTS  
 Heartburn and Acid Indigestion



**70 TABLETS**  
 (70 DOSES)

**TIPS FOR MANAGING HEARTBURN:**

- Do not lie flat or bend over soon after eating
- Do not eat late at night, or just before bedtime
- Certain foods or drinks are more likely to cause heartburn, such as rich, spicy, fatty, and fried foods, chocolate, caffeine, alcohol, even some fruits and vegetables
- Eat slowly and do not eat big meals
- If you are overweight, lose weight
- If you smoke, quit smoking
- Raise the head of your bed
- Wear loose fitting clothing around your stomach

Tamper Evident: Do Not Use if seal under cap is broken or missing.



NO COATING

FRONT

0.407" X 1.3"  
VARIABLE DATA AREA  
NO INK/NO COAT

<p>TAKE ANYTIME YOU NEED IT: before meal • during meal • after meal</p> <p><b>Tagamet</b> HD 200</p> <p>Cimetidine Tablets 200 mg Acid Reducer</p> <p><b>JUST ONE TABLET</b></p> <p>RELIEVES and PREVENTS Heartburn and Acid Indigestion</p> <p>70 TABLETS (70 DOSES)</p> <p>© 2020 Dist. by Medtech Products Inc., Tarrytown, NY 10591. A Prestige Consumer Healthcare company. Made in Spain. TBUS013201</p>	<p><b>Drug Facts</b></p> <p><b>Active ingredient Purpose</b> (in each tablet) Cimetidine 200 mg ..... Acid reducer</p> <p><b>Uses</b></p> <ul style="list-style-type: none"> <li>relieves heartburn associated with acid indigestion and sour stomach</li> <li>prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain foods and beverages</li> </ul>	<p><b>Warnings</b></p> <p><b>Allergy alert:</b> Do not use if you are allergic to cimetidine or other acid reducers.</p> <p><b>Do not use</b></p> <ul style="list-style-type: none"> <li>if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.</li> <li>with other acid reducers</li> </ul> <p><b>Tamper Evident: Do Not Use if seal under cap is broken or missing.</b></p>
	<p>© 2020 Dist. by Medtech Products Inc., Tarrytown, NY 10591. A Prestige Consumer Healthcare company. Made in Spain. TBUS013201</p>	

BACK

<p><b>Drug Facts (continued)</b></p> <p>Ask a doctor before use if you have</p> <ul style="list-style-type: none"> <li>frequent chest pain</li> <li>frequent wheezing, particularly with heartburn</li> <li>unexplained weight loss</li> <li>nausea or vomiting</li> <li>stomach pain</li> <li>had heartburn over 3 months. This may be a sign of a more serious condition.</li> <li>heartburn with lightheadedness, sweating or dizziness</li> <li>chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness</li> </ul>	<ul style="list-style-type: none"> <li>kidney disease</li> <li>liver disease</li> </ul> <p>Ask a doctor or pharmacist before use if you are taking a prescription drug. Acid reducers may interact with certain prescription drugs.</p> <p>Stop use and ask a doctor if you need to take this product for more than 14 days stomach pain continues your heartburn continues or worsens</p> <p>If pregnant or breast-feeding, ask a health professional before use.</p> <p>Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.</p>	<p>0.5" hinge non printing area</p>
--	--	---

BASE

<p>0.5" hinge non printing area</p>	<p><b>Drug Facts (continued)</b></p> <p><b>Directions</b></p> <p>Adults and children 12 years and over:</p> <ul style="list-style-type: none"> <li>to relieve symptoms, swallow 1 tablet with a glass of water</li> <li>to prevent symptoms, swallow 1 tablet with a glass of water right before or any time up to 30 minutes before eating food or drinking beverages that cause heartburn</li> <li>do not take more than 2 tablets in 24 hours.</li> </ul>	<p>Children under 12 years: ask a doctor</p> <p><b>Other information</b> store at 15°-30°C (59°-86°F)</p> <p><b>Inactive ingredients</b> cellulose, corn starch, hypromellose, magnesium stearate, polyethylene glycol, polysorbate 80, povidone, sodium lauryl sulfate, sodium starch glycolate, titanium dioxide</p> <p><b>Questions?</b> 1-800-482-4394</p>
---	--	--

-----  
**This is a representation of an electronic record that was signed electronically. Following this are manifestations of any and all electronic signatures for this electronic record.**  
-----

/s/  
-----

NUSHIN F TODD  
03/15/2021 01:37:30 PM