



Tamper Evident: Each tablet individually sealed in a blister pack. Do not use if blister or foil is open or torn.

Tagamet
HB
200

NEW!

TAKE ANYTIME YOU NEED IT:
before meal • during meal • after meal

Tagamet
HB
200
Cimetidine Tablets 200 mg
Acid Reducer

JUST ONE TABLET
RELIEVES and PREVENTS
Heartburn and Acid Indigestion

ICY COOL
MINT
FLAVOR

Actual size

40
TABLETS
(40 DOSES)

Dist. by Medtech Products Inc., Tarrytown, NY 10591
 A Prestige Consumer Healthcare company
 ©2022 Trade dress is owned by Medtech Products Inc.
 All rights reserved. Made in Spain. TGUS00X101

Tips for Managing Heartburn

- Do not lie flat or bend over soon after eating
- Do not eat late at night, or just before bedtime
- Certain foods or drinks are more likely to cause heartburn, such as rich, spicy, fatty, and fried foods, chocolate, caffeine, alcohol, even some fruits and vegetables
- Eat slowly and do not eat big meals
- If you are overweight, lose weight
- If you smoke, quit smoking
- Raise the head of your bed
- Wear loose fitting clothing around your stomach

Drug Facts

Active ingredient (in each tablet)	Purpose
Cimetidine 200 mg	Acid reducer

Uses ■ relieves heartburn associated with acid indigestion and sour stomach
 ■ prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain foods and beverages

Warnings
Allergy alert: Do not use if you are allergic to cimetidine or other acid reducers.
Do not use ■ if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor. ■ with other acid reducers

Ask a doctor before use if you have ■ frequent chest pain ■ frequent wheezing, particularly with heartburn ■ unexplained weight loss ■ nausea or vomiting
 ■ stomach pain ■ had heartburn over 3 months. This may be a sign of a more serious condition. ■ heartburn with lightheadedness, sweating or dizziness
 ■ chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness ■ kidney disease ■ liver disease

Ask a doctor or pharmacist before use if you are taking ■ a prescription drug. Acid reducers may interact with certain prescription drugs.

Stop use and ask a doctor if ■ you need to take this product for more than 14 days ■ stomach pain continues ■ your heartburn continues or worsens

If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions
Adults and children 12 years and over:
 ■ to relieve symptoms, swallow 1 tablet with a glass of water
 ■ to prevent symptoms, swallow 1 tablet with a glass of water right before or any time up to 30 minutes before eating food or drinking beverages that cause heartburn ■ do not take more than 2 tablets in 24 hours ■ do not chew tablet.
Children under 12 years: ask a doctor



Other information ■ store at 15°-30°C (59°-86°F)
 ■ Keep outer carton for full information

Inactive ingredients cellulose, corn starch, FD&C Blue #1 Lake, flavor, hypromellose, magnesium stearate, polyethylene glycol, polysorbate 80, povidone, sodium lauryl sulfate, sodium starch glycolate, titanium dioxide

Questions? Call toll-free 1-800-482-4394 weekdays.

■ Keep outer carton for full information

VARIABLE DATA
 NO INK / NO COAT
 1.5" X 0.50"



MM/YY
 XXXXXX

LOT/EXP
 PRINTED OR DEBOSSED ONLINE
 FORMAT BELOW:

GLUE - NO COATING

NO COATING

12789-MM-YY

Dist. by Medtech Products Inc.
 Tarrytown, NY 10591
 A Prestige Consumer Healthcare company
 ©2021 Trade dress is owned by
 Medtech Products Inc. All rights reserved.
 Made in Spain. TGUS0XX101

Drug Facts

Active ingredient (in each tablet) **Purpose**
 Cimetidine 200 mg Acid reducer

Uses ■ relieves heartburn associated with acid indigestion and sour stomach
 ■ prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain foods and beverages

Warnings
Allergy alert: Do not use if you are allergic to cimetidine or other acid reducers.
Do not use ■ if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor. ■ with other acid reducers
Ask a doctor before use if you have ■ frequent chest pain ■ frequent wheezing, particularly with heartburn ■ unexplained weight loss ■ nausea or vomiting ■ stomach pain ■ had heartburn over 3 months. This may be a sign of a more serious condition. ■ heartburn with lightheadedness, sweating or dizziness ■ chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness ■ kidney disease ■ liver disease
Ask a doctor or pharmacist before use if you are taking ■ a prescription drug.

Drug Facts (continued)

Acid reducers may interact with certain prescription drugs.
Stop use and ask a doctor if ■ you need to take this product for more than 14 days ■ stomach pain continues ■ your heartburn continues or worsens
If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions Adults and children 12 years and over: ■ to relieve symptoms, swallow 1 tablet with a glass of water ■ to prevent symptoms, swallow 1 tablet with a glass of water **right before or any time up to 30 minutes before** eating food or drinking beverages that cause heartburn ■ do not take more than 2 tablets in 24 hours. ■ do not chew tablet. Children under 12 years: ask a doctor

Other information
 store at 15° - 30°C (59° - 86°F)

Inactive ingredients cellulose, corn starch, FD&C Blue #1 Lake, flavor, hypromellose, magnesium stearate, polyethylene glycol, polysorbate 80, povidone, sodium lauryl sulfate, sodium starch glycolate, titanium dioxide

Questions?
 Call toll-free 1-800-482-4394 weekdays.

NEW! VALUE SIZE!

TAKE ANYTIME YOU NEED IT:
 before meal · during meal · after meal

Tagamet[®] HB 200
 Cimetidine Tablets 200 mg Acid Reducer

JUST ONE TABLET

RELIEVES and PREVENTS
 Heartburn and Acid Indigestion

Actual size

ICY COOL MINT FLAVOR

70 TABLETS (70 DOSES)

TIPS FOR MANAGING HEARTBURN:

- Do not lie flat or bend over soon after eating
- Do not eat late at night, or just before bedtime
- Certain foods or drinks are more likely to cause heartburn, such as rich, spicy, fatty, and fried foods, chocolate, caffeine, alcohol, even some fruits and vegetables
- Eat slowly and do not eat big meals
- If you are overweight, lose weight
- If you smoke, quit smoking
- Raise the head of your bed
- Wear loose fitting clothing around your stomach

Tamper Evident: Do Not Use If seal under cap is broken or missing.

NO COATING

FPO
 0 42037 00128 5

FRONT

0.40" X 1.5"
VARIABLE DATA AREA
NO INK/NO COAT

NEW! TAKE ANYTIME YOU NEED IT:
before meals, during meals, after meals

Tagamet[®] HB 200
Cimetidine Tablets 200 mg
Acid Reducer

JUST ONE TABLET
RELIEVES and PREVENTS
Heartburn and Acid Indigestion

ICY COOL MINT FLAVOR

70 TABLETS
(70 DOSES)

Keep outer carton for full information. 14822-244-17

Drug Facts	Drug Facts (continued)
<p>Active ingredient Purpose (in each tablet) Cimetidine 200 mg Acid reducer</p> <p>Uses</p> <ul style="list-style-type: none"> relieves heartburn associated with acid indigestion and sour stomach prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain foods and beverages 	<p>Warnings Allergy alert: Do not use if you are allergic to cimetidine or other acid reducers.</p> <p>Do not use</p> <ul style="list-style-type: none"> if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor. with other acid reducers

©2021 Dist. by Medtch Products Inc., Terrytown, NY 10591, Made in Spain. TGU8013201

Tamper Evident: Do Not Use if seal under cap is broken or missing.

WEB: www.tagamet.com

BACK

<p>Drug Facts (continued)</p> <p>Ask a doctor before use if you have</p> <ul style="list-style-type: none"> frequent chest pain frequent wheezing, particularly with heartburn unexplained weight loss nausea or vomiting stomach pain had heartburn over 3 months. This may be a sign of a more serious condition. heartburn with lightheadedness, sweating or dizziness chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness 	<p>Drug Facts (continued)</p> <ul style="list-style-type: none"> kidney disease liver disease <p>Ask a doctor or pharmacist before use if you are taking a prescription drug. Acid reducers may interact with certain prescription drugs.</p> <p>Stop use and ask a doctor if you need to take this product for more than 14 days • stomach pain continues • your heartburn continues or worsens</p> <p>If pregnant or breast-feeding, ask a health professional before use.</p> <p>Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.</p>	<p>0.5" hinge non printing area</p>
---	---	---

BASE

<p>0.5" hinge non printing area</p>	<p>Drug Facts (continued)</p> <p>Directions</p> <p>Adults and children 12 years and over:</p> <ul style="list-style-type: none"> to relieve symptoms, swallow 1 tablet with a glass of water to prevent symptoms, swallow 1 tablet with a glass of water right before or any time up to 30 minutes before eating food or drinking beverages that cause heartburn • do not take more than 2 tablets in 24 hours • do not chew tablet. <p>Children under 12 years: ask a doctor</p>	<p>Drug Facts (continued)</p> <p>Other information store at 15° – 30°C (59° – 86°F)</p> <p>Inactive ingredients cellulose, corn starch, FD&C Blue #1 Lake, flavor, hypromellose, magnesium stearate, polyethylene glycol, polysorbate 80, povidone, sodium lauryl sulfate, sodium starch glycolate, titanium dioxide</p> <p>Questions? Call toll-free 1-800-482-4394 weekdays.</p>
---	---	--

This is a representation of an electronic record that was signed electronically. Following this are manifestations of any and all electronic signatures for this electronic record.

/s/

NUSHIN F TODD
05/24/2022 08:04:34 PM