







(Do not use if the seal under the cap is missing. Do not use if the cap is missing. Do not use if the seal is broken or missing.)

**Acid Reducer**

**Famotidine 10 mg** Acid Reducer

**Uses** • relieves heartburn associated with acid indigestion and sour stomach  
• prevents heartburn associated with acid indigestion and sour stomach brought on by you eating or drinking certain food and beverages

**Warnings** Allergy Alert: Do not use if you are allergic to famotidine or other acid reducers. Do not use if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. There may be signs of serious condition. See your

**ORIGINAL STRENGTH**  
NDC 10037-472-25

**Pepcid<sup>®</sup> AC**

Famotidine Tablets 10 mg Acid Reducer

**Just One Tablet!**

Prelvents & Relieves Heartburn Due to Acid Indigestion

90 Tablets

doctor • with other acid reducers. Ask a doctor before use if you have • had heartburn over 3 months. This may be a sign of a more serious condition.  
• heart burn with difficulty swallowing, or difficulty • chest pain or shoulder pain with shortness of breath, sweating, pain spreading to arms, neck or shoulders or lightheadedness • frequent chest pain • frequent vomiting, particularly with heartburn • unexplained weight loss • nausea or vomit • stomach pain. Stop use and ask a doctor if your heartburn continues or worsens. • you need to take this product for more than 14 days. If pregnant or breast-feeding, ask a health professional before use. If you get pregnant while taking this product, get medical help or contact a Poison Control Center right away. (1-800-232-1232)

**Directions** • adults and children 12 years and over • to relieve symptoms, swallow 1 tablet with a glass of water. Do not chew. • to prevent symptoms, swallow 1 tablet with a glass of water at any time from 15 to 30 minutes before eating food or drinking beverages that cause heartburn. • do not use more than 2 tablets in 24 hours • children under 12 years • ask a doctor

Famotidine Active ingredient in the Republic of Ireland. Processed in Canada. Distributed by JANSSEN JANSSEN (CONSUMER), INC. McKeesport, PA 15136, USA. ©McNeil-PPC, Inc. PEPCID and PEPCID AC are registered trademarks of McNeil-PPC.

Do not use if foil seal under bottle cap printed "SAFETY SEAL" is broken or missing.

**Active ingredient (in each tablet)**  
Famotidine 10 mg

**Purpose**  
Acid reducer

**Uses** • relieves heartburn associated with acid indigestion and sour stomach • prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain food and beverages

**Warnings** Allergy alert: Do not use if you are allergic to famotidine or other acid reducers. Do not use • if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor. • with other acid reducers. Ask a doctor before use if you have • had heartburn over 3 months. This may be a sign of a more serious condition. • heartburn with lightheadedness, sweating, or dizziness • chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness • frequent chest pain • frequent wheezing, particularly with heartburn

**ORIGINAL STRENGTH**  
NDC 140-07-012-02

**Pepcid<sup>®</sup> AC**  
Famotidine Tablets 10 mg Acid Reducer  
**Just One Tablet!**  
Prevents & Relieves Heartburn  
Due to Acid Indigestion  
**90 Tablets**

• unexplained weight loss • nausea or vomiting • stomach pain. Stop use and ask a doctor if • your heartburn continues or worsens • you need to take this product for more than 14 days. If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222) **Directions** • adults and children 12 years and over • to relieve symptoms, swallow 1 tablet with a glass of water. Do not chew. • to prevent symptoms, swallow 1 tablet with a glass of water at any time from 15 to 60 minutes before eating food or drinking beverages that cause heartburn • do not use more than 2 tablets in 24 hours • children under 12 years: ask a doctor Famotidine (active ingredient) made in Republic of Ireland, Processed in Canada 000000000000

E.P.  
KAMACHO/STY  
10/17/12  
LOT: 101

140-07-012-02  
FAMOTIDINE TABLETS 10 MG ACID REDUCER  
90 TABLETS  
NDC 140-07-012-02  
LOT: 101

**Famotidine Tablets 20 mg**  
**Famotidine Tablets 20 mg**  
**Famotidine Tablets 20 mg**  
**Famotidine Tablets 20 mg**

**SOI font: 22 pt Helvetica Bold Condensed Oblique**







**Famotidine Tablets 20 mg**  
**Famotidine Tablets 20 mg**  
**Famotidine Tablets 20 mg**  
**Famotidine Tablets 20 mg**

**SOI font: 19.35 pt Helvetica Bold Condensed Oblique**



<p>Do not use if the seal is under both caps printed SAFETYSEAL is broken or missing. <b>Active ingredient (in each tablet)</b> Famotidine 20 mg</p> <p><b>Purpose</b> Acid reducer</p> <p><b>Uses</b> • relieves heartburn associated with acid indigestion and sour stomach • prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain food and beverages</p> <p><b>Warnings</b> Allergy alert: Do not use if you are allergic to famotidine or other acid reducers. Do not use if you have trouble or pain swallowing food, vomit, or any red blood or bloody or black stools. There may be signs of a serious condition. See your doctor. *If you have kidney disease, consult</p>	<p><b>MAXIMUM STRENGTH</b> NDC 1987-415-16</p> <p><b>Pepcid AC</b> Famotidine Tablets 20 mg Acid Reducer <b>Just One Tablet!</b> Prevents &amp; Relieves Heartburn Due to Acid Indigestion <b>50 Tablets</b></p>	<p>Under the advice and supervision of a doctor • with other acid reducers. Ask a doctor before use if you have had heartburn over 1 month. This may be a sign of a more serious condition. • heartburn with difficulty swallowing, vomiting, and indigestion • chest pain or shoulder pain with shortness of breath, nausea, or pain spreading to arms, neck or shoulders, or light-headedness • frequent chest pain • frequent wheezing, particularly with heartburn • a cough and weight loss • nausea or vomiting • stomach pain. Stop use and ask a doctor if • your heartburn continues or worsens • you need to take this product for more than 14 days. If you experience dizziness, ask a health professional before use. Stop use if much of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-332-2222)</p> <p><b>Directions</b> • Adults and children 12 years and over • To relieve symptoms, swallow 1 tablet with a glass of water. Do not chew. • To prevent symptoms, swallow 1 tablet with a glass of water at any time from 10 to 60 minutes before eating food or drinking beverages that cause heartburn • do not use more than 2 tablets in 24 hours • children under 12 years • ask a doctor</p> <p>Famotidine (active ingredient) made in Republic of Ireland. Processed in Canada. Distributed by: JOHNSON &amp; JOHNSON CONSUMER DIVISION, Fort Washington, PA 19043 USA. ©2014 McNeil-PPC, with PEPCID and PEPCID AC as registered trademarks of McNeil-PPC.</p>
---	--	---

(b) (4)

Do not use if foil seal under bottle cap printed "SAFETY SEAL" is broken or missing.

**Active ingredient in each tablet**

Purpose	Active ingredient
Acid reducer	Famotidine 20 mg

**Uses**  
• relieves heartburn associated with acid indigestion and sour stomach • prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain food and beverages

**Warnings** Allergy alert: Do not use if you are allergic to famotidine or other acid reducers  
Do not use • if you have trouble or pain swallowing food • vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor. • if you have kidney disease, except under the advice and supervision of a doctor  
• with other acid reducers. Ask a doctor before use if you have • high heartburn over 3 months. This may be a sign of a more serious condition. • heartburn with lightheadedness, sweating, or dizziness • chest pain or shoulder pain with shortness of breath, sweating, pain spreading to arms, neck or shoulders, or lightheadedness • frequent chest pain • frequent wheezing, particularly with heartburn

**Precautions**  
• unexplained weight loss • nausea or vomiting • stomach pain. Stop use and ask a doctor if • your heartburn continues or worsens • you need to take this product for more than 14 days  
If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

**Directions** • adults and children 12 years and over: • to relieve symptoms, swallow 1 tablet with a glass of water. Do not chew. • to prevent symptoms, swallow 1 tablet with a glass of water at any time from 15 to 30 minutes before eating food or drinking beverages that cause heartburn  
• do not use more than 2 tablets in 24 hours  
• children under 12 years: ask a doctor  
Famotidine (active ingredient) made in Republic of Ireland, Processed in Canada  
00000000000000

**Maximum Strength**  
NDC 10337-855-10


**Pepcid<sup>®</sup> AC**  
Famotidine Tablets 20 mg Acid Reducer  
Just One Tablet!  
Prevents & Relieves Heartburn Due to Acid Indigestion  
50 Tablets

EXP: LOT: NDC 10337-855-10


Blister @ 100% Scale




Blister @ 200% Scale




Blister @ 100% Scale




Blister @ 200% Scale



Blister @ 100% Scale



Blister @ 200% Scale



---

**This is a representation of an electronic record that was signed electronically and this page is the manifestation of the electronic signature.**

---

/s/  
-----

VALERIE S PRATT  
06/08/2016