


This label may not be the latest approved by FDA.
For current labeling information, please visit <https://www.fda.gov/drugsatfda>



50 Pouches of
 Due to Acid Indigestion
 Prevents & Relieves Heartburn
 Just One Tablet!
 Acid Reducer
 Famotidine Tablets 20 mg


Pepcid[®] AC

MAXIMUM STRENGTH


actual size

Drug Facts	Purpose
Active ingredient (in each tablet) Famotidine 20 mg	Acid reducer
Uses • relieves heartburn associated with acid indigestion and sour stomach • prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain food and beverages	
Warnings Allergy alert: Do not use if you are allergic to famotidine or other acid reducers	
Do not use • if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor. • with other acid reducers	
Ask a doctor before use if you have • had heartburn over 3 months. This may be a sign of a more serious condition. • heartburn with lightheadedness, sweating, or dizziness • chest pain or shoulder pain with shortness of breath, sweating, pain spreading to arms, neck or shoulders, or lightheadedness • frequent chest pain • frequent wheezing, particularly with heartburn • unexplained weight loss • nausea or vomiting • stomach pain • kidney disease	
Drug Facts (continued) Ask a doctor or pharmacist before use if you are taking a prescription drug. Acid reducers may interact with certain prescription drugs. Stop use and ask a doctor if • your heartburn continues or worsens • you need to take this product for more than 14 days If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)	
Directions • adults and children 12 years and over: • to relieve symptoms, swallow 1 tablet with a glass of water. Do not chew. • to prevent symptoms, swallow 1 tablet with a glass of water at any time from 10 to 60 minutes before eating food or drinking beverages that cause heartburn. • do not use more than 2 tablets in 24 hours • children under 12 years: ask a doctor	
Other information • read the directions and warnings before use • keep the carton. It contains important information. • store at 20°-25°C (68°-77°F) • protect from moisture • do not use if pouch is torn or damaged	
Inactive ingredients carnauba wax, hydroxypropyl cellulose, hypromellose, magnesium stearate, microcrystalline cellulose, pregelatinized starch, talc, titanium dioxide	
Questions or comments? 1-800-755-4008 (toll-free) or 215-273-8755 (collect)	

Famotidine (active ingredient) made in Hungary
 Distributed by: JOHNSON & JOHNSON CONSUMER INC.
 McNeil Consumer Healthcare Division, Fort Washington, PA 19034 USA
 ©McNeil Consumer Pharmaceuticals Co. 2021
 Pepcid[®] and Pepcid AC[®] are registered trademarks of McNeil-PPC.



7 16837 85572 9



MAXIMUM STRENGTH

Pepcid[®] AC

Famotidine Tablets 20 mg
 Acid Reducer

Just One Tablet!
 Prevents & Relieves Heartburn
 Due to Acid Indigestion

**50 Pouches of
 1 Tablet per Pouch
 20 mg each
 (Individual Pouches)**

actual size


NDC 16837-855-52

MAXIMUM STRENGTH

Pepcid[®] AC

Famotidine Tablets 20 mg
 Acid Reducer

Just One Tablet!
 Prevents & Relieves Heartburn
 Due to Acid Indigestion



**50 Pouches of 1 Tablet per Pouch
 20 mg each (Individual Pouches)**

actual size

MAXIMUM STRENGTH

Pepcid[®] AC

Famotidine Tablets 20 mg
 Acid Reducer

Just One Tablet!
 Prevents & Relieves Heartburn
 Due to Acid Indigestion

**50 Pouches of
 1 Tablet per Pouch
 20 mg each
 (Individual Pouches)**

actual size

This is a representation of an electronic record that was signed electronically. Following this are manifestations of any and all electronic signatures for this electronic record.

/s/

NUSHIN F TODD
02/18/2022 12:46:41 PM