

---

## **Drug Facts**

---

<b>Active ingredient (in each brown oval capsule)</b>	<b>Purpose</b>
Solubilized ibuprofen equal to 200 mg ibuprofen..... (present as the free acid and potassium salt)	Pain reliever

---

**Use** treats migraine

---

### **Warnings**

**Allergy alert:** Ibuprofen may cause a severe allergic reaction which may include:

- hives
- facial swelling
- asthma (wheezing)
- shock

**Alcohol warning:** If you consume 3 or more alcoholic drinks every day, ask your doctor whether you should take ibuprofen or other pain relievers/fever reducers. Ibuprofen may cause stomach bleeding.

---

**Do not use** if you have ever had an allergic reaction to any other pain reliever/fever reducer

---

### **Ask a doctor before use if you have**

- never had migraines diagnosed by a health professional
  - a headache that is different from your usual migraines
  - the worst headache of your life
  - fever and stiff neck
  - headaches beginning after or caused by head injury, exertion, coughing or bending
  - experienced your first headache after the age of 50
  - daily headaches
  - a migraine so severe as to require bed rest
  - problems or serious side effects from taking pain relievers or fever reducers
  - stomach pain
- 

### **Ask a doctor or pharmacist before use if you are**

- under a doctor's care for any continuing medical condition
  - taking other drugs on a regular basis
  - taking another product containing ibuprofen, or any other pain reliever/fever reducer
- 

### **Stop use and ask a doctor if**

- an allergic reaction occurs. Seek medical help right away.
  - migraine headache pain is not relieved or gets worse after first dose
  - stomach pain occurs with the use of this product
  - new or unexpected symptoms occur
- 

**If pregnant or breast-feeding**, ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

---

**Directions**

adults:	<ul style="list-style-type: none"><li>• take 2 capsules with a glass of water</li><li>• if symptoms persist or worsen, ask your doctor</li><li>• do not take more than 2 capsules in 24 hours, unless directed by a doctor</li></ul>
under 18 years of age:	<ul style="list-style-type: none"><li>• ask a doctor</li></ul>

**Other information**

- read all directions and warnings before use. Keep carton.
- store at 20 - 25°C (68 - 77°F)
- avoid excessive heat 40°C (above 104°F)

---

**Inactive ingredients** black iron oxide, D&C yellow no. 10, FD&C green no. 3, FD&C red no. 40, gelatin, lecithin, light mineral oil, pharmaceutical glaze, polyethylene glycol, potassium hydroxide, purified water, simethicone, sorbitan, sorbitol

---

**Questions?** Call 1-800-88-ADVIL

---

Do not use if seal under bottle cap imprinted with "Sealed for Your Protection" is broken or missing

Distributed by WHITEHALL-ROBINS HEALTHCARE  
Madison, NJ 07940 Made in USA  
1999 WHITEHALL-ROBINS HEALTHCARE