

BACK



SEAL AREA FOR LOT & EXP  
 25 mm x 4.5 mm

AREA FOR LOT AND EXP  
 TO BE PRINTED IN DARK COLOR  
 26 mm x 5.5 mm

000068381

DRUG FACTS TEXT DEFINED	TYPE SIZE
• HEADINGS	4.5 pt
• SUBHEADINGS/BODY TEXT	4.5 pt
• LEADING	4.5 pt
• BULLETS	3.5 pt
• SPACE BEFORE BULLET	2 ems
• WARNING BOX LINE, HAIRLINES	N/A

**READ AND KEEP CARTON FOR COMPLETE WARNINGS AND INFORMATION**

**LIFT HERE**

<p><b>Drug Facts</b></p> <p><b>Active ingredient (in each capsule)</b> Solubilized ibuprofen (NSAID)*..... Fever reducer (present as the free acid and potassium salt) *nonsteroidal anti-inflammatory drug</p> <p><b>Uses</b></p> <ul style="list-style-type: none"> <li>temporarily relieves minor aches and pains due to: <ul style="list-style-type: none"> <li>headache</li> <li>backache</li> <li>the common cold</li> <li>minor pain of arthritis</li> <li>temporarily reduces fever</li> </ul> </li> <li>toothache</li> <li>menstrual cramps</li> <li>muscular aches</li> </ul> <p><b>Warnings</b></p> <p><b>Allergy alert:</b> Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:</p> <ul style="list-style-type: none"> <li>hives</li> <li>asthma (wheezing)</li> <li>skin reddening</li> <li>facial swelling</li> <li>shock</li> <li>rash</li> <li>blisters</li> </ul>	<p><b>Purpose</b> Pain reliever/fever reducer</p> <p><b>Drug Facts (continued)</b></p> <p>If an allergic reaction occurs, stop use and seek medical help right away.</p> <p><b>Stomach bleeding warning:</b> This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you</p> <ul style="list-style-type: none"> <li>are age 60 or older</li> <li>have had stomach ulcers or bleeding problems</li> <li>take a blood thinning (anticoagulant) or steroid drug</li> <li>take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)</li> <li>have 3 or more alcoholic drinks every day while using this product</li> <li>take more or for a longer time than directed</li> </ul> <p><b>Heart attack and stroke warning:</b> NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.</p>
--	---

PANEL 1

<p><b>Drug Facts (continued)</b></p> <p><b>Do not use</b></p> <ul style="list-style-type: none"> <li>if you have ever had an allergic reaction to any other pain reliever/fever reducer</li> <li>right before or after heart surgery</li> </ul> <p><b>Ask a doctor before use if</b></p> <ul style="list-style-type: none"> <li>stomach bleeding warning applies to you</li> <li>you have problems or serious side effects from taking pain relievers or fever reducers</li> <li>you have a history of stomach problems, such as heartburn</li> <li>you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke</li> <li>you are taking a diuretic</li> </ul> <p><b>Ask a doctor or pharmacist before use if you are</b></p> <ul style="list-style-type: none"> <li>under a doctor's care for any serious condition</li> <li>taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin</li> <li>taking any other drug</li> </ul> <p><b>When using this product</b></p> <ul style="list-style-type: none"> <li>take with food or milk if stomach upset occurs</li> </ul>	<p><b>Drug Facts (continued)</b></p> <p><b>Stop use and ask a doctor if</b></p> <p>you experience any of the following signs of stomach bleeding:</p> <ul style="list-style-type: none"> <li>feel faint</li> <li>vomit blood</li> <li>have bloody or black stools</li> <li>have stomach pain that does not get better</li> </ul> <p>you have symptoms of heart problems or stroke:</p> <ul style="list-style-type: none"> <li>chest pain</li> <li>trouble breathing</li> <li>weakness in one part or side of body</li> <li>slurred speech</li> <li>leg swelling</li> </ul> <p>pain gets worse or lasts more than 10 days</p> <p>fever gets worse or lasts more than 3 days</p> <p>redness or swelling is present in the painful area</p> <p>any new symptoms appear</p>
---	---

PANEL 2

<p><b>Drug Facts (continued)</b></p> <p><b>If pregnant or breast-feeding,</b> ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.</p> <p><b>Keep out of reach of children.</b> In case of overdose, get medical help or contact a Poison Control Center right away.</p> <p><b>Directions</b></p> <ul style="list-style-type: none"> <li>do not take more than directed</li> <li>the smallest effective dose should be used</li> <li>adults and children 12 years and over: take 1 capsule every 4 to 6 hours while symptoms persist</li> <li>if pain or fever does not respond to 1 capsule, 2 capsules may be used</li> <li>do not exceed 6 capsules in 24 hours, unless directed by a doctor</li> <li>children under 12 years: ask a doctor</li> </ul>	<p><b>Drug Facts (continued)</b></p> <p><b>Other information</b></p> <ul style="list-style-type: none"> <li>each capsule contains: potassium 20 mg</li> <li>read all warnings and directions before use. Keep carton.</li> <li>store at 20-25°C (68-77°F)</li> <li>avoid excessive heat above 40°C (104°F)</li> </ul> <p><b>Inactive ingredients</b> FD&amp;C green no. 3, gelatin, lecithin (soybean), medium-chain triglycerides, pharmaceutical ink, polyethylene glycol, potassium hydroxide, purified water, sorbitol sorbitan solution</p> <p><b>Questions or comments?</b> call toll free 1-800-88-ADVIL</p> <p>000068383</p>
--	--

BASE PANEL

000068383

DRUG FACTS TEXT DEFINED	TYPE SIZE
• DRUG FACTS TITLE	9 pt
• DRUG FACTS CONTINUED	8 pt
• HEADINGS	8 pt
• SUBHEADINGS/BODY TEXT	6 pt
• LEADING	6.5 pt
• # OF CHARACTERS PER INCH	<39
• BULLETS	5 pt
• SPACE BEFORE BULLET	2 ems
• BARLINES, HAIRLINES	1.5 pt, .5 pt
• SPACE BETWEEN HAIRLINES AND BOX END	2 spaces

UNPRINTED AREA  
FOR DATE & LOT  
(UNVARNISHED)

PRINTED AREA FOR DATE & LOT  
LASER CODING (VARNISHED)

**READ AND KEEP CARTON FOR COMPLETE WARNINGS AND INFORMATION**

**Drug Facts**  
**Active ingredient (in each capsule)** Solubilized ibuprofen equal to 200 mg ibuprofen (NSAID) \*  
**Purpose** Pain reliever/fever reducer  
**Nonsteroidal anti-inflammatory drug**

**Uses**

- temporarily relieves minor aches and pains due to:
  - headache
  - backache
  - the common cold
  - menstrual cramps
  - muscular aches
- minor pain of arthritis
- temporarily reduces fever

**Warnings**  
**Allergy alert:** ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- dizziness
- fainting

If an allergic reaction occurs, stop use and seek medical help right away.

**Drug Facts (continued)**

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take other drugs containing aspirin or NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

**Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

**Do not use** ■ if you have ever had an allergic reaction to any other pain reliever/fever reducer ■ right before or after heart surgery

**Ask a doctor before use if** ■ you have had an allergic reaction to any applies to you ■ you have indigestion or serious side effects from taking pain relievers or fever reducers ■ you have a history of stomach problems, such as heartburn ■ you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke ■ you are taking a diuretic

**Ask a doctor or pharmacist before use if you are** ■ under a doctor's care for any serious condition ■ taking aspirin for heart attack or stroke, because ibuprofen may decrease the benefit of aspirin ■ taking any other drug

**When using this product**

- take with food or milk, if stomach upset occurs

000068421

**FPO**

3 0573 0169 20 2

**Drug Facts (continued)**

**Stop use and ask a doctor if** ■ you experience any of the following signs of stomach bleeding: ■ feel faint ■ vomit blood ■ have bloody or black stools

- you have stomach pain that does not get better
- chest pain ■ trouble breathing ■ weakness in one part of body ■ slurred speech ■ big swelling
- pain gets worse or lasts more than 10 days ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present in the painful area ■ any new symptoms appear

**If pregnant or breast-feeding,** ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions** ■ **do not take more than directed**

- the smallest effective dose should be used
- adults and children 12 years and over: take 1 capsule every 4 to 6 hours while symptoms persist ■ if pain or fever does not respond to 1 capsule, 2 capsules may be used
- do not exceed 6 capsules in 24 hours, unless directed by a doctor
- children under 12 years: ask a doctor

**Other information**

- each capsule contains: potassium 20 mg

**Do Not Use if seal under bottle cap imprinted with "SEALED FOR YOUR PROTECTION" is broken or missing.**

**Questions or comments?** call toll free 1-800-88-ADVIL

**Inactive ingredients:** FD&C green no. 3, gelatin, lecithin (polyethylene glycol, potassium hydroxide, purified water, sorbitol sorbitan solution)

**Read all warnings and directions before use. Keep carton.**

- store at 20-25°C (68-77°F)
- avoid excessive heat above 40°C (104°F)

**Drug Facts (continued)**

**For most recent product information, visit [www.advil.com](http://www.advil.com)**

**Dist. by:** GSK CH, Warren, NJ 07059

**©2021 GSK or licensor**

**Trademarks owned or licensed by GSK**

**LIQUI-GELS® is a trademark or registered trademark of Caballini Pharma Solutions.**

000068421

DRUG FACTS MODIFIED TEXT DEFINED	TYPE SIZE
• DRUG FACTS TITLE	8 pt
• DRUG FACTS CONTINUED	7 pt
• HEADINGS	7 pt
• SUBHEADINGS/BODY TEXT	6 pt
• LEADING	6.25 pt
• # OF CHARACTERS PER INCH	<39
• BULLETS	5 pt
• SPACE BEFORE BULLET	2 ems
• BARLINES, HAIRLINES	1.5 pt, .5 pt
• SPACE BETWEEN HAIRLINES AND BOX END	2 spaces

Reference ID: 4926509

UNPRINTED AREA  
FOR DATE & LOT  
(UNVARNISHED)

PRINTED AREA FOR DATE & LOT  
LASER CODING (VARNISHED)

**READ AND KEEP CARTON FOR COMPLETE WARNINGS AND INFORMATION**

**Drug Facts**  
**Active ingredient (in each capsule)** Solubilized ibuprofen equal to 200 mg ibuprofen (NSAID)<sup>†</sup> Present as the free acid and potassium salt. Nonsteroidal anti-inflammatory drug.

**Purpose** Pain reliever/fever reducer.

**Uses**  
Temporarily relieves minor aches and pains due to:  

- headache
- backache
- menstrual cramps
- the common cold
- muscular aches
- minor pain of arthritis

Temporarily reduces fever.

**Warnings**  
**Major alert:** Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:  

- hives
- facial swelling
- chest pain
- skin redness
- rash
- dizziness
- asthma (wheezing)

If an allergic reaction occurs, stop use and seek medical help right away.

**Drug Facts (continued)**  
**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:  

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- have more or for a longer time than directed

**Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

**Do not use** if you have ever had an allergic reaction to any other pain reliever/fever reducer, right before or after heart surgery.

**Ask a doctor before use if:**  

- stomach bleeding warning applies to you
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic

**Ask a doctor or pharmacist before use if you are:**  

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because ibuprofen may decrease his benefit
- taking any other drug

**When using this product:**  

- take with food or milk if stomach upset occurs

**Drug Facts (continued)**  
**Stop use and ask a doctor if:**  

- you experience any of the following signs of stomach bleeding:  
  - feel faint
  - vomit blood
  - have blood or black stools
- have stomach pain that does not get better
- you have symptoms of heart problems or stroke:  
  - chest pain
  - trouble breathing
  - weakness in one part of your body
  - slurred speech
  - leg swelling
- pain or loss more than 10 days
- fever gets worse or lasts more than 3 days
- rashes or swellings present in the genital area
- any new symptoms appear

**Pregnant or breast-feeding:** ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions:** Do not take more than directed.  
**the smallest effective dose should be used.**  

- adults and children 12 years and over: take 1 capsule every 4 to 6 hours while symptoms persist
- if pain or fever does not respond to 1 capsule, 2 capsules may be used
- do not exceed 6 capsules in 24 hours, unless directed by a doctor
- children under 12 years: ask a doctor

**Other information**  

- each capsule contains potassium 20 mg

000071083

FPO

3 0573 0169 22 6

**Drug Facts (continued)**  
**Inactive ingredients:** FD&C green no. 3, gelatin, lecithin (soybean), medium-chain triglycerides, pharmaceutical ink, polyethylene glycol, potassium hydroxide, purified water, sorbitol sorbitan solution.

**Questions or comments?** call toll free 1-800-88-ADVIL.

**Do Not Use if seal under bottle cap imprinted with "SEALED for YOUR PROTECTION" is broken or missing.**

For most recent product information, visit [www.AdvIL.com](http://www.AdvIL.com).  
 Dist. by: GSK CH, Warren, NJ 07059  
 ©2021 GSK or licensee. All trademarks owned or licensed by GSK registered trademark of Ciplaent Pharma Solutions.  
 Made in Argentina.

50% MORE FREE

10 Free LIQUI-GELS®

Advil®

LIQUI-GELS®

Solubilized Ibuprofen Capsules, 200 mg  
Pain Reliever/Fever Reducer (NSAID)

28 30

Liquid Filled Capsules

000071083

DRUG FACTS MODIFIED TEXT DEFINED	TYPE SIZE
• DRUG FACTS TITLE	8 pt
• DRUG FACTS CONTINUED	7 pt
• HEADINGS	7 pt
• SUBHEADINGS/BODY TEXT	6 pt
• LEADING	6.25 pt
• # OF CHARACTERS PER INCH	<39
• BULLETS	5 pt
• SPACE BEFORE BULLET	2 ems
• BARLINES, HAIRLINES	1.5 pt, .5 pt
• SPACE BETWEEN HAIRLINES AND BOX END	2 spaces

**READ AND KEEP CARTON FOR COMPLETE WARNINGS AND INFORMATION**

**Drug Facts**  
**Active ingredient (in each capsule)** Solubilized ibuprofen equal to 200 mg ibuprofen (NSAID)\*  
**Purpose** Pain reliever/fever reducer (present as the free acid and potassium salt)  
\*nonsteroidal anti-inflammatory drug

**Uses**  
temporarily relieves minor aches and pains due to:  

- headache
- toothache
- backache
- menstrual cramps
- the common cold
- muscular aches
- minor pain of arthritis
- temporarily reduces fever

**Warnings**  
**Allergy alert:** Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin.  
Symptoms may include:  

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin redness
- rash
- dizziness

If an allergic reaction occurs, stop use and seek medical help right away.  
**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding.  
The chance is higher if you:  

- are age 60 or older
- have a stomach ulcer or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

**Drug Facts (continued)**  
**Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.  
**Do not use**  

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

**Ask a doctor before use if:**  

- stomach bleeding warning applies to you
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic

**Ask a doctor or pharmacist before use if you are:**  

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- taking any other drug

**When using this product:**  

- take with food or milk if stomach upset occurs

**Stop use and ask a doctor if:**  

- you experience any of the following signs of stomach bleeding:  
  - feel faint
  - vomit blood
  - have bloody or black stools
- you have stomach pain that does not get better
- you have symptoms of heart problems or stroke:  
  - chest pain
  - trouble breathing
  - weakness in one part or side of body
  - slurred speech
  - leg swelling
- your gums worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

**Drug Facts (continued)**  
**Impregnant or breast-feeding:** ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.  
**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**  

- do not take more than directed
- the smallest effective dose should be used
- adults and children 12 years and over: take 1 capsule every 4 to 6 hours while symptoms persist
- if pain or fever does not respond to 1 capsule, 2 capsules may be used
- do not exceed 6 capsules in 24 hours, unless directed by a doctor
- children under 12 years: ask a doctor

**Other information**  

- each capsule contains potassium 20 mg
- read all warnings and directions before use. Keep carton.
- store at 20-25°C (68-77°F)
- avoid excessive heat above 40°C (104°F)

**Inactive ingredients**  
FD&C green no. 3, gelatin, lecithin (soybean), medium-chain triglycerides, pharmaceutical ink, polyethylene glycol, potassium hydroxide, purified water, sorbitol sorbitan solution

**Questions or comments?**  
call toll free 1-800-88-40111

**Do Not Use if seal under bottle cap imprinted with "SEALED FOR YOUR PROTECTION" is broken or missing.**  
For most recent product information, visit [www.Advii.com](http://www.Advii.com)  
Dist. by: ESK Consumer Healthcare, Warren, NJ 07059  
©2021 ESK Consumer Healthcare. All rights reserved.  
Liqui-Gels is a trademark or registered trademark of Galderma Pharma Solutions.  
000071101

PRINTED AREA FOR DATE & LOT  
LASER CODING (VARNISHED)

UNPRINTED AREA  
FOR DATE & LOT  
(UNVARNISHED)

3 0573 0169 30 1

**FPO**

**Advii**  
**LIQUI·GELS**  
Solubilized Ibuprofen Capsules, 200 mg  
Pain Reliever/Fever Reducer (NSAID)  
40 Liquid Filled Capsules

000071101

DRUG FACTS TEXT DEFINED	TYPE SIZE
• DRUG FACTS TITLE	9 pt
• DRUG FACTS CONTINUED	8 pt
• HEADINGS	8 pt
• SUBHEADINGS/BODY TEXT	6 pt
• LEADING	6.5 pt
• # OF CHARACTERS PER INCH	<39
• BULLETS	5 pt
• SPACE BEFORE BULLET	2 ems
• BARLINES, HAIRLINES	1.5 pt, .5 pt
• SPACE BETWEEN HAIRLINES AND BOX END	2 spaces

**READ AND KEEP CARTON FOR COMPLETE WARNINGS AND INFORMATION**

**Drug Facts**

**Active ingredient (in each capsule)** Solubilized ibuprofen equal to 200 mg ibuprofen (NSAID)\*

**Purpose** Pain reliever/fever reducer (nonsteroidal anti-inflammatory drug)

**Uses**

- temporarily relieves minor aches and pains due to:
  - headache
  - backache
  - the common cold
  - minor pain of arthritis
- temporarily reduces fever

**Warnings**

**Allergy alert:** Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- stomach pain
- stomach bleeding
- skin redness
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away!

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- are age 60 or older
- have had stomach ulcers or bleeding problems
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product

**Hear attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

0 4 0 5 7 3 0 1 6 9 4 0 3

FPO

00001770

**Drug Facts (continued)**

**Do not use**

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

**Ask a doctor before use if**

- stomach bleeding warning applies to you
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have a history of stomach problems, such as heartburn, kidney disease, asthma, or had a stroke
- you are taking a diuretic

**Ask a doctor or pharmacist before use if you are**

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- taking any other drug

**When using this product:**

- take with food or milk if stomach upset occurs

**Stop use and ask a doctor if**

- you experience any of the following signs of stomach bleeding:
  - red or black stools
  - vomit blood
  - have bloody or black stools
  - have stomach pain that does not get better
  - you have symptoms of heart problems or stroke:
    - chest pain
    - trouble breathing
    - weakness in one part of side of body
    - slurred speech
    - leg swelling
  - pain gets worse or lasts more than 10 days
  - ever gets worse or lasts more than 3 days
  - redness or swelling is present in the painful area
  - any new symptoms appear

**If pregnant or breast-feeding, ask a health professional before use.** It is especially important not to use ibuprofen a 20 weeks or later in pregnancy, unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Do Not Use if seal under bottle cap imprinted with "SEALED for YOUR PROTECTION" is broken or missing.**

For most recent product information, visit [www.Advii.com](http://www.Advii.com)

Distributed by: GSK Consumer Healthcare, Warren, NJ 07059

©2021 GSK group of companies or its licensors

Trademarks are owned by or licensed to the GSK group of companies

LIQUI-GELS® is a trademark or registered trademark of Catalent Pharma Solutions.

**Drug Facts (continued)**

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**

- do not take more than directed
- the smallest effective dose should be used
- adults and children 12 years and over: take 1 capsule every 4 to 6 hours while symptoms persist
- if pain or fever does not respond to 1 capsule, 2 capsules may be used
- do not exceed 6 capsules in 24 hours, unless directed by a doctor
- children under 12 years: ask a doctor

**Other information**

- each capsule contains: potassium 20 mg
- read all warnings and directions before use.
- keep carton.
- store at 20-25°C (68-77°F)
- avoid excessive heat above 40°C (104°F)

**Inactive ingredients**

F3&C green no. 3, gelatin, lecithin (soybean), medium-chain triglycerides, pharmaceutical ink, polyethylene glycol, potassium hydroxide, purified water, sorbitol sorbitan solution

**Questions or comments?**

call toll free 1-800-38-ADVIL

000071770

DRUG FACTS TEXT DEFINED	TYPE SIZE
• DRUG FACTS TITLE	9.25 pt
• DRUG FACTS CONTINUED	8.25 pt
• HEADINGS	8.25 pt
• SUBHEADINGS/BODY TEXT	6.25 pt
• LEADING	6.5 pt
• # OF CHARACTERS PER INCH	<39
• BULLETS	5.25 pt
• SPACE BEFORE BULLET	2 ems
• BARLINES, HAIRLINES	1.5 pt, .5 pt
• SPACE BETWEEN HAIRLINES AND BOX END	2 spaces



UNPRINTED AREA FOR DATE & LOT CODING (UNVARNISHED)  
PRINTED AREA FOR DATE & LOT LASER CODING (VARNISHED)

000071632

DRUG FACTS TEXT DEFINED	TYPE SIZE
• DRUG FACTS TITLE	9.25 pt
• DRUG FACTS CONTINUED	8.25 pt
• HEADINGS	8.25 pt
• SUBHEADINGS/BODY TEXT	6.25 pt
• LEADING	6.5 pt
• # OF CHARACTERS PER INCH	<39
• BULLETS	5.25 pt
• SPACE BEFORE BULLET	2 ems
• BARLINES, HAIRLINES	1.5 pt, .5 pt
• SPACE BETWEEN HAIRLINES AND BOX END	2 spaces



TOP  
PANEL 1

**20 FREE LIQUI-GELS** Easy Open ARTHRITIS CAP

THIS PACKAGE FOR HOUSEHOLDS WITHOUT YOUNG CHILDREN

**Advil LIQUI-GELS**

Solubilized Ibuprofen Capsules, 200 mg Pain Reliever/Fever Reducer (NSAID)

**160-180 Liquid Filled Capsules**

**Drug Facts**

**Active ingredient (in each capsule) Purpose**  
Solubilized ibuprofen equal to 200 mg ibuprofen (NSAID)\* .....Pain reliever/Fever reducer (present as the free acid and potassium salt) \*nonsteroidal anti-inflammatory drug

**Uses**

- temporarily relieves minor aches and pains due to:
  - headache
  - toothache
  - backache
  - menstrual cramps
  - the common cold
  - muscular aches
  - minor pain of arthritis
  - temporarily reduces fever

000071637

L3\_0573\_0149\_91\_4

Do Not Use if seal under bottle cap imprinted with "SEALED for YOUR PROTECTION" is broken or missing.

LIFT HERE for More Drug Facts

FOR LOT & EXPIRATION AREA. UNVARNISHED AND PRINTED IN BLACK OR DARK BLUE.

BASE  
PANEL 3

**Advil**

For most recent product information, visit [www.Advil.com](http://www.Advil.com)  
Distributed by: GSK Consumer Healthcare, Warren, NJ 07059  
©2021 GSK group of companies or its licensor  
Trademarks are owned by or licensed to the GSK group of companies  
LIQUI-GELS® is a trademark or registered trademark of Catalent Pharma Solutions.

**Drug Facts (continued)**

**If pregnant or breast-feeding**, ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.  
**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**

- do not take more than directed
- the smallest effective dose should be used
- adults and children 12 years and over: take 1 capsule every 4 to 6 hours while symptoms persist
- if pain or fever does not respond to 1 capsule, 2 capsules may be used
- do not exceed 6 capsules in 24 hours, unless directed by a doctor
- children under 12 years: ask a doctor

**Other information**

- each capsule contains: potassium 20 mg
- read all warnings and directions before use.
- store at 20-25°C (68-77°F)
- avoid excessive heat above 40°C (104°F)

**Inactive ingredients**  
FD&C green no. 3, gelatin, lecithin (soybean), medium-chain triglycerides, pharmaceutical ink, polyethylene glycol, potassium hydroxide, purified water, sorbitol sorbitan solution

**Questions or comments?** call toll free 1-800-88-ADVIL

RELEASED BACK  
PANEL 2

**Drug Facts (continued)**

**Warnings**

**Allergy alert:** Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

**Stomach bleeding warning:**  
This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

**Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

**Do not use**

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

**Drug Facts (continued)**

**Ask a doctor before use if**

- stomach bleeding warning applies to you
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic

**Ask a doctor or pharmacist before use if you are**

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- taking any other drug

**When using this product**

- take with food or milk if stomach upset occurs

**Stop use and ask a doctor if**

- you experience any of the following signs of stomach bleeding:
  - feel faint
  - vomit blood
  - have bloody or black stools
  - have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
  - chest pain
  - trouble breathing
  - weakness in one part or side of body
  - slurred speech
  - leg swelling
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

000071637

DRUG FACTS TEXT DEFINED	TYPE SIZE
• DRUG FACTS TITLE	9.0 pt
• DRUG FACTS CONTINUED	8.0 pt
• HEADINGS	8.0 pt
• SUBHEADINGS/BODY TEXT	6.0 pt
• LEADING	6.5 pt
• # OF CHARACTERS PER INCH	<39
• BULLETS	5 pt
• SPACE BEFORE BULLET	2 ems
• BARLINES, HAIRLINES	1.5 pt, .5 pt
• SPACE BETWEEN HAIRLINES AND BOX END	2 spaces

**READ AND KEEP CARTON FOR COMPLETE WARNINGS AND INFORMATION**

**Drug Facts**  
**Active Ingredient (in each capsule)** Solubilized Ibuprofen (NSAID) 200 mg Ibuprofen (NSAID) (present as the free acid and potassium salt) \*nonsteroidal anti-inflammatory drug

**Purpose** Pain reliever/fever reducer

**Uses**  
temporarily relieves minor aches and pains due to:  

- headache
- menstrual cramps
- minor pain of arthritis
- toothache
- the common cold
- muscle aches

temporarily reduces fever

**Warnings**  
**Allergy alert:** Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:  

- hives
- skin swelling
- asthma (breathing)
- wheezing
- dizziness
- fainting

If an allergic reaction occurs, stop use and seek medical help right away.  
**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:  

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product

**Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack and stroke. The risk is higher if you use more than directed or for longer than directed.

**Do not use**  

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

**Ask a doctor before use if**  

- stomach bleeding warning applies to you
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have a history of stomach problems, such as heartburn
- you are taking a diuretic
- you are taking a diuretic

**Ask a doctor or pharmacist before use if you are**  

- under a doctor's care for any serious condition
- taking aspirin
- taking aspirin
- taking any other drug

**Drug Facts (continued)**

**When using this product**  

- take with food or milk if stomach upset occurs

**Stop use and ask a doctor if**  

- you experience any of the following signs of stomach bleeding:  
  - heart burn
  - vomit blood
  - have bloody or black stools
- have stomach pain that does not get better
- you have symptoms of heart problems or stroke:  
  - chest pain
  - trouble breathing
  - weakness in one part or side of body
  - slurred speech
  - eye swelling
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- rash or skin sores appear in the painful area
- any new symptoms appear

**If pregnant or breast-feeding,** ask a health professional before use. It's especially important not to use ibuprofen at 20 weeks or later in pregnancy unless directly directed to do so by a doctor because it may cause problems in the fetus. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**  

- do not take more than directed
- the smallest effective dose should be used
- adults and children 12 years and over: take 1 capsule every 4 to 6 hours while symptoms persist. capsule, 2 capsules may be used if needed. Do not take more than 6 capsules in 24 hours. Unless directed by a doctor
- children under 12 years: ask a doctor

**Other information**  

- each capsule contains: potassium 20 mg
- read all warnings and directions before use. Keep carton.
- store at 20-25°C (68°-77°F)
- avoid excessive heat above 40°C (104°F)

**Inactive ingredients:** FD-C green no. 3, gelatin, behenyl sorbitan, medium-chain triglycerides, pharmaceutical ink, polyethylene glycol, potassium hydroxide, purified water, sorbitol sorbitan solution

**Questions or comments?** call toll free 1-800-86-ADVIL

**Do Not Use If you see under bottle cap imprinted with "SEALED FOR YOUR PROTECTION" is broken or missing.**

For most recent product information, visit [www.Advii.com](http://www.Advii.com)  
Distributed by GSK Consumer Healthcare Warren, NJ 07059  
©2021 GSK group of companies or its licensor  
LIQUI-GEL.S® is a trademark or registered trademark of Catalent Pharma Solutions.

**UNPRINTED AREA FOR DATE & LOT LASER CODING (UNVARNISHED)**

000071634

3 0573 0169 51 6

FPO

000071634

**PRINTED AREA FOR DATE & LOT LASER CODING (VARNISHED)**

000071634

DRUG FACTS TEXT DEFINED	TYPE SIZE
• DRUG FACTS TITLE	9.0 pt
• DRUG FACTS CONTINUED	8.0 pt
• HEADINGS	8.0 pt
• SUBHEADINGS/BODY TEXT	6.0 pt
• LEADING	6.5 pt
• # OF CHARACTERS PER INCH	<39
• BULLETS	5. pt
• SPACE BEFORE BULLET	2 ems
• BARLINES, HAIRLINES	1.5 pt, .5 pt
• SPACE BETWEEN HAIRLINES AND BOX END	2 spaces

TOP LAYER FACE

**GREAT VALUE!**

# Advil<sup>®</sup>

## LIQUI-GELS<sup>®</sup>

Solubilized Ibuprofen Capsules, 200 mg  
Pain Reliever/Fever Reducer (NSAID)

**200**  
Liquid Filled Capsules

**GREAT VALUE!**

# Advil<sup>®</sup>

## LIQUI-GELS<sup>®</sup>

Solubilized Ibuprofen Capsules, 200 mg  
Pain Reliever/Fever Reducer (NSAID)

**Drug Facts**

**Active ingredient (in each capsule) Purpose**  
Solubilized ibuprofen equal to 200 mg ibuprofen (NSAID)\* .....Pain reliever/Fever reducer (present as the free acid and potassium salt)  
\*nonsteroidal anti-inflammatory drug

**Uses**

- temporarily relieves minor aches and pains due to:
  - headache
  - toothache
  - backache
  - menstrual cramps
  - the common cold
  - muscular aches
  - minor pain of arthritis
  - temporarily reduces fever

000071158

Do Not Use if seal under bottle cap imprinted with "SEALED for YOUR PROTECTION" is broken or missing.

**FPO**

3 0573 0169 49 3

UNVARNISHED & PRINTED AREA FOR LOT & EXP. (LINES DO NOT PRINT)

TOP LAYER BACK

**Drug Facts (continued)**

**Warnings**

**Allergy alert:** Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

**Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

**Do not use**

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

**Ask a doctor before use if**

- stomach bleeding warning applies to you
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic

**Drug Facts (continued)**

**Ask a doctor or pharmacist before use if you are**

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- taking any other drug

**When using this product**

- take with food or milk if stomach upset occurs

**Stop use and ask a doctor if**

- you experience any of the following signs of stomach bleeding:
  - feel faint
  - vomit blood
  - have bloody or black stools
  - have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
  - chest pain
  - trouble breathing
  - weakness in one part or side of body
  - slurred speech
  - leg swelling
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

**If pregnant or breast-feeding,** ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

BASE LAYER

**Drug Facts (continued)**

**Directions**

- do not take more than directed
- the smallest effective dose should be used
- adults and children 12 years and over: take 1 capsule every 4 to 6 hours while symptoms persist
- if pain or fever does not respond to 1 capsule, 2 capsules may be used
- do not exceed 6 capsules in 24 hours, unless directed by a doctor
- children under 12 years: ask a doctor

**Other information**

- each capsule contains: potassium 20 mg
- read all warnings and directions before use.
- store at 20-25°C (68-77°F)
- avoid excessive heat above 40°C (104°F)

**Inactive ingredients**  
FD&C green no. 3, gelatin, lecithin (soybean), medium-chain triglycerides, pharmaceutical ink, polyethylene glycol, potassium hydroxide, purified water, sorbitol sorbitan solution

**Questions or comments?** call toll free 1-800-88-ADVIL

# Advil<sup>®</sup>

For most recent product information, visit [www.Advil.com](http://www.Advil.com)  
Distributed by: GSK Consumer Healthcare, Warren, NJ 07059  
©2021 GSK group of companies or its licensor  
Trademarks are owned by or licensed to the GSK group of companies  
LIQUI-GELS<sup>®</sup> is a trademark or registered trademark of Catalent Pharma Solutions.

000071158

DRUG FACTS TEXT DEFINED	TYPE SIZE
• DRUG FACTS TITLE	9.0 pt
• DRUG FACTS CONTINUED	8.0 pt
• HEADINGS	8.0 pt
• SUBHEADINGS/BODY TEXT	6.0 pt
• LEADING	6.5 pt
• # OF CHARACTERS PER INCH	<39
• BULLETS	5 pt
• SPACE BEFORE BULLET	2 ems
• BARLINES, HAIRLINES	1.5 pt, .5 pt
• SPACE BETWEEN HAIRLINES AND BOX END	2 spaces

TOP  
PANEL 1

FOR LOT & EXPIRATION AREA.  
UNVARNISHED AND PRINTED  
IN BLACK OR DARK BLUE.

BASE  
PANEL 3

RELEASED BACK  
PANEL 2

000068422

DRUG FACTS TEXT DEFINED	TYPE SIZE
• DRUG FACTS TITLE	9.0 pt
• DRUG FACTS CONTINUED	8.0 pt
• HEADINGS	8.0 pt
• SUBHEADINGS/BODY TEXT	6.0 pt
• LEADING	6.5 pt
• # OF CHARACTERS PER INCH	<39
• BULLETS	5 pt
• SPACE BEFORE BULLET	2 ems
• BARLINES, HAIRLINES	1.5 pt, .5 pt
• SPACE BETWEEN HAIRLINES AND BOX END	2 spaces





**READ AND KEEP CARTON FOR COMPLETE WARNINGS AND INFORMATION**

**Drug Facts**

**Active ingredient (in each capsule)** Solubilized ibuprofen equal to 200 mg ibuprofen (NSAID)\* (present as the free acid and potassium salt) (nonsteroidal anti-inflammatory drug)

**Purpose**

**Uses**  
 ■ temporarily relieves minor aches and pains due to:  
 ■ headache ■ toothache ■ backache  
 ■ menstrual cramps ■ the common cold  
 ■ muscular aches ■ minor pain of arthritis  
 ■ temporarily reduces fever

**Warnings**

**Allergy alert:** Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:  
 ■ hives ■ facial swelling ■ asthma (wheezing)  
 ■ shock ■ skin reddening ■ rash ■ blisters  
 If an allergic reaction occurs, stop use and seek medical help right away.  
**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you  
 ■ are age 60 or older ■ have had stomach ulcers or bleeding problems  
 ■ take a blood thinning (anticoagulant) or steroid drug  
 ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)  
 ■ have 3 or more alcoholic drinks every day while using this product  
 ■ take more or for a longer time than directed  
**Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.  
**Do not use**  
 ■ if you have ever had an allergic reaction to any other pain reliever/fever reducer  
 ■ right before or after heart surgery  
**Ask a doctor before use if** ■ stomach bleeding warning applies to you  
 ■ you have problems or serious side effects from taking pain relievers or fever reducers  
 ■ you have a history of stomach problems, such as heartburn  
 ■ you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke  
 ■ you are taking a diuretic

**Drug Facts (continued)**

**Ask a doctor or pharmacist before use if you are**  
 ■ under a doctor's care for any serious condition  
 ■ taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin  
 ■ taking any other drug

**When using this product**  
 ■ take with food or milk if stomach upset occurs

**Stop use and ask a doctor if**  
 ■ you experience any of the following signs of stomach bleeding:  
 ■ feel faint ■ vomit blood ■ have bloody or black stools  
 ■ have stomach pain that does not get better  
 ■ you have symptoms of heart problems or stroke:  
 ■ chest pain ■ trouble breathing  
 ■ weakness in one part or side of body  
 ■ slurred speech ■ leg swelling  
 ■ pain gets worse or lasts more than 10 days  
 ■ fever gets worse or lasts more than 3 days  
 ■ redness or swelling is present in the painful area  
 ■ any new symptoms appear

**If pregnant or breast-feeding,** ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.  
**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**

■ do not take more than directed  
 ■ the smallest effective dose should be used  
 ■ adults and children 12 years and over: take 1 capsule every 4 to 6 hours while symptoms persist  
 ■ if pain or fever does not respond to 1 capsule, 2 capsules may be used  
 ■ do not exceed 6 capsules in 24 hours, unless directed by a doctor  
 ■ children under 12 years: ask a doctor

**Other information** ■ each capsule contains: potassium 20 mg  
 ■ read all warnings and directions before use. Keep carton.  
 ■ store at 20-25°C (68-77°F)  
 ■ avoid excessive heat above 40°C (104°F)

**Inactive ingredients** FD&C green no. 3, gelatin, lecithin (soybean), medium-chain triglycerides, pharmaceutical ink, polyethylene glycol, potassium hydroxide, purified water, sorbitol sorbitan solution

**Questions or comments?** call toll free 1-800-88-ADVIL

Do Not Use if seal under bottle cap imprinted with "SEALED for YOUR PROTECTION" is broken or missing.

For most recent product information, visit [www.Advil.com](http://www.Advil.com)  
 Distributed by: GSK Consumer Healthcare, Warren, NJ 07059  
 ©2021 GSK group of companies or its licensor  
 Trademarks are owned by or licensed to the GSK group of companies  
 LIQUI-GELS® is a trademark or registered trademark of Catalent Pharma Solutions.



000071631

PRINTED AREA FOR DATE & LOT  
LASER CODING (VARNISHED)

000071631

DRUG FACTS TEXT DEFINED	TYPE SIZE
• DRUG FACTS TITLE	9 pt
• DRUG FACTS CONTINUED	8 pt
• HEADINGS	8 pt
• SUBHEADINGS/BODY TEXT	6 pt
• LEADING	6.5 pt
• # OF CHARACTERS PER INCH	<39
• BULLETS	5 pt
• SPACE BEFORE BULLET	2 ems
• BARLINES, HAIRLINES	1.5 pt, .5 pt
• SPACE BETWEEN HAIRLINES AND BOX END	2 spaces

**READ AND KEEP CARTON FOR COMPLETE WARNINGS AND INFORMATION**

**Drug Facts**  
*(Active ingredient in each capsule)*  
Solubilized ibuprofen equal to 200 mg ibuprofen (NSAID) present as the free acid and potassium salt, nonsteroidal anti-inflammatory drug

**Purpose**  
Pain reliever/fever reducer

**Uses**

- temporarily relieves minor aches and pains due to:
  - headache
  - backache
  - muscular cramps
  - the common cold
  - muscular aches
- minor pain of arthritis
- temporarily reduces fever

**Warnings**  
Always alert: ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:
 

- hives
- facial swelling
- swelling (swelling)
- skin redness
- stomach pain
- stomach ulcers
- stomach bleeding
- stomach ulcers
- stomach pain
- stomach ulcers

 If an allergic reaction occurs, stop use and seek medical help right away.

**Do Not Use if seal under bottle cap imprinted with "SEALED FOR YOUR PROTECTION" is broken or missing.**

**Questions or comments?** call toll free 1-800-88-ADVIL.

**Inactive ingredients**  
FD&C green no. 3, gelatin, medium-chain triglycerides, pharmaceutical ink, polyethylene glycol, potassium hydroxide, purified water, sorbitol sorbitan solution.

**Other information** ■ each capsule contains potassium 20 mg ■ read all warnings and directions before use. Keep carton. ■ store at 20-25°C (68-77°F) ■ avoid excessive heat above 40°C (104°F)

**Drug Facts (continued)**

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take other drugs containing aspirin or other NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed

**Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

**Do not use** ■ if you have ever had an allergic reaction to any other pain reliever/fever reducer ■ right before or after heart surgery

**Ask a doctor before use if** ■ stomach bleeding warning applies to you ■ you have problems or serious side effects from taking pain relievers or fever reducers ■ you have a history of stomach problems, such as heartburn ■ you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke ■ you are taking a diuretic. ■ you are taking aspirin ■ you are taking aspirin for heart attack or stroke, because ibuprofen may decrease the benefit of aspirin ■ taking any other drug

**When using this product** ■ take with food or milk if stomach upset occurs

**Do Not Use if seal under bottle cap imprinted with "SEALED FOR YOUR PROTECTION" is broken or missing.**

**Questions or comments?** call toll free 1-800-88-ADVIL.

**Inactive ingredients**  
FD&C green no. 3, gelatin, medium-chain triglycerides, pharmaceutical ink, polyethylene glycol, potassium hydroxide, purified water, sorbitol sorbitan solution.

**Other information** ■ each capsule contains potassium 20 mg ■ read all warnings and directions before use. Keep carton. ■ store at 20-25°C (68-77°F) ■ avoid excessive heat above 40°C (104°F)

**Drug Facts (continued)**

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take other drugs containing aspirin or other NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed

**Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

**Do not use** ■ if you have ever had an allergic reaction to any other pain reliever/fever reducer ■ right before or after heart surgery

**Ask a doctor before use if** ■ stomach bleeding warning applies to you ■ you have problems or serious side effects from taking pain relievers or fever reducers ■ you have a history of stomach problems, such as heartburn ■ you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke ■ you are taking a diuretic. ■ you are taking aspirin ■ you are taking aspirin for heart attack or stroke, because ibuprofen may decrease the benefit of aspirin ■ taking any other drug

**When using this product** ■ take with food or milk if stomach upset occurs

**Stop use and ask a doctor if** ■ you experience any of the following signs of stomach bleeding: ■ feel faint ■ vomit blood ■ have bloody or black stools ■ have stomach pain that does not get better ■ have symptoms of heart problems or stroke: ■ chest pain ■ trouble breathing ■ weakness in one part or side of body ■ slurred speech ■ egg swelling ■ pain gets worse or lasts more than 3 days ■ redness or swelling is present in the painful area ■ any new symptoms appear

**If pregnant or breast-feeding,** ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**  
■ do not take more than directed ■ the smallest effective dose should be used ■ adults and children 12 years and over: take 1 capsule every 4 to 6 hours while symptoms persist ■ pain or fever does not respond to 1 capsule, 2 capsules may be used ■ do not exceed 6 capsules in 24 hrs, unless directed by a doctor ■ children under 12 years: ask a doctor

LIQUI-GEL® is a trademark or registered trademark of Catalent Pharma Solutions.  
000068424

**FPO**

3 0573 1769 20 3

**UNPRINTED AREA FOR DATE & LOT (UNVARNISHED)**

**PRINTED AREA FOR DATE & LOT LASER CODING (VARNISHED)**

**Advil LIQUI-GELS minis**  
Solubilized Ibuprofen Capsules, 200 mg Pain Reliever/Fever Reducer (NSAID)  
20 Liquid Filled Capsules

\*Compared to Advil® Liqui-Gels®

000068424

MODIFIED DRUG FACTS TEXT DEFINED	TYPE SIZE
• DRUG FACTS TITLE	8 pt
• DRUG FACTS CONTINUED	7 pt
• HEADINGS	7 pt
• SUBHEADINGS/BODY TEXT	6 pt
• LEADING	6.25 pt
• # OF CHARACTERS PER INCH	<39
• BULLETS	5 pt
• SPACE BEFORE BULLET	2 ems
• BARLINES, HAIRLINES	1.5 pt, .5 pt
• SPACE BETWEEN HAIRLINES AND BOX END	2 spaces

BACK





SEAL AREA FOR LOT & EXP  
 25 mm x 4.5 mm

AREA FOR LOT AND EXP  
 TO BE PRINTED IN DARK COLOR  
 26 mm x 5.5 mm

000071419

DRUG FACTS TEXT DEFINED	TYPE SIZE
• HEADINGS	4.55 pt
• SUBHEADINGS/BODY TEXT	4.55 pt
• LEADING	4.6 pt
• BULLETS	3.55 pt
• SPACE BEFORE BULLET	2 ems
• BARLINES, HAIRLINES	N/A


0.375" DIA. HOLE (STRIP OUT)



LOT & EXP AREA

000070109

DRUG FACTS TEXT DEFINED	TYPE SIZE
• DRUG FACTS TITLE	9 pt
• DRUG FACTS CONTINUED	8 pt
• HEADINGS	8 pt
• SUBHEADINGS/BODY TEXT	6 pt
• LEADING	6.5 pt
• # OF CHARACTERS PER INCH	<39
• BULLETS	5 pt
• SPACE BEFORE BULLET	2 ems
• BARLINES, HAIRLINES	1.5 pt, .5 pt
• SPACE BETWEEN HAIRLINES AND BOX END	2 spaces



PRINTED AREA FOR DATE & LOT  
LASER CODING (VARNISHED)

000071492

DRUG FACTS MODIFIED TEXT DEFINED	TYPE SIZE
• DRUG FACTS TITLE	8 pt
• DRUG FACTS CONTINUED	7 pt
• HEADINGS	7 pt
• SUBHEADINGS/BODY TEXT	6 pt
• LEADING	6.25 pt
• # OF CHARACTERS PER INCH	<39
• BULLETS	5 pt
• SPACE BEFORE BULLET	2 ems
• BARLINES, HAIRLINES	1.5 pt, .5 pt
• SPACE BETWEEN HAIRLINES AND BOX END	2 spaces



UNPRINTED AREA FOR DATE & LOT CODING (UNVARNISHED)  
PRINTED AREA FOR DATE & LOT LASER CODING (VARNISHED)

000071414

DRUG FACTS MODIFIED TEXT DEFINED	TYPE SIZE
• DRUG FACTS TITLE	9 pt
• DRUG FACTS CONTINUED	8 pt
• HEADINGS	8 pt
• SUBHEADINGS/BODY TEXT	6 pt
• LEADING	6.5 pt
• # OF CHARACTERS PER INCH	<39
• BULLETS	5 pt
• SPACE BEFORE BULLET	2 ems
• BARLINES, HAIRLINES	1.5 pt, .5 pt
• SPACE BETWEEN HAIRLINES AND BOX END	2 spaces

**READ AND KEEP CARTON FOR COMPLETE WARNINGS AND INFORMATION**

**Drug Facts**

**Active Ingredient (in each capsule)**  
Solubilized ibuprofen equal to 200 mg ibuprofen (NSAID) (present as the free acid and potassium salt) nonsteroidal anti-inflammatory drug

**Purpose**  
Pain reliever/fever reducer

**Uses**  
temporarily relieves minor aches and pains due to:  
 ■ headache  
 ■ backache  
 ■ the common cold  
 ■ minor pain of arthritis  
 ■ temporarily reduces fever

**Warnings**  
**Allergy alert:** Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:  
 ■ hives  
 ■ facial swelling  
 ■ asthma (wheezing)  
 ■ stomach pain  
 ■ rash  
 ■ blisters  
 If an allergic reaction occurs, stop use and seek medical help right away.  
**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you  
 ■ are age 60 or older  
 ■ have had stomach ulcers or bleeding problems  
 ■ take a blood thinning (anticoagulant) or steroid drug  
 ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)  
 ■ have 3 or more alcoholic drinks every day while using this product  
**Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

000071414  
 7 08 6921 1769 80 3  
 FPO

**Drug Facts (continued)**

**Do not use**  
 ■ if you have ever had an allergic reaction to any other pain reliever/fever reducer  
 ■ right before or after heart surgery  
**Ask a doctor before use if**  
 ■ stomach bleeding warning applies to you  
 ■ you have problems or serious side effects from taking pain relievers or fever reducers  
 ■ you have a history of stomach problems, such as heartburn  
 ■ you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke  
 ■ you are taking a diuretic  
**Ask a doctor or pharmacist before use if you are**  
 ■ under a doctor's care for any serious condition  
 ■ taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin  
 ■ taking any other drug

**When using this product**

take with food or milk if stomach upset occurs  
**Stop use and ask a doctor if**  
 ■ you experience any of the following signs of stomach bleeding:  
 ■ feel faint  
 ■ vomit blood  
 ■ have blood or black stools  
 ■ have stomach pain that does not get better  
 ■ you have symptoms of heart problems or stroke:  
 ■ chest pain  
 ■ trouble breathing  
 ■ weakness in one part or side of body  
 ■ slurred speech  
 ■ leg swelling  
 ■ pain gets worse or lasts more than 10 days  
 ■ fever gets worse or lasts more than 3 days  
 ■ redness or swelling is present in the painful area  
 ■ any new symptoms appear

**Drug Facts (continued)**

**If pregnant or breast-feeding,** ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.  
**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**

**do not take more than directed**  
 ■ the smallest effective dose should be used  
 ■ adults and children 12 years and over: take 1 capsule every 4 to 6 hours while symptoms persist  
 ■ if pain or fever does not respond to 1 capsule, 2 capsules may be used  
 ■ do not exceed 6 capsules in 24 hours, unless directed by a doctor  
 ■ children under 12 years: ask a doctor

**Other Information**

each capsule contains: potassium 20 mg  
 ■ read all warnings and directions before use. Keep carton.  
 ■ store at 20-25°C (68-77°F)  
 ■ avoid excessive heat above 40°C (104°F)  
**Inactive ingredients:** FD&C green no. 3, gelatin, medium-chain triglycerides, pharmaceutical ink, polyethylene glycol, potassium hydroxide, purified water, sorbitol sorbitan solution

**Questions or comments?** call toll free 1-800-88-ADVIL

**Do Not Use if seal under bottle cap imprinted with "SEALED FOR YOUR PROTECTION" is broken or missing.**

For most recent product information, visit [www.Advil.com](http://www.Advil.com)  
 Distributed by: GSK Consumer Healthcare, Warren, NJ 07059  
 ©2021 GSK group of companies or its licensor  
 Trademarks are owned by or licensed to the GSK group of companies  
 For US Patent or Application status see [www.products.gsk.com](http://www.products.gsk.com)  
 LIQUI-GELS® is a trademark or registered trademark of Calient Pharma Solutions.

**Smaller Capsule Same Strength\***

# Advil LIQUI-GELS minis

Solubilized Ibuprofen Capsules, 200 mg Pain Reliever/Fever Reducer (NSAID)

80 Liquid Filled Capsules

\*Compared to Advil® Liqui-Gels®



UNPRINTED AREA FOR DATE & LOT CODING (UNVARNISHED)  
PRINTED AREA FOR DATE & LOT LASER CODING (VARNISHED)

**READ AND KEEP CARTON FOR COMPLETE WARNINGS AND INFORMATION**

**Drug Facts**  
**Active Ingredient (in each capsule)** Solubilized ibuprofen equal to 200 mg ibuprofen (NSAID) (present as the free acid and potassium salt) nonsteroidal anti-inflammatory drug

**Purpose** Pain reliever/fever reducer

**Uses**  
temporarily relieves minor aches and pains due to:  

- headache
- backache
- the common cold
- minor pain of arthritis
- temporarily reduces fever

**Warnings**  
**Allergy alert:** Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:  

- hives
- facial swelling
- asthma (wheezing)
- stomach pain
- skin redness
- rash
- blisters

 If an allergic reaction occurs, stop use and seek medical help right away.  
**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:  

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product

**Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

**Drug Facts (continued)**

**Do not use**  

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

**Ask a doctor before use if**  

- stomach bleeding warning applies to you
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have a history of stomach problems, such as heartburn, kidney disease, asthma, or had a stroke
- you are taking a diuretic

**Ask a doctor or pharmacist before use if you are**  

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- taking any other drug

**When using this product**  

- take with food or milk if stomach upset occurs

**Stop use and ask a doctor if**  

- you experience any of the following signs of stomach bleeding:  
  - feel faint
  - vomit blood
  - have bloody or black stools
  - have stomach pain that does not get better
- you have symptoms of heart problems or stroke:  
  - chest pain
  - trouble breathing
  - wheezing in one part or side of body
  - slurred speech
  - leg swelling
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

**Drug Facts (continued)**

**If pregnant or breast-feeding, ask a health professional before use.** It is especially important not to use ibuprofen at 20 weeks or later in pregnancy, unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.  
**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**  

- do not take more than directed
- the smallest effective dose should be used
- adults and children 12 years and over: take 1 capsule every 4 to 6 hours while symptoms persist
- if pain or fever does not respond to 1 capsule, 2 capsules may be used
- do not exceed 6 capsules in 24 hours, unless directed by a doctor
- children under 12 years: ask a doctor

**Other information**  

- each capsule contains: potassium 20 mg
- read all warnings and directions before use. Keep carton.
- store at 20-25°C (68-77°F)
- avoid excessive heat above 40°C (104°F)

**Inactive ingredients**  
 FD&C green no. 3; gelatin, medium-chain triglycerides, pharmaceutical ink, polyethylene glycol, potassium hydroxide, purified water, sorbitol sorbitan solution

**Questions or comments?** call toll free 1-800-86-ADVIL

**Do Not Use if seal under bottle cap imprinted with "SEALED for YOUR PROTECTION" is broken or missing.**

For most recent product information, visit [www.advil.com](http://www.advil.com)  
 Distributed by GSK Consumer Healthcare, Warren, NJ 07059  
 ©2021 GSK group of companies or its licensor  
 Patents: [www.productpatents.com](http://www.productpatents.com)  
 LIQUI-GELS® is a trademark or registered trademark of Galeant Pharma Solutions.

000071417

DRUG FACTS TEXT DEFINED	TYPE SIZE
• DRUG FACTS TITLE	9 pt
• DRUG FACTS CONTINUED	8 pt
• HEADINGS	8 pt
• SUBHEADINGS/BODY TEXT	6 pt
• LEADING	6.5 pt
• # OF CHARACTERS PER INCH	<39
• BULLETS	5 pt
• SPACE BEFORE BULLET	2 ems
• BARLINES, HAIRLINES	1.5 pt, .5 pt
• SPACE BETWEEN HAIRLINES AND BOX END	2 spaces

**READ AND KEEP CARTON FOR COMPLETE WARNINGS AND INFORMATION**

**Drug Facts**  
**Active Ingredient (in each capsule)** Solubilized ibuprofen (NSAID) 200 mg ibuprofen (NSAID) (see important information about NSAIDs on the inside of this anti-inflammatory drug)

**Purpose** Pain reliever/fever reducer (nonsteroidal anti-inflammatory drug)

**Uses** temporarily relieves minor aches and pains due to:  
 ■ headache ■ cold/flu ■ menstrual cramps  
 ■ the common cold ■ muscle aches  
 ■ minor pain of arthritis  
 ■ temporarily reduces fever

**Warnings**  
**Allergy alert:** Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:  
 ■ hives ■ facial swelling ■ asthma (wheezing)  
 ■ stomach pain ■ skin redness  
 ■ rash ■ dizziness

**Caution:** If an allergic reaction occurs, stop use and seek medical help right away. If an allergic reaction occurs, stop use and seek medical help right away.

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:  
 ■ are age 60 or older  
 ■ have had stomach ulcers or bleeding problems  
 ■ take a blood thinning (anticoagulant) or steroid drug  
 ■ take aspirin, ibuprofen, naproxen, or other NSAIDs (aspirin, ibuprofen, naproxen, or other NSAIDs)

**Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

**Do not use**  
 ■ if you have ever had an allergic reaction to any other pain reliever/fever reducer  
 ■ right before or after heart surgery

**Ask a doctor before you use it**  
 ■ stomach bleeding warning applies to you  
 ■ you have problems or serious side effects from taking pain relievers or fever reducers  
 ■ you have a history of stomach problems, such as heartburn  
 ■ you have high blood pressure, heart disease, liver cirrhosis, kidney disease, or had a stroke  
 ■ you are taking a diuretic

**Drug Facts (continued)**

**Ask a doctor or pharmacist before use if you are**  
 ■ under a doctor's care for any serious condition  
 ■ taking aspirin for heart attack or stroke, because ibuprofen may decrease the benefit of aspirin  
 ■ taking any other drug

**When using this product**  
 ■ take with food or milk if stomach upset occurs

**Stop use and ask a doctor if**  
 ■ you experience any of the following signs of stomach bleeding:  
 ■ red, dark, or vomit blood  
 ■ black, tarry stools  
 ■ you have symptoms of heart problems or stroke:  
 ■ chest pain ■ trouble breathing  
 ■ weakness in one part or side of body  
 ■ slurred speech ■ leg swelling  
 ■ pain, numbness, or tingling in the arms or legs  
 ■ stomach pain that does not get better  
 ■ any new symptoms appear

**If pregnant or breast-feeding, ask a health professional before use.** It is especially important not to use ibuprofen at 20 weeks or later in pregnancy, unless explicitly directed to do so by a doctor, because it may cause problems in the unborn child or complications during delivery. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions** ■ **do not take more than directed**  
 ■ the smallest effective dose should be used  
 ■ adults and children 12 years and over: take 1 capsule every 4 to 6 hours while symptoms persist. capsule, 2 capsules may be used  
 ■ do not exceed 6 capsules in 24 hours, unless directed by a doctor  
 ■ children under 12 years: ask a doctor

**Other information** ■ each capsule contains potassium 20 mg

**Inactive ingredients** FD&C green no. 3, gelatin, medium-chain triglycerides, pharmaceutical ink, polyethylene glycol, potassium hydroxide, purified water, sodium stearoyl sulfonate

**Questions or comments?** call toll free 1-800-88-ADVIL

**Do Not Use if seal under bottle cap imprinted with "SEALED FOR YOUR PROTECTION" is broken or missing.**  
 For most recent product information, visit [www.advil.com](http://www.advil.com)  
 Distributed by: GSK Consumer Healthcare, Warren, NJ 07099  
 ©2021 GSK group of companies or its licensor.  
 Trademarks are owned by or licensed to the GSK group of companies  
 US Patent and Application status see [www.advil.com](http://www.advil.com)  
 LIQUI-GEL® is a trademark or registered trademark of Cardinal Health Solutions.

UNPRINTED AREA FOR DATE & LOT LASER CODING (UNVARNISHED)

PRINTED AREA FOR DATE & LOT LASER CODING (VARNISHED)

000071416

3 0573 1769 92 0

FPO

**20 FREE Liquid Filled Capsules**

**Advil LIQUI-GELS minis**

Solubilized Ibuprofen Capsules, 200 mg Pain Reliever/Fever Reducer (NSAID)  
\*Compared to Advil® Liqui-Gels®

**160 180 Liquid Filled Capsules**

**Smaller Capsule Same Strength\***

000071416

DRUG FACTS TEXT DEFINED	TYPE SIZE
• DRUG FACTS TITLE	9 pt
• DRUG FACTS CONTINUED	8 pt
• HEADINGS	8 pt
• SUBHEADINGS/BODY TEXT	6 pt
• LEADING	6.5 pt
• # OF CHARACTERS PER INCH	<39
• BULLETS	5 pt
• SPACE BEFORE BULLET	2 ems
• BARLINES, HAIRLINES	1.5 pt, .5 pt
• SPACE BETWEEN HAIRLINES AND BOX END	2 spaces

TOP  
PANEL 1

FOR LOT & EXPIRATION AREA.  
UNVARNISHED AND PRINTED  
IN BLACK OR DARK BLUE.

BASE  
PANEL 3

RELEASED BACK  
PANEL 2

000071768

DRUG FACTS TEXT DEFINED	TYPE SIZE
• DRUG FACTS TITLE	9 pt
• DRUG FACTS CONTINUED	8 pt
• HEADINGS	8 pt
• SUBHEADINGS/BODY TEXT	6 pt
• LEADING	6.5 pt
• # OF CHARACTERS PER INCH	<39
• BULLETS	5 pt
• SPACE BEFORE BULLET	2 ems
• BARLINES, HAIRLINES	1.5 pt, .5 pt
• SPACE BETWEEN HAIRLINES AND BOX END	2 spaces

TOP LAYER FACE

**Advil<sup>®</sup> LIQUI-GELS<sup>®</sup>**  
Solubilized Ibuprofen Capsules, 200 mg  
Pain Reliever/Fever Reducer (NSAID)

**minis**  
Advil  
200  
Liquid Filled  
Capsules

*Smaller Capsule Same Strength\**

000068425

Do Not Use if seal under bottle cap imprinted with "SEALED for YOUR PROTECTION" is broken or missing.

**FPO**

3 0573 1769 13 5

LIFT HERE for More Drug Facts

\*Compared to Advil<sup>®</sup> Liqui-Gels<sup>®</sup>

UNVARNISHED & PRINTED AREA FOR LOT & EXP. (LINES DO NOT PRINT)

TOP LAYER BACK

<p><b>Drug Facts (continued)</b></p> <p><b>Warnings</b> <b>Allergy alert:</b> Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:  <ul style="list-style-type: none"> <li>■ hives</li> <li>■ skin reddening</li> <li>■ facial swelling</li> <li>■ rash</li> <li>■ asthma (wheezing)</li> <li>■ blisters</li> <li>■ shock</li> </ul>                     If an allergic reaction occurs, stop use and seek medical help right away.  <b>Stomach bleeding warning:</b> This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you  <ul style="list-style-type: none"> <li>■ are age 60 or older</li> <li>■ have had stomach ulcers or bleeding problems</li> <li>■ take a blood thinning (anticoagulant) or steroid drug</li> <li>■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)</li> <li>■ have 3 or more alcoholic drinks every day while using this product</li> <li>■ take more or for a longer time than directed</li> </ul> <b>Heart attack and stroke warning:</b> NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.  <b>Do not use</b>  <ul style="list-style-type: none"> <li>■ if you have ever had an allergic reaction to any other pain reliever/fever reducer</li> <li>■ right before or after heart surgery</li> </ul> <b>Ask a doctor before use if</b>  <ul style="list-style-type: none"> <li>■ stomach bleeding warning applies to you</li> <li>■ you have problems or serious side effects from taking pain relievers or fever reducers</li> <li>■ you have a history of stomach problems, such as heartburn</li> <li>■ you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke</li> <li>■ you are taking a diuretic</li> </ul> </p>	<p><b>Drug Facts (continued)</b></p> <p><b>Ask a doctor or pharmacist before use if you are</b>  <ul style="list-style-type: none"> <li>■ under a doctor's care for any serious condition</li> <li>■ taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin</li> <li>■ taking any other drug</li> </ul> <b>When using this product</b>  <ul style="list-style-type: none"> <li>■ take with food or milk if stomach upset occurs</li> </ul> <b>Stop use and ask a doctor if</b>  <ul style="list-style-type: none"> <li>■ you experience any of the following signs of stomach bleeding:  <ul style="list-style-type: none"> <li>■ feel faint</li> <li>■ vomit blood</li> <li>■ have bloody or black stools</li> <li>■ have stomach pain that does not get better</li> </ul> </li> <li>■ you have symptoms of heart problems or stroke:  <ul style="list-style-type: none"> <li>■ chest pain</li> <li>■ trouble breathing</li> <li>■ weakness in one part or side of body</li> <li>■ slurred speech</li> <li>■ leg swelling</li> </ul> </li> <li>■ pain gets worse or lasts more than 10 days</li> <li>■ fever gets worse or lasts more than 3 days</li> <li>■ redness or swelling is present in the painful area</li> <li>■ any new symptoms appear</li> </ul> <b>If pregnant or breast-feeding,</b> ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.                 </p>
--	--

BASE LAYER

<p><b>Drug Facts (continued)</b></p> <p><b>Keep out of reach of children.</b> In case of overdose, get medical help or contact a Poison Control Center right away.</p> <p><b>Directions</b>  <ul style="list-style-type: none"> <li>■ do not take more than directed</li> <li>■ the smallest effective dose should be used</li> <li>■ adults and children 12 years and over: take 1 capsule every 4 to 6 hours while symptoms persist</li> <li>■ if pain or fever does not respond to 1 capsule, 2 capsules may be used</li> <li>■ do not exceed 6 capsules in 24 hours, unless directed by a doctor</li> <li>■ children under 12 years: ask a doctor</li> </ul> </p> <p><b>Other information</b>  <ul style="list-style-type: none"> <li>■ each capsule contains: potassium 20 mg</li> <li>■ read all warnings and directions before use.</li> <li>■ store at 20-25°C (68-77°F)</li> <li>■ avoid excessive heat above 40°C (104°F)</li> </ul> </p> <p><b>Inactive ingredients</b>                      FD&amp;C green no. 3, gelatin, medium-chain triglycerides, pharmaceutical ink, polyethylene glycol, potassium hydroxide, purified water, sorbitol sorbitan solution                 </p> <p><b>Questions or comments?</b> call toll free 1-800-88-ADVIL</p>	<h1>Advil<sup>®</sup></h1> <p>For most recent product information, visit <a href="http://www.Advil.com">www.Advil.com</a>                      For US Patent or Application status see <a href="http://www.productpats.com">www.productpats.com</a>                      Distributed by: GSK Consumer Healthcare, Warren, NJ 07059                      Trademarks are owned by or licensed to the GSK group of companies                      ©2021 GSK group of companies or its licensor                      LIQUI-GELS<sup>®</sup> is a trademark or registered trademark of Catalent Pharma Solutions.</p>
---	--

000068425

DRUG FACTS MODIFIED TEXT DEFINED	TYPE SIZE
• DRUG FACTS TITLE	9 pt
• DRUG FACTS CONTINUED	8 pt
• HEADINGS	8 pt
• SUBHEADINGS/BODY TEXT	6 pt
• LEADING	6.5 pt
• # OF CHARACTERS PER INCH	<39
• BULLETS	5 pt
• SPACE BEFORE BULLET	2 ems
• BARLINES, HAIRLINES	1.5 pt, .5 pt
• SPACE BETWEEN HAIRLINES AND BOX END	2 spaces

TOP LAYER FACE

000071418

Do Not Use if seal under bottle cap imprinted with "SEALED for YOUR PROTECTION" is broken or missing.

3/16" X 3/16" DATA MATRIX CODE (ECC 200), PRINTED IN BLACK OR DARK BLUE ON UNPRINTED WHITE BACKGROUND WITH A MINIMUM OF 1/8" QUIET ZONE ALL 4 SIDES.

UNVARNISHED & PRINTED AREA FOR LOT & EXP. (LINES DO NOT PRINT)

TOP LAYER BACK

<p><b>Drug Facts (continued)</b></p> <p><b>Warnings</b>  <b>Allergy alert:</b> Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:          ■ hives ■ facial swelling ■ asthma (wheezing) ■ shock          ■ skin reddening ■ rash ■ blisters          If an allergic reaction occurs, stop use and seek medical help right away.  <b>Stomach bleeding warning:</b> This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you          ■ are age 60 or older          ■ have had stomach ulcers or bleeding problems          ■ take a blood thinning (anticoagulant) or steroid drug          ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)          ■ have 3 or more alcoholic drinks every day while using this product          ■ take more or for a longer time than directed  <b>Heart attack and stroke warning:</b> NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.  <b>Do not use</b>          ■ if you have ever had an allergic reaction to any other pain reliever/fever reducer          ■ right before or after heart surgery  <b>Ask a doctor before use if</b>          ■ stomach bleeding warning applies to you          ■ you have problems or serious side effects from taking pain relievers or fever reducers          ■ you have a history of stomach problems, such as heartburn          ■ you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke          ■ you are taking a diuretic</p>	<p><b>Drug Facts (continued)</b></p> <p><b>Ask a doctor or pharmacist before use if you are</b>          ■ under a doctor's care for any serious condition          ■ taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin          ■ taking any other drug  <b>When using this product</b>          ■ take with food or milk if stomach upset occurs  <b>Stop use and ask a doctor if</b>          ■ you experience any of the following signs of stomach bleeding:          ■ feel faint          ■ vomit blood          ■ have bloody or black stools          ■ have stomach pain that does not get better          ■ you have symptoms of heart problems or stroke:          ■ chest pain          ■ trouble breathing          ■ weakness in one part or side of body          ■ slurred speech          ■ leg swelling          ■ pain gets worse or lasts more than 10 days          ■ fever gets worse or lasts more than 3 days          ■ redness or swelling is present in the painful area          ■ any new symptoms appear  <b>If pregnant or breast-feeding,</b> ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.</p>
--	--

NO COPY HINGED AREA

BASE LAYER

<p><b>Drug Facts (continued)</b></p> <p><b>Keep out of reach of children.</b> In case of overdose, get medical help or contact a Poison Control Center right away.</p> <p><b>Directions</b>          ■ do not take more than directed          ■ the smallest effective dose should be used          ■ adults and children 12 years and over: take 1 capsule every 4 to 6 hours while symptoms persist          ■ if pain or fever does not respond to 1 capsule, 2 capsules may be used          ■ do not exceed 6 capsules in 24 hours, unless directed by a doctor          ■ children under 12 years: ask a doctor</p> <p><b>Other information</b>          ■ each capsule contains: potassium 20 mg          ■ read all warnings and directions before use.          ■ store at 20-25°C (68-77°F)          ■ avoid excessive heat above 40°C (104°F)</p> <p><b>Inactive ingredients</b>          FD&amp;C green no. 3, gelatin, medium-chain triglycerides, pharmaceutical ink, polyethylene glycol, potassium hydroxide, purified water, sorbitol sorbitan solution</p> <p><b>Questions or comments?</b> call toll free 1-800-68-ADVIL</p>	<h1>Advil®</h1> <p>For most recent product information, visit <a href="http://www.Advil.com">www.Advil.com</a>          For US Patent or Application status see <a href="http://www.productpats.com">www.productpats.com</a></p> <p>Dist. by: GSK Consumer Healthcare, Warren, NJ 07059</p> <p>Trademarks owned or licensed by GSK          ©2021 GSK or licensor          LIQUI-GELS® is a trademark or registered trademark of Catalent Pharma Solutions.</p>	<p>NO COPY AREA</p>
--	---	---------------------

000071418

DRUG FACTS MODIFIED TEXT DEFINED	TYPE SIZE
• DRUG FACTS TITLE	9 pt
• DRUG FACTS CONTINUED	8 pt
• HEADINGS	8 pt
• SUBHEADINGS/BODY TEXT	6 pt
• LEADING	6.5 pt
• # OF CHARACTERS PER INCH	<39
• BULLETS	5 pt
• SPACE BEFORE BULLET	2 ems
• BARLINES, HAIRLINES	1.5 pt, .5 pt
• SPACE BETWEEN HAIRLINES AND BOX END	2 spaces



**READ AND KEEP CARTON FOR COMPLETE WARNINGS AND INFORMATION**

**Drug Facts**

**Active ingredient (in each capsule)**  
Solubilized ibuprofen equal to 200 mg ibuprofen (NSAID)\*. Pain reliever/fever reducer (present as the free acid and potassium salt)  
\*nonsteroidal anti-inflammatory drug

**Purpose**

**Uses**  
temporarily relieves minor aches and pains due to:  
headache, toothache, backache, menstrual cramps, the common cold, muscular aches, minor pain of arthritis, temporarily reduces fever

**Warnings**

**Allergy alert:** Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:  
hives, facial swelling, asthma (wheezing), shock, skin reddening, rash, blisters  
If an allergic reaction occurs, stop use and seek medical help right away.  
**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:  
are age 60 or older, have had stomach ulcers or bleeding problems, take a blood thinning (anticoagulant) or steroid drug, take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others), have 3 or more alcoholic drinks every day while using this product, take more or for a longer time than directed.  
**Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

**Do not use**  
if you have ever had an allergic reaction to any other pain reliever/fever reducer, right before or after heart surgery

**Ask a doctor before use if**  
stomach bleeding warning applies to you, you have problems or serious side effects from taking pain relievers or fever reducers, you have a history of stomach problems, such as heartburn

**Drug Facts (continued)**

you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke, you are taking a diuretic

**Ask a doctor or pharmacist before use if you are**  
under a doctor's care for any serious condition, taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin, taking any other drug

**When using this product** take with food or milk if stomach upset occurs

**Stop use and ask a doctor if**  
you experience any of the following signs of stomach bleeding:  
feel faint, vomit blood, have bloody or black stools, have stomach pain that does not get better, you have symptoms of heart problems or stroke:  
chest pain, trouble breathing, weakness in one part or side of body, slurred speech, leg swelling, pain gets worse or lasts more than 10 days, fever gets worse or lasts more than 3 days, redness or swelling is present in the painful area, any new symptoms appear

**If pregnant or breast-feeding**, ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.  
**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions** do not take more than directed, the smallest effective dose should be used, adults and children 12 years and over: take 1 capsule every 4 to 6 hours while symptoms persist, if pain or fever does not respond to 1 capsule, 2 capsules may be used, do not exceed 6 capsules in 24 hours, unless directed by a doctor, children under 12 years: ask a doctor

**Other information** each capsule contains: potassium 20 mg, read all warnings and directions before use. Keep carton, store at 20-25°C (68-77°F), avoid excessive heat above 40°C (104°F)

**Inactive ingredients** FD&C green no. 3, gelatin, medium-chain triglycerides, pharmaceutical ink, polyethylene glycol, potassium hydroxide, purified water, sorbitol sorbitan solution

**Questions or comments?** call toll free 1-800-88-ADVIL

Do Not Use if seal under bottle cap imprinted with "SEALED for YOUR PROTECTION" is broken or missing.

For most recent product information, visit [www.Advil.com](http://www.Advil.com)  
Distributed by: GSK Consumer Healthcare, Warren, NJ 07059  
©2021 GSK group of companies or its licensor  
Trademarks are owned by or licensed to the GSK group of companies  
For US Patent or Application status see [www.productpats.com](http://www.productpats.com)  
LIQUI-GELS® is a trademark or registered trademark of Catalent Pharma Solutions.



000071633

PRINTED AREA FOR DATE & LOT  
LASER CODING (VARNISHED)

000071633

DRUG FACTS TEXT DEFINED	TYPE SIZE
• DRUG FACTS TITLE	9 pt
• DRUG FACTS CONTINUED	8 pt
• HEADINGS	8 pt
• SUBHEADINGS/BODY TEXT	6 pt
• LEADING	6.5 pt
• # OF CHARACTERS PER INCH	<39
• BULLETS	5 pt
• SPACE BEFORE BULLET	2 ems
• BARLINES, HAIRLINES	1.5 pt, .5 pt
• SPACE BETWEEN HAIRLINES AND BOX END	2 spaces





**READ AND KEEP CARTON FOR COMPLETE WARNINGS AND INFORMATION**

**Drug Facts**  
*(in each brown oval capsule)*

**Purpose**  
Solubilized ibuprofen equal to 200 mg ibuprofen (NSAID) (present as the free acid and potassium salt) nonsteroidal anti-inflammatory drug

**Use**  
■ treats migraine

**Warnings**  
**Allergy alert:** Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:  
■ facial swelling  
■ hives  
■ asthma (wheezing)  
■ shock  
■ skin redness  
■ rash  
■ blisters

If an allergic reaction occurs, stop use and seek medical help right away.

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you  
■ are age 60 or older  
■ have had stomach ulcers or bleeding problems  
■ take a blood thinning (anticoagulant) or steroid drug  
■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)  
■ have 3 or more alcoholic drinks every day while using this product.  
■ take more or for a longer time than directed

**Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.  
**Medication overdose headache warning:** Headaches may worsen if this product is used for 10 or more days per month.

**Do not use**  
■ if you have ever had an allergic reaction to any other pain reliever/fever reducer  
■ right before or after heart surgery

**Ask a doctor before use if**  
■ you have never had migraines diagnosed by a health professional  
■ you have a headache that is different from your usual migraines  
■ you have the worst headache of your life  
■ you have fever and stiff neck  
■ you have headaches beginning after or caused by head injury, exertion, coughing or bending  
■ you have experienced your first headache after the age of 50  
■ you have daily headaches  
■ you have a migraine so severe as to require bed rest  
■ you have stomach pain, vomiting, or diarrhea  
■ you have had stomach bleeding, vomiting, or black stools from taking pain relievers or fever reducers  
■ you have a history of stomach problems, such as heartburn  
■ you have high blood pressure, heart disease, new or worse heartburn, kidney disease, asthma, or had a stroke  
■ you are taking a diuretic

**Ask a doctor or pharmacist before use if you are**  
■ under a doctor's care for any serious condition  
■ taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin  
■ taking any other drug

**When using this product:**  
■ take with food or milk if stomach upset occurs

**Stop use and ask a doctor if**  
■ you experience any of the following signs of stomach bleeding:  
■ feel faint  
■ vomit blood  
■ have bloody or black stools  
■ have stomach pain that does not get better  
■ you have symptoms of heart problems or stroke:  
■ chest pain  
■ weakness in one part or side of body  
■ slurred speech  
■ leg swelling

■ minimize headache pain is not relieved or gets worse after the first dose  
■ any new symptoms appear

**If pregnant or breast-feeding, ask a health professional before use.** It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**  
■ do not take more than directed  
■ the smallest effective dose should be used

adults  
■ take 2 capsules with a glass of water  
■ if symptoms persist or worsen, ask your doctor  
■ do not take more than 2 capsules in 24 hours, unless directed by a doctor

under 18 years of age  
■ ask a doctor

PRINTED AREA FOR DATE & LOT  
LASER CODING (VARNISHED)

UNPRINTED AREA  
FOR DATE & LOT  
(UNVARNISHED)

For most recent product information, visit [www.Advii.com](http://www.Advii.com)  
Distributed by: GSKCH, Warren, NJ 07059  
©2021 GSK or licensor  
Trademarks owned or licensed by GSK  
000071635

**FPO**

3 0 5 7 3 0 1 6 8 3 0 4

**Advil**  
**MIGRAINE**  
Solubilized Ibuprofen Capsules, 200 mg  
Pain Reliever (NSAID)  
40 Liquid Filled Capsules

**Drug Facts (continued)**

**Other information**  
■ each capsule contains potassium 20 mg  
■ read all directions and warnings before use.  
■ avoid excessive heat above 40°C (104°F)  
■ store at 20-25°C (68-77°F)  
Keep carton, water, sorbitol sorbitan solution  
■ Inactive ingredients: D&C yellow no. 10, FD&C green no. 3, FD&C red no. 40, gelatin, pharmaceutical ink, polyethylene glycol, potassium hydroxide, purified water, sorbitol sorbitan solution

**Questions or comments?**  
call toll free 1-800-88-ADVIL

Do Not Use if seal under bottle cap imprinted with "SEALED for YOUR PROTECTION" is broken or missing.

000071635

DRUG FACTS TEXT DEFINED	TYPE SIZE
• DRUG FACTS TITLE	9 pt
• DRUG FACTS CONTINUED	8 pt
• HEADINGS	8 pt
• SUBHEADINGS/BODY TEXT	6 pt
• LEADING	6.5 pt
• # OF CHARACTERS PER INCH	<39
• BULLETS	5 pt
• SPACE BEFORE BULLET	2 ems
• BARLINES, HAIRLINES	1.5 pt, .5 pt
• SPACE BETWEEN HAIRLINES AND BOX END	2 spaces

# Advil

## MIGRAINE

Solubilized Ibuprofen Capsules, 200 mg  
Pain Reliever (NSAID)

80  
Liquid Filled  
Capsules

READ AND KEEP CARTON FOR COMPLETE WARNINGS AND INFORMATION

**Drug Facts**

**Active ingredient**  
(in each brown oval capsule) Solubilized ibuprofen equal to 200 mg ibuprofen (NSAID)\*

**Use**

- treats migraine

**Warnings**

**Allergy alert:** Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- asthma (wheezing)
- skin reddening
- facial swelling
- rash
- blisters
- stomach bleeding
- shock

If an allergic reaction occurs, stop use and seek medical help right away.

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

**Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

**Medication overdose headache warning:** Headaches may worsen if this product is used for 10 or more days per month.

**Purpose**  
Pain reliever

80  
Liquid Filled  
Capsules

Advil

Do Not Use if seal under bottle cap imprinted with "SEALED for YOUR PROTECTION" is broken or missing.  
For most recent product information, visit [www.Advil.com](http://www.Advil.com)  
©2021 GSK Group of companies or its licensor  
Distributed by: GSK Consumer Healthcare, Warren, NJ 07089  
Trademarks are owned by or licensed to the GSK group of companies

**Drug Facts (continued)**

**Do not use**

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

**Ask a doctor before use if**

- you have never had migraines diagnosed by a health professional
- you have a headache that is different from your usual migraines
- you have the worst headache of your life
- you have fever and stiff neck
- you have headaches beginning after or caused by head injury, exertion, coughing or sneezing
- you have experienced your first headache after the age of 50
- you have daily headaches
- you have a migraine so severe as to require bed rest
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic

**Ask a doctor or pharmacist before use if you are**

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- taking any other drug

**When using this product**

- take with food or milk if stomach upset occurs

**Stop use and ask a doctor if**

- you experience any of the following signs of stomach bleeding:
  - feel faint
  - vomit blood
  - have bloody or black stools
- have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
  - chest pain
  - trouble breathing
  - weakness in one part or side of body
  - slurred speech
  - leg swelling
- migraine headache pain is not relieved or gets worse after the first dose
- any new symptoms appear

**Drug Facts (continued)**

**If pregnant or breast-feeding, ask a health professional before use.** It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**

- do not take more than directed
- the smallest effective dose should be used

adults

- take 2 capsules with a glass of water
- if symptoms persist or worsen, ask your doctor
- do not take more than 2 capsules in 24 hours, unless directed by a doctor
- ask a doctor

under 18 years of age

**Other information**

- each capsule contains potassium 20 mg
- read all directions and warnings before use. Keep carton.
- store at 20-25°C (68°-77°F)
- avoid excessive heat above 40°C (104°F)

**Inactive ingredients**

D&C yellow no. 10, FD&C green no. 3, FD&C red no. 40, gelatin, pharmaceutical ink, polyethylene glycol, potassium hydroxide, purified water, sorbitol sorbitan solution

**Questions or comments?**  
call toll free 1-800-88-ADVIL

000071157

FPO

3 0573 0168 40 3

UNPRINTED AREA FOR COMPLETE WARNINGS AND INFORMATION

UNPRINTED AREA FOR DATE & LOT CODING (UNVARNISHED)

PRINTED AREA FOR DATE & LOT LASER CODING (VARNISHED)

000071157

DRUG FACTS TEXT DEFINED	TYPE SIZE
• DRUG FACTS TITLE	9 pt
• DRUG FACTS CONTINUED	8 pt
• HEADINGS	8 pt
• SUBHEADINGS/BODY TEXT	6 pt
• LEADING	6.5 pt
• # OF CHARACTERS PER INCH	<39
• BULLETS	5 pt
• SPACE BEFORE BULLET	2 ems
• BARLINES, HAIRLINES	1.5 pt, .5 pt
• SPACE BETWEEN HAIRLINES AND BOX END	2 spaces

-----  
**This is a representation of an electronic record that was signed electronically. Following this are manifestations of any and all electronic signatures for this electronic record.**  
-----

/s/  
-----

VALERIE S PRATT  
01/25/2022 02:41:10 PM