



PRINTED AREA FOR DATE & LOT
LASER CODING (VARNISHED)

214561

| DRUG FACTS TEXT DEFINED | TYPE SIZE |
|---------------------------------------|---------------|
| • DRUG FACTS TITLE | 9 pt |
| • DRUG FACTS CONTINUED | 8 pt |
| • HEADINGS | 8 pt |
| • SUBHEADINGS/BODY TEXT | 6 pt |
| • LEADING | 6.5 pt |
| • # OF CHARACTERS PER INCH | <39 |
| • BULLETS | 5 pt |
| • SPACE BEFORE BULLET | 2 ems |
| • BARLINES, HAIRLINES | 1.5 pt, .5 pt |
| • SPACE BETWEEN HAIRLINES AND BOX END | 2 spaces |

READ AND KEEP CARTON FOR COMPLETE WARNINGS AND INFORMATION

Drug Facts

Active ingredient (in each capsule) Solubilized ibuprofen (present as the free acid and potassium salt) *nonsteroidal anti-inflammatory drug

Purpose Pain reliever/Fever reducer

Uses

- temporarily relieves minor aches and pains due to:
 - headache
 - toothache
 - backache
 - menstrual cramps
 - the common cold
 - muscular aches
 - minor pain of arthritis
- temporarily reduces fever

Warnings

- Allergy alert:** Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:
- hives
 - facial swelling
 - asthma (wheezing)
 - shock
 - skin reddening
 - rash
 - blisters
- If an allergic reaction occurs, stop use and seek medical help right away.
- Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:
- are age 60 or older
 - have had stomach ulcers or bleeding problems
 - take a blood thinning (anticoagulant) or steroid drug
 - take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
 - have 3 or more alcoholic drinks every day while using this product
 - take more or for a longer time than directed
- Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
 - right before or after heart surgery
- Ask a doctor before use if**
- stomach bleeding warning applies to you
 - you have problems or serious side effects from taking pain relievers or fever reducers
 - you have a history of stomach problems, such as heartburn
 - you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
 - you are taking a diuretic

Drug Facts (continued)

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
- have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
 - chest pain
 - trouble breathing
 - weakness in one part or side of body
 - slurred speech
 - leg swelling
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not take more than directed
- the smallest effective dose should be used
- adults and children 12 years and over: take 1 capsule every 4 to 6 hours while symptoms persist
- if pain or fever does not respond to 1 capsule, 2 capsules may be used
- do not exceed 6 capsules in 24 hours, unless directed by a doctor
- children under 12 years: ask a doctor

Other information

- each capsule contains: potassium 20 mg
- read all warnings and directions before use. Keep carton.
- store at 20-25°C (68-77°F)
- avoid excessive heat above 40°C (104°F)

Inactive ingredients: FD&C green no. 3, gelatin, lecithin (soybean), medium-chain triglycerides, pharmaceutical ink, polyethylene glycol, potassium hydroxide, purified water, sorbitol sorbitan solution

Questions or comments? call toll free 1-800-88-ADVIL

Advil



VALUE PACK
2x80 Count

RELIEF AT HOME & ON-THE-MOVE

VALUE PACK
2x80 Count

Advil
LIQUI-GELS

Solubilized Ibuprofen
Capsules, 200 mg Pain Reliever/
Fever Reducer (NSAID)
2x80 Count Bottles Inside

Actual Size
160
Liquid Filled
Capsules

VALUE PACK
2x80 Count

Advil
LIQUI-GELS

Solubilized Ibuprofen
Capsules, 200 mg Pain Reliever/
Fever Reducer (NSAID)
2x80 Count
Bottles Inside

Actual Size
160
Liquid Filled
Capsules

VALUE PACK
2x80 Count

Advil
LIQUI-GELS

Solubilized Ibuprofen
Capsules, 200 mg Pain Reliever/
Fever Reducer (NSAID)
2x80 Count
Bottles Inside

Actual Size

Do Not Use if seal under bottle cap imprinted with "SEALED for YOUR PROTECTION" is broken or missing.

Distributed by: Haleon, Warren, NJ 07059
For most recent product information, visit www.Advil.com
Trademarks are owned by or licensed to the Haleon group of companies.
©2024 Haleon group of companies or its licensor.
LIQUI-GELS® is a trademark or registered trademark of Catalent Pharms Solutions.





PRINTED AREA FOR DATE & LOT
LASER CODING (VARNISHED)

214562

| DRUG FACTS TEXT DEFINED | TYPE SIZE |
|---------------------------------------|---------------|
| • DRUG FACTS TITLE | 9 pt |
| • DRUG FACTS CONTINUED | 8 pt |
| • HEADINGS | 8 pt |
| • SUBHEADINGS/BODY TEXT | 6 pt |
| • LEADING | 6.5 pt |
| • # OF CHARACTERS PER INCH | <39 |
| • BULLETS | 5 pt |
| • SPACE BEFORE BULLET | 2 ems |
| • BARLINES, HAIRLINES | 1.5 pt, .5 pt |
| • SPACE BETWEEN HAIRLINES AND BOX END | 2 spaces |

This is a representation of an electronic record that was signed electronically. Following this are manifestations of any and all electronic signatures for this electronic record.

/s/

NUSHIN F TODD
01/08/2025 09:10:41 AM