

1. The first step is to identify the problem. This involves understanding the current situation and what needs to be improved.

2. The second step is to set clear goals. These should be specific, measurable, achievable, relevant, and time-bound (SMART).

3. The third step is to develop a plan. This involves identifying the resources needed and the steps to be taken to achieve the goals.

4. The fourth step is to implement the plan. This involves putting the plan into action and monitoring progress.

5. The fifth step is to evaluate the results. This involves comparing the actual results with the goals and identifying areas for improvement.

6. The sixth step is to make adjustments. This involves making changes to the plan or implementation based on the evaluation.

7. The seventh step is to communicate the results. This involves sharing the findings with stakeholders and providing feedback.

8. The eighth step is to document the process. This involves keeping records of the plan, implementation, and results for future reference.

9. The ninth step is to review the process. This involves reflecting on the overall experience and identifying lessons learned.

10. The tenth step is to celebrate success. This involves recognizing the achievements and rewarding the team for their efforts.

NDA 20-448/S-004

Page 4

20-ct bottle label:

Active ingredient (in each tablet) Loperamide HCl 2 mg.....Anti-diarrheal Use controls symptoms of diarrhea, including Travelers' Diarrhea Warnings Allergy alert: Do not use if you have ever	Purpose NDC 50580-397-20 Imodium® A-D Loperamide HCl/ Anti-Diarrheal	drowsiness or dizziness may occur. Be careful when driving or operating machinery. Stop use and ask a doctor if: • symptoms get worse • diarrhea lasts for more than 2 days • you get abdominal swelling or bulging. These may be signs of a serious condition. If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222) Directions • drink plenty of clear fluids to help prevent dehydration caused by diarrhea • take only on an empty stomach (1 hour before or 2 hours after a meal) • find right dose on chart. If possible, use weight to dose; otherwise, use age.								
LOT EXP.	20 Chewable Cool Mint Flavored Tablets EZ CHEWS	<table border="1"> <tr> <td>adults and children 12 years and over</td> <td>chew 2 tablets after the first loose stool; but no more than 4 tablets in 24 hours</td> </tr> <tr> <td>children 9-11 years (60-95 lbs)</td> <td>chew 1 tablet after the first loose stool; chew 1/2 tablet after each subsequent loose stool; but no more than 3 tablets in 24 hours</td> </tr> <tr> <td>children 6-8 years (48-59 lbs)</td> <td>chew 1 tablet after the first loose stool; chew 1/2 tablet after each subsequent loose stool; but no more than 2 tablets in 24 hours</td> </tr> <tr> <td>children under 6 years (up to 47 lbs)</td> <td>ask a doctor</td> </tr> </table>	adults and children 12 years and over	chew 2 tablets after the first loose stool; but no more than 4 tablets in 24 hours	children 9-11 years (60-95 lbs)	chew 1 tablet after the first loose stool; chew 1/2 tablet after each subsequent loose stool; but no more than 3 tablets in 24 hours	children 6-8 years (48-59 lbs)	chew 1 tablet after the first loose stool; chew 1/2 tablet after each subsequent loose stool; but no more than 2 tablets in 24 hours	children under 6 years (up to 47 lbs)	ask a doctor
adults and children 12 years and over	chew 2 tablets after the first loose stool; but no more than 4 tablets in 24 hours									
children 9-11 years (60-95 lbs)	chew 1 tablet after the first loose stool; chew 1/2 tablet after each subsequent loose stool; but no more than 3 tablets in 24 hours									
children 6-8 years (48-59 lbs)	chew 1 tablet after the first loose stool; chew 1/2 tablet after each subsequent loose stool; but no more than 2 tablets in 24 hours									
children under 6 years (up to 47 lbs)	ask a doctor									
XXXXXXX	had a rash or other allergic reaction to loperamide HCl. Do not use if you have bloody or black stool. Ask a doctor before use if you have: • fever • mucus in the stool • a history of liver disease. Ask a doctor or pharmacist before use if you are taking antibiotics When using this product • tiredness,	Other information • store between 20-25°C (68-77°F) • do not use if printed foil seal under bottle cap is open or torn Questions or comments? call 1-877-895-3665 (English) or 1-888-466-8746 (Spanish) DISTRIBUTED BY: MCNEIL-PPC, INC., FORT WASHINGTON, PA 19034 USA © MCN-PPC, Inc. '06								