


Children's Nasacort Allergy 24HR 60 Sprays Partial Cover Clamshell - Outside Layout

11.05.14

NAS_Childrens_Allergy_24HR_60sprays



Children's Nasacort Allergy 24HR 60 Sprays Partial Cover Clamshell - Drug Facts Only

11.05.14

NAS_Childrens_Allergy_24HR_60sprays

(b) (4)

(b) (4)

Drug Facts		Drug Facts (continued)	
Active ingredient (in each spray) Triamcinolone acetonide (glucocorticoid) 55 mcg Nasal allergy symptom reliever		Directions Read insert (inside package) on how to: ■ get a new bottle ready (primed) before first use ■ prime bottle again if not used for more than 2 weeks ■ use the spray ■ clean the spray nozzle	
Uses temporarily relieves these symptoms of hay fever or other upper respiratory allergies: ■ nasal congestion ■ runny nose ■ sneezing ■ itchy nose		ADULTS AND CHILDREN 12 YEARS OF AGE AND OLDER adults and children 12 years of age and older ■ once daily, spray 2 times into each nostril while sniffing gently ■ once your allergy symptoms improve, reduce to 1 spray in each nostril per day	
Warnings Do not use ■ in children under 2 years of age ■ if you have ever had an allergic reaction to any of the ingredients Ask a doctor before use if you ■ have had recent nose ulcers or nose surgery ■ have had a nose injury that has not healed ■ are using a steroid medicine for asthma, allergies or skin rash ■ have an eye infection ■ have or had glaucoma or cataracts		CHILDREN 2 TO UNDER 12 YEARS OF AGE ■ the growth rate of some children may be slower while using this product. Talk to your child's doctor if your child needs to use the spray for longer than two months a year. children 6 to under 12 years of age ■ an adult should supervise use ■ once daily, spray 1 time into each nostril while sniffing gently ■ if allergy symptoms do not improve, increase to 2 sprays in each nostril per day. Once allergy symptoms improve, reduce to 1 spray in each nostril per day.	
When using this product ■ the growth rate of some children may be slower ■ some symptoms may get better on the first day of treatment. It may take up to one week of daily use to feel the most symptom relief. ■ do not share this bottle with anyone else as this may spread germs ■ remember to tell your doctor about all the medicines you take, including this one		children 2 to under 6 years of age ■ an adult should supervise use ■ once daily, spray 1 time into each nostril while sniffing gently	
Stop use and ask a doctor if ■ you have, or come into contact with someone who has, chickenpox, measles or tuberculosis ■ you have or develop symptoms of an infection such as a persistent fever ■ you have any change in vision ■ you have severe or frequent nosebleeds		children under 2 years of age ■ do not use	
If pregnant or breast-feeding , ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.		Other information ■ do not use if sealed package is torn or opened ■ keep package and insert. They contain important information. ■ store between 20°-25°C (68°-77°F)	

(b) (4)


Drug Facts (continued) Inactive ingredients benzalkonium chloride, carboxymethylcellulose sodium, dextrose, edetate disodium, hydrochloric acid or sodium hydroxide (for pH adjustment), microcrystalline cellulose, polysorbate 80, purified water Questions or comments? call toll-free 1-800-633-1610

(b) (4)

Children's Nasacort Allergy 24HR 60 Sprays Partial Cover Clamshell - Inside Layout

11.05.14

NAS_Childrens_Allergy_24HR_60sprays



Children's Nasacort Allergy 24HR 60 LBL


11.05.14

NAS_Childrens_Allergy_24HR_60_LBL


PAGE 1 (Front Page)




PAGE 2 (Inside Front Page)




PAGE 3




PAGE 4




PAGE 5




PAGE 6



PAGE 7



PAGE 8



BASE LABEL

