

# Nasacort Allergy 24HR 30 LBL


08.23.16

NAS\_Allergy\_24HR\_30\_LBL


## PAGE 1 (Front Page)




## PAGE 2 (Inside Front Page)




## PAGE 3




## PAGE 4




## PAGE 5





## PAGE 6



## PAGE 7



## PAGE 8



## BASE LABEL

# Nasacort Allergy 24HR 60 LBL


08.23.16

NAS\_Allergy\_24HR\_60\_LBL


## PAGE 1 (Front Page)




## PAGE 2 (Inside Front Page)




## PAGE 3




## PAGE 4




## PAGE 5




## PAGE 6




## PAGE 7



## PAGE 8



## BASE LABEL



# Nasacort Allergy 24HR 120 LBL


08.23.16

NAS\_Allergy\_24HR\_120\_LBL\_r3\_v2


## PAGE 1 (Front Page)




## PAGE 2 (Inside Front Page)




## PAGE 3




## PAGE 4




## PAGE 5





## PAGE 6



## PAGE 7



## PAGE 8



## BASE LABEL

# Nasacort Allergy 24HR 60 Sprays

08.23.16

NAS\_Allergy\_60 Spray\_for IRC

|   |   |                                      |   |
|---|---|--------------------------------------|---|
| <p><b>Drug Facts</b></p>  | <p><b>Drug Facts (continued)</b></p>  | <p><b>Nasacort® Allergy 24HR</b></p> | <p>NDC XXXXX-XXXX-X</p> <p>Multi-Symptom Nasal Allergy Spray</p> <h2>Nasacort® Allergy 24HR</h2> <p>triamcinolone acetonide nasal spray 55 mcg per spray<br/>allergy symptom reliever (glucocorticoid)*</p> <p><b>24HR</b><br/>Relief of:</p> <ul style="list-style-type: none"> <li>Nasal Congestion</li> <li>Sneezing</li> <li>Runny Nose</li> <li>Itchy Nose</li> </ul> <p>0.37 fl. oz. (10.8 ml)</p> <p><b>60</b><br/>SPRAYS</p> <p><small>* Triamcinolone acetonide is a steroid medicine known as a glucocorticoid.</small></p> |
| <p><b>Active ingredient (in each spray)</b> Purpose</p> <p>Triamcinolone acetonide<br/>55 mcg (glucocorticoid) .....Allergy symptom reliever</p> <p><b>Uses</b><br/>temporarily relieves these symptoms of hay fever or other upper respiratory allergies:<br/>                 ■ nasal congestion ■ runny nose ■ sneezing ■ itchy nose</p> <p><b>Warnings</b><br/><b>Do not use</b><br/>                 ■ in children under 2 years of age<br/>                 ■ if you have ever had an allergic reaction to any of the ingredients</p> <p><b>Ask a doctor before use if you</b><br/>                 ■ have had recent nose ulcers or nose surgery<br/>                 ■ have had a nose injury that has not healed<br/>                 ■ are using a steroid medicine for asthma, allergies or skin rash<br/>                 ■ have an eye infection<br/>                 ■ have or had glaucoma or cataracts</p> <p><b>When using this product</b><br/>                 ■ the growth rate of some children may be slower<br/>                 ■ some symptoms may get better on the first day of treatment. It may take up to one week of daily use to feel the most symptom relief.<br/>                 ■ do not share this bottle with anyone else as this may spread germs<br/>                 ■ remember to tell your doctor about all the medicines you take, including this one</p> <p><b>Stop use and ask a doctor if</b><br/>                 ■ you have, or come into contact with someone who has, chickenpox, measles or tuberculosis<br/>                 ■ you have or develop symptoms of an infection such as a persistent fever<br/>                 ■ you have any change in vision<br/>                 ■ you have severe or frequent nosebleeds</p> <p><b>If pregnant or breast-feeding</b>, ask a health professional before use.<br/> <b>Keep out of reach of children.</b> In case of overdose, get medical help or contact a Poison Control Center right away.</p> | <p><b>Directions</b><br/>Read insert (inside package) on how to:</p> <ul style="list-style-type: none"> <li>get a new bottle ready (primed) before first use</li> <li>prime bottle again if not used for more than 2 weeks</li> <li>use the spray</li> <li>clean the spray nozzle</li> </ul> <p><b>ADULTS AND CHILDREN 12 YEARS OF AGE AND OLDER</b></p> <p><b>adults and children 12 years of age and older</b><br/>                 ■ once daily, spray 2 times into each nostril while sniffing gently<br/>                 ■ once your allergy symptoms improve, reduce to 1 spray in each nostril per day</p> <p><b>CHILDREN 2 TO UNDER 12 YEARS OF AGE</b></p> <p><b>children 6 to under 12 years of age</b><br/>                 ■ the growth rate of some children may be slower while using this product. Talk to your child's doctor if your child needs to use the spray for longer than two months a year.<br/>                 ■ an adult should supervise use<br/>                 ■ once daily, spray 1 time into each nostril while sniffing gently<br/>                 ■ if allergy symptoms do not improve, increase to 2 sprays in each nostril per day. Once allergy symptoms improve, reduce to 1 spray in each nostril per day.</p> <p><b>children 2 to under 6 years of age</b><br/>                 ■ an adult should supervise use<br/>                 ■ once daily, spray 1 time into each nostril while sniffing gently</p> <p><b>children under 2 years of age</b><br/>                 ■ do not use</p> <p><b>do not use more than directed</b><br/>                 ■ if you forget a dose, do not double the next dose<br/>                 ■ do not spray into eyes or mouth<br/>                 ■ if allergy symptoms do not improve after one week, stop using and talk to a doctor<br/>                 ■ do not use for the common cold<br/>                 ■ shake well before each use</p> <p><b>Other information</b><br/>                 ■ do not use if sealed package is torn or opened<br/>                 ■ keep package and insert. They contain important information.<br/>                 ■ store between 20°-25°C (68°-77°F)</p> |                                      |   |
| <p>Dist. by: Chatten, Inc.,<br/>a Sanofi Company<br/>Chattanooga, TN 37409-0219<br/>© 2016 Origin France</p>  | <p>UPC FPO</p>  | <p>LOT<br/>EXP<br/>00XXXXXX</p>      |   |

# Nasacort Allergy 24HR 60 Sprays

08.23.16

NAS\_Allergy\_60 Spray\_for IRC

Prescription strength allergy relief.  
Without a prescription.

# Nasacort<sup>®</sup> Allergy 24HR

24 hour symptom relief  
from indoor & outdoor allergies

Read insert (inside package)  
for complete instructions  
on how to:

- ▶ Get a new bottle ready (primed) before first use
- ▶ Prime bottle again if not used for more than 2 weeks
- ▶ Use the spray
- ▶ Clean the spray nozzle

**BEFORE FIRST USE,  
A NEW BOTTLE  
MUST BE PRIMED**

This label may not be the latest approved by FDA.  
For current labeling information, please visit <https://www.fda.gov/drugsatfda>

# Nasacort Allergy 24HR 60 Sprays with Peel-Away IRC

08.23.16

NAS\_Allergy\_60 Sprays\_IRC

**SAVE \$2.00 NOW!** PEEL HERE ►

NDC XXXXX-XXXX-X

Multi-Symptom  
Nasal Allergy Spray

**Nasacort®  
Allergy 24HR**

MANUFACTURER'S COUPON | EXPIRATION DATE XX/XX/XX

**SAVE \$2.00**  
When You Buy Nasacort® Allergy 24HR

**CONSUMER:** There is a limit of one coupon per purchase, and the coupon can't be combined with other offers. Good only in the U.S. and overseas commissaries on the purchase of the specified product/size. Void where prohibited or if altered, copied, transferred, auctioned, or sold. You must pay sales tax. Cash value: 1/100 of 1¢. **RETAILER:** We will reimburse you for the face value of this coupon plus 8¢ if redeemed in compliance with our Coupon Reimbursement Policy (available at address below). Invoices proving purchase of sufficient stock to cover coupons presented for redemption must be provided upon request. Failure to follow the policy may constitute fraud and may void all coupons submitted. Mail to: Chatterm, Inc., PO Box 880445, El Paso, TX 88588-0445

UPC FPO

©2016 Chatterm, Inc. XXXXXXXXXXXX

# Nasacort Allergy 24HR 60 Sprays with Peel-Away IRC

08.23.16

NAS\_Allergy\_60 Sprays\_IRC



# Nasacort Allergy 24HR 120 Sprays Bonus Pack

08.22.16

NAS\_Allergy\_120plus30\_Bonus\_r3

|   |  |  |  |
|---|--|--|--|
| <p><b>Drug Facts</b></p> <p><b>Active ingredient (in each spray)</b> Purpose<br/>                 Triamcinolone acetonide<br/>                 55 mcg (glucocorticoid) .....Allergy symptom reliever</p> <p><b>Uses</b><br/>                 temporarily relieves these symptoms of hay fever or other upper respiratory allergies:<br/>                 ■ nasal congestion ■ runny nose ■ sneezing ■ itchy nose</p> <p><b>Warnings</b><br/> <b>Do not use</b><br/>                 ■ in children under 2 years of age<br/>                 ■ if you have ever had an allergic reaction to any of the ingredients</p> <p><b>Ask a doctor before use if you</b><br/>                 ■ have had recent nose ulcers or nose surgery<br/>                 ■ have had a nose injury that has not healed<br/>                 ■ are using a steroid medicine for asthma, allergies or skin rash<br/>                 ■ have an eye infection<br/>                 ■ have or had glaucoma or cataracts</p> <p><b>When using this product</b><br/>                 ■ the growth rate of some children may be slower<br/>                 ■ some symptoms may get better on the first day of treatment. It may take up to one week of daily use to feel the most symptom relief.<br/>                 ■ do not share this bottle with anyone else as this may spread germs<br/>                 ■ remember to tell your doctor about all the medicines you take, including this one</p> <p><b>Stop use and ask a doctor if</b><br/>                 ■ you have, or come into contact with someone who has, chickenpox, measles or tuberculosis<br/>                 ■ you have or develop symptoms of an infection such as a persistent fever<br/>                 ■ you have any change in vision<br/>                 ■ you have severe or frequent nosebleeds</p> <p><b>If pregnant or breast-feeding</b>, ask a health professional before use.<br/> <b>Keep out of reach of children.</b> In case of overdose, get medical help or contact a Poison Control Center right away.</p> |  | <p><b>Drug Facts (continued)</b></p> <p><b>Directions</b><br/>                 Read insert (inside package) on how to:<br/>                 ■ get a new bottle ready (primed) before first use<br/>                 ■ prime bottle again if not used for more than 2 weeks<br/>                 ■ use the spray<br/>                 ■ clean the spray nozzle</p> <p><b>ADULTS AND CHILDREN 12 YEARS OF AGE AND OLDER</b><br/>                 adults and children 12 years of age and older<br/>                 ■ once daily, spray 2 times into each nostril while sniffing gently<br/>                 ■ once your allergy symptoms improve, reduce to 1 spray in each nostril per day</p> <p><b>CHILDREN 2 TO UNDER 12 YEARS OF AGE</b><br/>                 ■ the growth rate of some children may be slower while using this product. Talk to your child's doctor if your child needs to use the spray for longer than two months a year.</p> <p>children 6 to under 12 years of age<br/>                 ■ an adult should supervise use<br/>                 ■ once daily, spray 1 time into each nostril while sniffing gently<br/>                 ■ if allergy symptoms do not improve, increase to 2 sprays in each nostril per day. Once allergy symptoms improve, reduce to 1 spray in each nostril per day.</p> <p>children 2 to under 6 years of age<br/>                 ■ an adult should supervise use<br/>                 ■ once daily, spray 1 time into each nostril while sniffing gently</p> <p>children under 2 years of age<br/>                 ■ do not use</p> <p><b>do not use more than directed</b><br/>                 ■ if you forget a dose, do not double the next dose<br/>                 ■ do not spray into eyes or mouth<br/>                 ■ if allergy symptoms do not improve after one week, stop using and talk to a doctor<br/>                 ■ do not use for the common cold<br/>                 ■ shake well before each use</p> <p><b>Other information</b><br/>                 ■ do not use if sealed package is torn or opened<br/>                 ■ keep package and insert. They contain important information.<br/>                 ■ store between 20° - 25°C (68° - 77°F)</p> |  |
| <p>Dist. by: Chatterm, Inc.,<br/>                 a Sanofi Company<br/>                 Chattanooga, TN 37409-0219<br/>                 © 2016 Origin France</p>  |  | <p>UPC FPO</p>   |  |
| <p>LOT</p> <p>EXP</p> <p>XXXXXXXXXX</p>   |  | <p><b>25% MORE</b> › <b>BONUS</b> <sup>30 SPRAY</sup> BOTTLE</p> <p>Original Prescription Strength</p> <p>NDC XXXXX-XXXX-X</p> <p>Multi-Symptom Nasal Allergy Spray</p> <p><b>Nasacort® Allergy 24HR</b></p> <p>triamcinolone acetonide nasal spray 55 mcg per spray allergy symptom reliever (glucocorticoid)*</p> <p><b>24 HR</b><br/>                 Relief of:</p> <ul style="list-style-type: none"> <li>› Nasal Congestion</li> <li>› Sneezing</li> <li>› Runny Nose</li> <li>› Itchy Nose</li> </ul> <p><b>120</b> SPRAYS</p> <p><b>150</b> Total Sprays</p> <p>* Triamcinolone acetonide is a steroid medicine known as a glucocorticoid.</p> <p>One 120 spray bottle (16.9 ml) / One 30 spray bottle (6.7 ml)</p>  |  |

# Nasacort Allergy 24HR 120 Sprays Bonus Pack - Inside Panels

08.22.16

NAS\_Allergy\_120plus30\_Bonus\_r3



Prescription strength allergy relief.  
Without a prescription.

# Nasacort<sup>®</sup> Allergy 24HR

24 hour symptom relief  
from indoor & outdoor allergies

Read insert (inside package)  
for complete instructions  
on how to:

- › Get a new bottle ready (primed) before first use
- › Prime bottle again if not used for more than 2 weeks
- › Use the spray
- › Clean the spray nozzle

**BEFORE FIRST USE,  
A NEW BOTTLE  
MUST BE PRIMED**

**SHAKE WELL  
BEFORE EACH USE**



# Nasacort Allergy 24HR 120 Sprays Multi Pack

08.22.16

NAS\_Allergy\_120plus30\_MultiPack\_r4v2

|  |  |  |  |
|--|--|--|--|
| <p><b>Drug Facts</b></p> <p><b>Active ingredient (in each spray)</b>      <b>Purpose</b></p> <p>Triamcinolone acetonide<br/>55 mcg (glucocorticoid) .....Allergy symptom reliever</p> <p><b>Uses</b></p> <p>temporarily relieves these symptoms of hay fever or other upper respiratory allergies:</p> <ul style="list-style-type: none"> <li>nasal congestion</li> <li>runny nose</li> <li>sneezing</li> <li>itchy nose</li> </ul> <p><b>Warnings</b></p> <p><b>Do not use</b></p> <ul style="list-style-type: none"> <li>in children under 2 years of age</li> <li>if you have ever had an allergic reaction to any of the ingredients</li> </ul> <p><b>Ask a doctor before use if you</b></p> <ul style="list-style-type: none"> <li>have had recent nose ulcers or nose surgery</li> <li>have had a nose injury that has not healed</li> <li>are using a steroid medicine for asthma, allergies or skin rash</li> <li>have an eye infection</li> <li>have or had glaucoma or cataracts</li> </ul> <p><b>When using this product</b></p> <ul style="list-style-type: none"> <li>the growth rate of some children may be slower</li> <li>some symptoms may get better on the first day of treatment. It may take up to one week of daily use to feel the most symptom relief.</li> <li>do not share this bottle with anyone else as this may spread germs</li> <li>remember to tell your doctor about all the medicines you take, including this one</li> </ul> <p><b>Stop use and ask a doctor if</b></p> <ul style="list-style-type: none"> <li>you have, or come into contact with someone who has, chickenpox, measles or tuberculosis</li> <li>you have or develop symptoms of an infection such as a persistent fever</li> <li>you have any change in vision</li> <li>you have severe or frequent nosebleeds</li> </ul> <p><b>If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children.</b> In case of overdose, get medical help or contact a Poison Control Center right away.</p> |  | <p><b>Drug Facts (continued)</b></p> <p><b>Directions</b></p> <p>Read insert (inside package) on how to:</p> <ul style="list-style-type: none"> <li>get a new bottle ready (primed) before first use</li> <li>prime bottle again if not used for more than 2 weeks</li> <li>use the spray</li> <li>clean the spray nozzle</li> </ul> <p><b>ADULTS AND CHILDREN 12 YEARS OF AGE AND OLDER</b></p> <p><b>adults and children 12 years of age and older</b></p> <ul style="list-style-type: none"> <li>once daily, spray 2 times into each nostril while sniffing gently</li> <li>once your allergy symptoms improve, reduce to 1 spray in each nostril per day</li> </ul> <p><b>CHILDREN 2 TO UNDER 12 YEARS OF AGE</b></p> <ul style="list-style-type: none"> <li>the growth rate of some children may be slower while using this product. Talk to your child's doctor if your child needs to use the spray for longer than two months a year.</li> </ul> <p><b>children 6 to under 12 years of age</b></p> <ul style="list-style-type: none"> <li>an adult should supervise use</li> <li>once daily, spray 1 time into each nostril while sniffing gently</li> <li>if allergy symptoms do not improve, increase to 2 sprays in each nostril per day. Once allergy symptoms improve, reduce to 1 spray in each nostril per day.</li> </ul> <p><b>children 2 to under 6 years of age</b></p> <ul style="list-style-type: none"> <li>an adult should supervise use</li> <li>once daily, spray 1 time into each nostril while sniffing gently</li> </ul> <p><b>children under 2 years of age</b></p> <ul style="list-style-type: none"> <li>do not use</li> </ul> <p><b>do not use more than directed</b></p> <ul style="list-style-type: none"> <li>if you forget a dose, do not double the next dose</li> <li>do not spray into eyes or mouth</li> <li>if allergy symptoms do not improve after one week, stop using and talk to a doctor</li> <li>do not use for the common cold</li> <li>shake well before each use</li> </ul> <p><b>Other information</b></p> <ul style="list-style-type: none"> <li>do not use if sealed package is torn or opened</li> <li>keep package and insert. They contain important information.</li> <li>store between 20° - 25°C (68° - 77°F)</li> </ul> |  |
|--|--|--|--|

Dist. by: Chatterm, Inc.,  
a Sanofi Company  
Chattanooga, TN 37409-0219  
© 2016 Origin France

UPC FPO

LOT      EXP

XXXXXXXXXX

**MULTI-PACK** › **FREE** 30 SPRAY BOTTLE

Original Prescription Strength

NDC XXXXX-XXXX-X

Multi-Symptom Nasal Allergy Spray

## Nasacort® Allergy 24HR

triamcinolone acetonide nasal spray 55 mcg per spray  
allergy symptom reliever (glucocorticoid)\*

**24 HR**

Relief of:

- › Nasal Congestion
- › Sneezing
- › Runny Nose
- › Itchy Nose

**150**

Total Sprays

\* Triamcinolone acetonide is a steroid medicine known as a glucocorticoid.

One 120 spray bottle (16.9 ml) / One 30 spray bottle (6.7 ml)

# Nasacort Allergy 24HR 120 Sprays Multi Pack - Inside Panels

08.22.16

NAS\_Allergy\_120plus30\_MultiPack\_r4v2



# Nasacort Allergy 24HR Instructions - Front

08.24.16

NAS\_Allergy\_24HR\_INSERT

Product Information  
XXXXXXXXXX

## Nasacort<sup>®</sup> Allergy 24HR

### Multi-Symptom Nasal Allergy Spray

triamcinolone acetonide nasal spray 55 mcg per spray  
allergy symptom reliever  
(glucocorticoid)

#### IMPORTANT INFORMATION


- Read both sides of this insert for complete instructions on how to get a bottle ready (primed), how to use the spray bottle and how to clean the spray nozzle.
- Keep this insert as it contains important information.

#### STEPS TO GET A NEW BOTTLE READY FOR USE (PRIMED)

**1. Before first use, a new bottle must be primed.**

- a. Remove cap (Figure A).
- b. Shake bottle.
- c. Press and release spray nozzle until a fine mist is produced (as shown in Figure B). This may take several times. Take care not to spray in face.

Figure A



Cap  
Spray Nozzle  
Bottle


#### PRIME BOTTLE AGAIN IF NOT USED FOR MORE THAN 2 WEEKS

- Repeat steps for priming a bottle (see above).

#### USE INSTRUCTIONS

1. Blow nose gently to clear nostrils.
2. Remove cap, then shake bottle.
3. Hold bottle with thumb under bottle and spray nozzle between fingers (as shown in Figure B).
4. Press against the outside of your nose with your finger to close off one nostril (as shown in Figure C).
5. Place the tip of the spray nozzle into the other nostril. The spray nozzle should not reach far into the nose. Aim nozzle toward back of nose (as shown in Figure C).

Figure B



**DO NOT** spray toward nasal septum (the wall between the 2 nostrils) (as shown in Figure D).




Figure C  
**Correct Position**




Figure D  
**Wrong Position**


6. While sniffing gently, spray into nostril

**IMPORTANT: For complete dosing instructions, See "Directions for use" on next side.**

7. Repeat steps "4" through "6" for the other nostril.
8. After using the nasal spray, wipe nozzle with a tissue and replace cap by pressing it down over the spray nozzle.

*NOTE: Avoid blowing nose for 15 minutes after use.*

*If nozzle does not spray properly, see cleaning instructions on next side.*



(continued on side 2)

**SIDE 1**

# Nasacort Allergy 24HR Instructions - Back

08.24.16

NAS\_Allergy\_24HR\_INSERT

## DIRECTIONS FOR USE

- **BEFORE first use, you must get a new bottle ready (primed).**
- **If bottle is not used for more than 2 weeks, prime bottle again.**

### Dosing

#### ADULTS AND CHILDREN 12 YEARS OF AGE AND OLDER

- |  |   |
|--|---|
| <b>adults and children 12 years of age and older</b> | <ul style="list-style-type: none"><li>• once daily, spray 2 times into each nostril while sniffing gently</li><li>• once your allergy symptoms improve, reduce to 1 spray in each nostril per day</li></ul> |
|--|---|

#### CHILDREN 2 TO UNDER 12 YEARS OF AGE

- the growth rate of some children may be slower while using this product. Talk to your child's doctor if your child needs to use the spray for longer than two months a year.

- |  |  |
|--|--|
| <b>children 6 to under 12 years of age</b> | <ul style="list-style-type: none"><li>• an adult should supervise use</li><li>• once daily, spray 1 time into each nostril while sniffing gently</li><li>• if allergy symptoms do not improve, increase to 2 sprays in each nostril per day. Once allergy symptoms improve, reduce to 1 spray in each nostril per day.</li></ul> |
|--|--|

- |   |  |
|---|--|
| <b>children 2 to under 6 years of age</b> | <ul style="list-style-type: none"><li>• an adult should supervise use</li><li>• once daily, spray 1 time into each nostril while sniffing gently</li></ul> |
|---|--|


- |                                      |   |
|--------------------------------------|---|
| <b>children under 2 years of age</b> | <ul style="list-style-type: none"><li>• <b>do not use</b></li></ul> |
|--------------------------------------|---|

### Other important information for use

- Do not use more than directed.
- If you forget a dose, do **not** double the next dose.
- Do not spray into eyes or mouth.
- If you get the spray in your eyes, rinse well with water.
- If allergy symptoms do not improve after one week, stop using and talk to a doctor.
- Do not share this bottle with anyone else as this may spread germs.



### IF PUMP DOES NOT SPRAY PROPERLY, THE NOZZLE MAY BE BLOCKED

1. Never try to unblock nozzle with a pin or any other object (as shown in Figure E).
2. Clean the nozzle as shown below.



### CLEANING INSTRUCTIONS

1. Gently pull spray nozzle away from bottle (as shown in Figure F).
2. Rinse **SPRAY NOZZLE ONLY** under warm water (as shown in Figure G).
3. Shake or tap to remove excess water.
4. Re-attach spray nozzle to bottle.
5. Press and release spray nozzle until a fine spray is produced, taking care not to spray in face.



Nasacort Allergy 24HR pump is now ready to use.

### Where can I get more information?

1-800-633-1610

Store between 20°-25°C (68°-77°F)

Distributed by: Chattem, Inc., a Sanofi Company  
Chattanooga, TN 37409-0219 XXXXXX-XX © 2016 Chattem, Inc.



SIDE 2

# Nasacort Children's Allergy 24HR 60 LBL


08.23.16

NAS\_Childrens\_Allergy\_24HR\_60\_LBL


## PAGE 1 (Front Page)




## PAGE 2 (Inside Front Page)




## PAGE 3




## PAGE 4




## PAGE 5





## PAGE 6



## PAGE 7



## PAGE 8



Carton For IRC Sticker  
Children's Nasacort Allergy 24HR 60 Sprays  
08.23.16

NAS\_Childrens\_Allergy\_60 Sprays\_for IRC

|  |   |
|--|---|
| <p><b>Drug Facts</b></p>   | <p><b>Drug Facts (continued)</b></p>  |
| <p><b>Active ingredient (in each spray)</b>      <b>Purpose</b><br/>Triamcinolone acetonide<br/>55 mcg (glucocorticoid) .....Allergy symptom reliever</p>  | <p><b>Directions</b><br/>Read insert (inside package) on how to:<br/>■ get a new bottle ready (primed) before first use<br/>■ prime bottle again if not used for more than 2 weeks<br/>■ use the spray<br/>■ clean the spray nozzle</p>   |
| <p><b>Uses</b><br/>temporarily relieves these symptoms of hay fever or other upper respiratory allergies:<br/>■ nasal congestion   ■ runny nose   ■ sneezing   ■ itchy nose</p>  | <p><b>ADULTS AND CHILDREN 12 YEARS OF AGE AND OLDER</b><br/>adults and children 12 years of age and older<br/>■ once daily, spray 2 times into each nostril while sniffing gently<br/>■ once your allergy symptoms improve, reduce to 1 spray in each nostril per day</p>   |
| <p><b>Warnings</b><br/><b>Do not use</b><br/>■ in children under 2 years of age<br/>■ if you have ever had an allergic reaction to any of the ingredients</p>  | <p><b>CHILDREN 2 TO UNDER 12 YEARS OF AGE</b><br/>■ the growth rate of some children may be slower while using this product. Talk to your child's doctor if your child needs to use the spray for longer than two months a year.</p>  |
| <p><b>Ask a doctor before use if you</b><br/>■ have had recent nose ulcers or nose surgery<br/>■ have had a nose injury that has not healed<br/>■ are using a steroid medicine for asthma, allergies or skin rash<br/>■ have an eye infection<br/>■ have or had glaucoma or cataracts</p>  | <p>children 6 to under 12 years of age<br/>■ an adult should supervise use<br/>■ once daily, spray 1 time into each nostril while sniffing gently<br/>■ if allergy symptoms do not improve, increase to 2 sprays in each nostril per day. Once allergy symptoms improve, reduce to 1 spray in each nostril per day.</p> |
| <p><b>When using this product</b><br/>■ the growth rate of some children may be slower<br/>■ some symptoms may get better on the first day of treatment. It may take up to one week of daily use to feel the most symptom relief.<br/>■ do not share this bottle with anyone else as this may spread germs<br/>■ remember to tell your doctor about all the medicines you take, including this one</p> | <p>children 2 to under 6 years of age<br/>■ an adult should supervise use<br/>■ once daily, spray 1 time into each nostril while sniffing gently</p>  |
| <p><b>Stop use and ask a doctor if</b><br/>■ you have, or come into contact with someone who has, chickenpox, measles or tuberculosis<br/>■ you have or develop symptoms of an infection such as a persistent fever<br/>■ you have any change in vision<br/>■ you have severe or frequent nosebleeds</p>   | <p>children under 2 years of age<br/>■ do not use</p>   |
| <p><b>If pregnant or breast-feeding</b>, ask a health professional before use. <b>Keep out of reach of children.</b> In case of overdose, get medical help or contact a Poison Control Center right away.</p>  | <p><b>Other information</b><br/>■ do not use if sealed package is torn or opened<br/>■ keep package and insert. They contain important information.<br/>■ store between 20°- 25°C (68°- 77°F)</p>   |

Dist. by: Chattem, Inc.,  
a Sanofi Company  
Chattanooga, TN 37409-0219  
© 2016 Origin France

UPC FPO

Children's  
Nasacort®  
Allergy 24HR

**Drug Facts (continued)**  
**Inactive ingredients**  
benzalkonium chloride,  
carboxymethylcellulose sodium,  
dextrose, edetate disodium,  
hydrochloric acid or sodium  
hydroxide (for pH adjustment),  
microcrystalline cellulose,  
polysorbate 80, purified water

**Questions or comments?**  
call toll-free 1-800-633-1610

LOT  
EXP  
XXXXXXXXXX

NDC XXXXX-XXXX-X

Multi-Symptom  
Nasal Allergy Spray

ages 2 years & older

**Children's  
Nasacort®  
Allergy 24HR**

triamcinolone acetonide nasal spray 55 mcg per spray  
allergy symptom reliever  
(glucocorticoid)\*

**24 HR**  
Relief of:

- Nasal Congestion
- Sneezing
- Runny Nose
- Itchy Nose

0.37 fl. oz.  
(10.8 ml)

**60**  
SPRAYS

\* Triamcinolone acetonide is a steroid medicine known as a glucocorticoid.

Carton For IRC Sticker  
Children's Nasacort Allergy 24HR 60 Sprays - Inside Panels  
08.23.16

NAS\_Childrens\_Allergy\_60 Sprays\_for IRC



This label may not be the latest approved by FDA.  
For current labeling information, please visit <https://www.fda.gov/drugsatfda>

# Children's Nasacort Allergy 24HR 60 Sprays with Peel-Away IRC

08.23.16

NAS\_Childrens\_Allergy\_IRC



This label may not be the latest approved by FDA.  
For current labeling information, please visit <https://www.fda.gov/drugsatfda>

# Children's Nasacort Allergy 24HR 60 Sprays

## Rendering with IRC

### 08.23.16

NAS\_Childrens\_Allergy\_24HR\_IRC



# Children's Nasacort Allergy 24HR 60 Sprays Value Pack

## 08.22.16

NAS\_Childrens\_Allergy\_60x2\_ValuePack\_r1\_v2

|   |  |   |  |
|---|--|---|--|
| <p><b>Drug Facts</b></p> <p><b>Active ingredient (in each spray)</b>      <b>Purpose</b></p> <p>Triamcinolone acetonide<br/>55 mcg (glucocorticoid) .....Allergy symptom reliever</p> <p><b>Uses</b></p> <p>temporarily relieves these symptoms of hay fever or other upper respiratory allergies:</p> <ul style="list-style-type: none"> <li>nasal congestion</li> <li>runny nose</li> <li>sneezing</li> <li>itchy nose</li> </ul> <p><b>Warnings</b></p> <p><b>Do not use</b></p> <ul style="list-style-type: none"> <li>in children under 2 years of age</li> <li>if you have ever had an allergic reaction to any of the ingredients</li> </ul> <p><b>Ask a doctor before use if you</b></p> <ul style="list-style-type: none"> <li>have had recent nose ulcers or nose surgery</li> <li>have had a nose injury that has not healed</li> <li>are using a steroid medicine for asthma, allergies or skin rash</li> <li>have an eye infection</li> <li>have or had glaucoma or cataracts</li> </ul> <p><b>When using this product</b></p> <ul style="list-style-type: none"> <li>the growth rate of some children may be slower</li> <li>some symptoms may get better on the first day of treatment. It may take up to one week of daily use to feel the most symptom relief.</li> <li>do not share this bottle with anyone else as this may spread germs</li> <li>remember to tell your doctor about all the medicines you take, including this one</li> </ul> <p><b>Stop use and ask a doctor if</b></p> <ul style="list-style-type: none"> <li>you have, or come into contact with someone who has, chickenpox, measles or tuberculosis</li> <li>you have or develop symptoms of an infection such as a persistent fever</li> <li>you have any change in vision</li> <li>you have severe or frequent nosebleeds</li> </ul> <p><b>If pregnant or breast-feeding</b>, ask a health professional before use. <b>Keep out of reach of children.</b> In case of overdose, get medical help or contact a Poison Control Center right away.</p> |  | <p><b>Drug Facts (continued)</b></p> <p><b>Directions</b></p> <p>Read insert (inside package) on how to:</p> <ul style="list-style-type: none"> <li>get a new bottle ready (primed) before first use</li> <li>prime bottle again if not used for more than 2 weeks</li> <li>use the spray</li> <li>clean the spray nozzle</li> </ul> <p><b>ADULTS AND CHILDREN 12 YEARS OF AGE AND OLDER</b></p> <p><b>adults and children 12 years of age and older</b></p> <ul style="list-style-type: none"> <li>once daily, spray 2 times into each nostril while sniffing gently</li> <li>once your allergy symptoms improve, reduce to 1 spray in each nostril per day</li> </ul> <p><b>CHILDREN 2 TO UNDER 12 YEARS OF AGE</b></p> <ul style="list-style-type: none"> <li>the growth rate of some children may be slower while using this product. Talk to your child's doctor if your child needs to use the spray for longer than two months a year.</li> <li>an adult should supervise use</li> </ul> <p><b>children 6 to under 12 years of age</b></p> <ul style="list-style-type: none"> <li>once daily, spray 1 time into each nostril while sniffing gently</li> <li>if allergy symptoms do not improve, increase to 2 sprays in each nostril per day. Once allergy symptoms improve, reduce to 1 spray in each nostril per day.</li> </ul> <p><b>children 2 to under 6 years of age</b></p> <ul style="list-style-type: none"> <li>an adult should supervise use</li> <li>once daily, spray 1 time into each nostril while sniffing gently</li> </ul> <p><b>children under 2 years of age</b></p> <ul style="list-style-type: none"> <li><b>do not use</b></li> </ul> <p><b>do not use more than directed</b></p> <ul style="list-style-type: none"> <li>if you forget a dose, do not double the next dose</li> <li>do not spray into eyes or mouth</li> <li>if allergy symptoms do not improve after one week, stop using and talk to a doctor</li> <li>do not use for the common cold</li> <li>shake well before each use</li> </ul> <p><b>Other information</b></p> <ul style="list-style-type: none"> <li>do not use if sealed package is torn or opened</li> <li>keep package and insert. They contain important information.</li> <li>store between 20°- 25°C (68°- 77°F)</li> </ul> |  |
|---|--|---|--|

Dist. by: Chatterm, Inc.,  
a Sanofi Company  
Chattanooga, TN 37409-0219  
© 2016 Origin France

UPC FPO

LOT      EXP     

XXXXXXXXXX

**Children's Nasacort® Allergy 24HR**

**Drug Facts (continued)**

**Inactive ingredients**  
benzalkonium chloride, carboxymethylcellulose sodium, dextrose, edetate disodium, hydrochloric acid or sodium hydroxide (for pH adjustment), microcrystalline cellulose, polysorbate 80, purified water

**Questions or comments?**  
call toll-free 1-800-633-1610

VALUE-PACK › 2 BOTTLES  
COMPARE TO ONE 60 SPRAY BOTTLE

Original Prescription Strength

NDC XXXX-XXXX-X

Multi-Symptom Nasal Allergy Spray

**Children's Nasacort® Allergy 24HR**

ages 2 years & older

triamcinolone acetonide nasal spray 55 mcg per spray  
allergy symptom reliever (glucocorticoid)\*

**24 HR**  
Relief of:

- Nasal Congestion
- Sneezing
- Runny Nose
- Itchy Nose

60 SPRAYS

**120**  
Total Sprays

\* Triamcinolone acetonide is a steroid medicine known as a glucocorticoid.

Two bottles / 60 sprays per bottle (10.8 ml) each

# Children's Nasacort Allergy 24HR 60 Sprays Value Pack - Inside Panels

08.22.16

NAS\_Childrens\_Allergy\_60x2\_ValuePack\_r1\_v2



# Children's Nasacort Allergy 24HR Instructions - Front

08.24.16

NAS\_Childrens\_Allergy\_24HR\_INSERT

Product Information  
XXXXXXXXXX

## Children's Nasacort<sup>®</sup> Allergy 24HR

### Multi-Symptom Nasal Allergy Spray

triamcinolone acetate nasal spray 55 mcg per spray  
allergy symptom reliever  
(glucocorticoid)

### IMPORTANT INFORMATION


- Read both sides of this insert for complete instructions on how to get a bottle ready (primed), how to use the spray bottle and how to clean the spray nozzle.
- Keep this insert as it contains important information.

### STEPS TO GET A NEW BOTTLE READY FOR USE (PRIMED)

**1. Before first use, a new bottle must be primed.**

- Remove cap (Figure A).
- Shake bottle.
- Press and release spray nozzle until a fine mist is produced (as shown in Figure B). This may take several times. Take care not to spray in face.

Figure A



Cap

Spray Nozzle

Bottle


### PRIME BOTTLE AGAIN IF NOT USED FOR MORE THAN 2 WEEKS

- Repeat steps for priming a bottle (see above).

### USE INSTRUCTIONS


- Blow nose gently to clear nostrils.
- Remove cap, then shake bottle.
- Hold bottle with thumb under bottle and spray nozzle between fingers (as shown in Figure B).
- Press against the outside of your nose with your finger to close off one nostril (as shown in Figure C).
- Place the tip of the spray nozzle into the other nostril. The spray nozzle should not reach far into the nose. Aim nozzle toward back of nose (as shown in Figure C).

Figure B




**DO NOT** spray toward nasal septum (the wall between the 2 nostrils) (as shown in Figure D).

Figure C



Correct Position

Figure D



Wrong Position

- While sniffing gently, spray into nostril

**IMPORTANT: For complete dosing instructions, See "Directions for use" on next side.**

- Repeat steps "4" through "6" for the other nostril.
- After using the nasal spray, wipe nozzle with a tissue and replace cap by pressing it down over the spray nozzle.

*NOTE: Avoid blowing nose for 15 minutes after use.*

*If nozzle does not spray properly, see cleaning instructions on next side.*

10

(continued on side 2)

**SIDE 1**

# Children's Nasacort Allergy 24HR Instructions - Back

08.24.16

NAS\_Childrens\_Allergy\_24HR\_INSERT

## DIRECTIONS FOR USE

- **BEFORE first use, you must get a new bottle ready (primed).**
- **If bottle is not used for more than 2 weeks, prime bottle again.**

### Dosing

#### ADULTS AND CHILDREN 12 YEARS OF AGE AND OLDER

- |  |  |
|--|--|
| <b>adults and children 12 years of age and older</b> | <ul style="list-style-type: none"> <li>• once daily, spray 2 times into each nostril while sniffing gently</li> <li>• once your allergy symptoms improve, reduce to 1 spray in each nostril per day</li> </ul> |
|--|--|

#### CHILDREN 2 TO UNDER 12 YEARS OF AGE

- the growth rate of some children may be slower while using this product. Talk to your child's doctor if your child needs to use the spray for longer than two months a year.

- |  |  |
|--|--|
| <b>children 6 to under 12 years of age</b> | <ul style="list-style-type: none"> <li>• an adult should supervise use</li> <li>• once daily, spray 1 time into each nostril while sniffing gently</li> <li>• if allergy symptoms do not improve, increase to 2 sprays in each nostril per day. Once allergy symptoms improve, reduce to 1 spray in each nostril per day.</li> </ul> |
|--|--|

- |   |   |
|---|---|
| <b>children 2 to under 6 years of age</b> | <ul style="list-style-type: none"> <li>• an adult should supervise use</li> <li>• once daily, spray 1 time into each nostril while sniffing gently</li> </ul> |
|---|---|


- |                                      |   |
|--------------------------------------|---|
| <b>children under 2 years of age</b> | <ul style="list-style-type: none"> <li>• <b>do not use</b></li> </ul> |
|--------------------------------------|---|

### Other important information for use

- Do not use more than directed.
- If you forget a dose, do **not** double the next dose.
- Do not spray into eyes or mouth.
- If you get the spray in your eyes, rinse well with water.
- If allergy symptoms do not improve after one week, stop using and talk to a doctor.
- Do not share this bottle with anyone else as this may spread germs.


### IF PUMP DOES NOT SPRAY PROPERLY, THE NOZZLE MAY BE BLOCKED

1. Never try to unblock nozzle with a pin or any other object (as shown in Figure E).
2. Clean the nozzle as shown below.



### CLEANING INSTRUCTIONS

1. Gently pull spray nozzle away from bottle (as shown in Figure F).
2. Rinse **SPRAY NOZZLE ONLY** under warm water (as shown in Figure G).
3. Shake or tap to remove excess water.
4. Re-attach spray nozzle to bottle.
5. Press and release spray nozzle until a fine spray is produced, taking care not to spray in face.



**Children's Nasacort Allergy 24HR**  
 pump is now ready to use.

### Where can I get more information?

1-800-633-1610

Store between 20°-25°C (68°-77°F)



Distributed by: Chattem, Inc., a Sanofi Company  
 Chattanooga, TN 37409-0219 XXXXXX-XX © 2016 Chattem, Inc.

**SIDE 2**