


LATO STAMPA



Drug Facts (continued)

- changes in vision occur
- redness or irritation of the eye lasts
- condition worsens or lasts more than 72 hours

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away. Accidental swallowing by infants and children may lead to coma and marked reduction in body temperature.

Directions

- adults and children 6 years of age and over: put 1 or 2 drops in the affected eye(s) up to four times a day
- children under 6 years of age: consult a doctor

Other information

- store at 15° to 25°C (59° to 77°F)

Inactive ingredients
benzalkonium chloride, boric acid, edetate disodium, purified water, sodium borate, sodium hydroxide and/or hydrochloric acid

Questions?
888-734-7648; or Outside US, dial collect 215-273-8755

Distributed by:
JOHNSON & JOHNSON CONSUMER INC.
Skillman, NJ 08558 Made in Belgium
© J&JCI 2016
www.visine.com 30035144

VALUE TWIN PACK

Visine:A Sterile
ANTIHISTAMINE + REDNESS
RELIEVER EYE DROPS
MULTI-ACTION
EYE ALLERGY
Relief

Clinically proven to relieve itchy, red, allergy eyes

2 BOTTLES:
1/2 FL OZ (15 mL) each
NET 1 FL OZ (30 mL)

VALUE TWIN PACK

Visine:A Sterile
ANTIHISTAMINE + REDNESS
RELIEVER EYE DROPS
MULTI-ACTION
EYE ALLERGY
Relief

Clinically proven to relieve itchy, red, allergy eyes

2 BOTTLES:
1/2 FL OZ (15 mL) each
NET 1 FL OZ (30 mL)

Caution: Do not use if VISINE® imprinted neckband on bottle is broken or missing.

Drug Facts

Active ingredients	Purpose
Naphazoline HCl 0.025%.....	Redness reliever
Pheniramine maleate 0.3%.....	Antihistamine

Use Temporarily relieves itchy, red eyes due to: pollen ragweed grass animal hair and dander

Warnings
For external use only

Do not use if you are sensitive to any ingredient in this product.

Ask a doctor before use if you have

- heart disease
- high blood pressure
- narrow angle glaucoma
- trouble urinating

When using this product

- pupils may become enlarged temporarily causing light sensitivity
- do not touch tip of container to any surface to avoid contamination
- replace cap after each use
- remove contact lenses before using
- do not use if this solution changes color or becomes cloudy
- overuse may cause more eye redness
- some users may experience a brief stinging sensation

Stop use and ask a doctor if

- you feel eye pain

LOT:
EXP:

BOTTOM

3 12547 49370 3

STANDARD FORMAT
Helvetica Condensed
100% Horizontal Scale

Drug Facts - 9 pt Helv Bold Italic
Drug Facts (continued) -
Drug Facts - 8 pt Helv Bold Italic
(continued) - 8 pt Helv Reg
Headings - 8 pt Helv Bold Italic
Sub Heads - 6 pt Helv Bold
Text - 6 pt Helv Reg, 6.5 pt leading
Square Bullets - 5 pt
2.5 pt Outline Box
2.5 pt Barline
0.5 pt Hairline

MM 3/64" - Gap (Border-Type)

Net Content Statement

PDP Area: (width x length)	Net Content Size:	NET CONTENT	AREA OF PDP
1/2" H	7.148 in ²	5.50 IN OZ (150.00 ML)	5.50 IN OZ (150.00 ML)
1/4" H	.8875"	2.75 IN OZ (75.00 ML)	2.75 IN OZ (75.00 ML)
3/16" H		1.83 IN OZ (50.00 ML)	1.83 IN OZ (50.00 ML)
1/8" H		0.92 IN OZ (25.00 ML)	0.92 IN OZ (25.00 ML)
1/2" H		5.50 IN OZ (150.00 ML)	5.50 IN OZ (150.00 ML)



This is a representation of an electronic record that was signed electronically and this page is the manifestation of the electronic signature.

/s/

KAREN M MAHONEY

08/26/2016

Signing for Dr. Valerie Pratt