



Children's Motrin
 contains
Ibuprofen Oral Suspension (NSAID)
 Grape Flavor

NDC 50580-603-04

For Ages 2 to 11 Years
Children's Motrin
 contains **Ibuprofen Oral Suspension (NSAID)**
 100 mg per 5 mL
Pain Reliever/Fever Reducer

Lasts up to 8 hours

Alcohol Free

Grape Flavor
 4 fl oz (120 mL)

3 0045-0603-04 3

LOT:

EXP:

Drug Facts (continued)
Ask a doctor before use if
 ■ stomach bleeding warning applies to your child
 ■ child has a history of stomach problems, such as heartburn
 ■ child has problems or serious side effects from taking pain relievers or fever reducers
 ■ child has not been drinking fluids
 ■ child has lost a lot of fluid due to vomiting or diarrhea
 ■ child has high blood pressure, heart disease, liver cirrhosis, or kidney disease
 ■ child has asthma
 ■ child is taking a diuretic
Ask a doctor or pharmacist before use if the child is
 ■ under a doctor's care for any serious condition
 ■ taking any other drug
When using this product
 ■ take with food or milk if stomach upset occurs
 ■ the risk of heart attack or stroke may increase if you use more than directed or for longer than directed
Stop use and ask a doctor if
 ■ child experiences any of the following signs of stomach bleeding
 ■ feels faint ■ vomits blood
 ■ has bloody or black stools
 ■ has stomach pain that does not get better
 ■ the child does not get any relief within first day (24 hours) of treatment
 ■ fever or pain gets worse or lasts more than 3 days
 ■ redness or swelling is present in the painful area
 ■ any new symptoms appear
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions
 ■ this product does not contain directions or complete warnings for adult use
 ■ do not give more than directed
 ■ shake well before using
 ■ mL = milliliter; tsp = teaspoonful
 ■ find right dose on chart - if possible, use weight to dose; otherwise use age.
 ■ use only enclosed dosing cup. Do not use any other dosing device.

Drug Facts (continued)
 ■ If needed, repeat dose every 6-8 hours
 ■ do not use more than 4 times a day
 ■ replace original bottle cap to maintain child resistance
Dosing Chart

Weight (lb)	Age (yr)	Dose (mL or tsp)*
under 24	under 2 years	ask a doctor
24-35 lbs	2-3 years	5 mL (1 tsp)
36-47 lbs	4-5 years	7.5 mL (1½ tsp)
48-59 lbs	6-8 years	10 mL (2 tsp)
60-71 lbs	9-10 years	12.5 mL (2½ tsp)
72-95 lbs	11 years	15 mL (3 tsp)

* or as directed by a doctor

Other information
 ■ each 5 mL (1 tsp) contains: sodium 2 mg
 ■ store between 20-25°C (68-77°F)
 ■ do not use if carton is opened or if carton tape or bottle wrap imprinted "SAFETY SEAL®" is broken or missing

Inactive ingredients acesulfame potassium, anhydrous citric acid, D&C red no. 33, FD&C blue no. 1, FD&C red no. 40, flavors, glycerin, polysorbate 80, pregelatinized starch, purified water, sodium benzoate, sucrose, xanthan gum

Questions or comments?
 Call 1-877-895-3665 (toll-free) or 215-273-8755 (collect)

McNeil
 Distributed by:
 McNeil Consumer Healthcare
 Division of McNEIL-PPC, Inc.
 Fort Washington, PA 19034 USA
 © McNEIL-PPC, INC. 2015
www.motrin.com 00000000

Important: Read all product information before using. Keep this box for important information. This product is intended for use in children.

Drug Facts
Active ingredient Purpose (in each 5 mL = 1 teaspoonful)
 Ibuprofen 100 mg (NSAID)*.....Pain reliever/fever reducer
 *nonsteroidal anti-inflammatory drug

Uses temporarily:
 ■ relieves minor aches and pains due to the common cold, flu, sore throat, headache and toothache
 ■ reduces fever

Warnings
Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:
 ■ hives ■ facial swelling
 ■ asthma (wheezing) ■ shock
 ■ skin reddening ■ rash ■ blisters
 If an allergic reaction occurs, stop use and seek medical help right away.
Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if your child:
 ■ has had stomach ulcers or bleeding problems
 ■ takes a blood thinning (anticoagulant) or steroid drug
 ■ takes other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
 ■ takes more or for a longer time than directed
Sore throat warning: Severe or persistent sore throat or sore throat accompanied by high fever, headache, nausea, and vomiting may be serious. Consult doctor promptly. Do not use more than 2 days or administer to children under 3 years of age unless directed by doctor.
Do not use
 ■ if the child has ever had an allergic reaction to any other pain reliever/fever reducer
 ■ right before or after heart surgery

USA
 000000
 000

63,5 mm
146 mm

NDC 50580-683-04

Children's Motrin®

contains **Ibuprofen**
Oral Suspension (NSAID)
 100 mg per 5 mL

Pain Reliever/Fever Reducer 4 fl oz (120 mL)

Grape Flavor

Important: Read all product information before using. Keep the box for important information. This product is intended for use in children.

Active ingredient (in each 5 mL = 1 teaspoonful) **Purpose**
 Ibuprofen 100 mg (NSAID) Pain reliever/fever reducer
 Nonsteroidal anti-inflammatory drug

Uses: Temporarily relieves minor aches and pains due to the common cold, flu, sore throat, headache and toothache; reduces fever.

Warnings:
Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: hives; facial swelling; wheezing; shock; skin redness; or rash/blebules. If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if your child has had stomach ulcers or bleeding problems, takes a blood-thinning (anti-coagulant) or steroid drug, or takes other prescription or non-prescription NSAIDs (aspirin, ibuprofen, naproxen, or others). Do not use more or for a longer time than directed. **Stomach warning:** Severe or persistent sore throat or sore throat accompanied by high fever, headache, nausea, and vomiting may be serious. Consult doctor promptly. Do not use more than 2 days or administer to children under 3 years of age unless directed by doctor. Do not use if the child has ever had an allergic reaction to any other pain reliever/fever reducer, or right before or after an allergy shot. **Ask a doctor before use if:** stomach bleeding warning applies to you or child; child has a history of stomach problems, such as heartburn; child has problems or serious side effects from taking pain relievers or fever reducers; child has not been drinking fluids; child has lost a lot of fluid due to vomiting or diarrhea; child has high blood pressure; heart disease; asthma; or kidney disease; child has asthma; child is taking a diuretic. **Ask a doctor or pharmacist before use if the child is under a doctor's care for any serious condition or is taking any other drug.** When using this product, do not give food or milk. If stomach upset occurs, do not use more than directed or for longer than directed. Stop use and ask a doctor if child experiences any of the following signs of stomach bleeding: vomit that is bloody or black stools; or if stomach pain that does not go away; or if the child does not get any relief within 1 day (24 hours) of the stomach pain; or pain gets worse or lasts more than 3 days; or redness or swelling is present in the mouth or on any new symptoms appear. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222). **Directions:** This product does not contain directions or complete warnings for adult use. Do not give more than directed. Shake well before using. mL = milliliter; tsp = 1 teaspoon; 1/2 tsp = 1/2 teaspoon. If possible, use weight to dose, otherwise use age. Use only enclosed dosing cup. Do not use any other dosing device. If needed, repeat dose every 6-8 hours. Do not use more than 4 times a day. Replace original bottle cap to maintain freshness.

Weight (lb)	Age (yr)	Dose (mL or tsp)*
under 24	under 2 years	ask a doctor
24-35 lbs	2-3 years	5 mL (1 tsp)
36-47 lbs	4-5 years	7.5 mL (1 1/2 tsp)
48-59 lbs	6-8 years	10 mL (2 tsp)
60-71 lbs	9-10 years	12.5 mL (2 1/2 tsp)
72-95 lbs	11 years	15 mL (3 tsp)

*or as directed by a doctor

Other information: Each 5 mL (1 tsp) contains sodium 2 mg, inactive between 20-25°C (68-77°F). Do not use if bottle is unimprinted "SAFETY SEAL" is broken or missing. **Active ingredients:** sodium ibuprofen, potassium, ethylhexylcinnamate, D-C-6-ethyl-11,11-dimethyl-2,2,2-trifluoroethane-1,1-diol, FD&C blue no. 1, FD&C red no. 40, flavor, glycerin, polyethylene glycol, purified water, sodium benzoate, sucrose, anthranquinone. **Directions or comments?** Call 1-877-899-5683 (toll-free) or 215-273-8789 (collect).

Manufactured by
 Mallinckrodt Pharmaceuticals, Inc.
 One Mallinckrodt Place
 Fort Washington, PA 19074 USA
 © Mallinckrodt, Inc. 2015
 www.childrensmotrin.com

LOT: XXXXXXXXXX

EXP: XXXXXXXXXX

(b) (4)

Reference ID: 3733010



Drug Facts (continued)

Ask a doctor before use if

- stomach bleeding warning applies to your child
- child has a history of stomach problems, such as heartburn
- child has problems or serious side effects from taking pain relievers or fever reducers
- child has not been drinking fluids
- child has lost a lot of fluid due to vomiting or diarrhea
- child has high blood pressure, heart disease, liver cirrhosis, or kidney disease
- child has asthma
- child is taking a diuretic

Ask a doctor or pharmacist before use if the child is

- under a doctor's care for any serious condition
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs
- the risk of heart attack or stroke may increase if you use more than directed or for longer than directed

Stop use and ask a doctor if

- child experiences any of the following signs of stomach bleeding
 - feels faint
 - vomits blood
 - has bloody or black stools
- has stomach pain that does not get better
- the child does not get any relief within first day (24 hours) of treatment
- fever or pain gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions

- this product does not contain directions or complete warnings for adult use
- do not give more than directed
- shake well before using
- mL = milliliter; tsp = teaspoonful
- find right dose on chart— if possible, use weight to dose; otherwise use age.
- use only enclosed dosing cup. Do not use any other dosing device.

Drug Facts (continued)

- if needed, repeat dose every 6-8 hours
- do not use more than 4 times a day
- replace original bottle cap to maintain child resistance

Dosing Chart

Weight (lb)	Age (yr)	Dose (mL or tsp)*
under 24	under 2 years	ask a doctor
24-35 lbs	2-3 years	5 mL (1 tsp)
36-47 lbs	4-5 years	7.5 mL (1½ tsp)
48-59 lbs	6-8 years	10 mL (2 tsp)
60-71 lbs	9-10 years	12.5 mL (2½ tsp)
72-95 lbs	11 years	15 mL (3 tsp)

* or as directed by a doctor

Other information

- each 5 mL (1 tsp) contains: sodium 2 mg
- store between 20-25°C (68-77°F)
- do not use if carton is opened or if carton tape or bottle wrap imprinted "SAFETY SEAL®" is broken or missing

Inactive ingredients acesulfame potassium, anhydrous citric acid, D&C yellow no. 10, FD&C red no. 40, flavors, glycerin, polysorbate 80, pregelatinized starch, purified water, sodium benzoate, sucrose, xanthan gum

Questions or comments?
Call 1-877-895-3665 (toll-free) or 215-273-8755 (collect)

McNeil

Distributed by:
McNeil Consumer Healthcare
Division of McNEIL-PPC, Inc.
Fort Washington, PA 19034 USA
© McNEIL-PPC, INC. 2015
www.motrin.com 00000000

Children's Motrin
contains
Ibuprofen Oral Suspension (NSAID)
Original Berry Flavor

NDC 50580-601-20

For Ages 2 to 11 Years

Children's Motrin
contains Ibuprofen
Oral Suspension (NSAID)
100 mg per 5 mL
Pain Reliever/Fever Reducer

Lasts up to
8 hours

Alcohol Free

Original
Berry Flavor

Important: Read all product information before using. Keep this box for important information. This product is intended for use in children.

Drug Facts

Active ingredient	Purpose
Ibuprofen 100 mg (NSAID)*	Pain reliever/fever reducer

*nonsteroidal anti-inflammatory drug

Uses temporarily:

- relieves minor aches and pains due to the common cold, flu, sore throat, headache and toothache
- reduces fever

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if your child:

- has had stomach ulcers or bleeding problems
- takes a blood thinning (anticoagulant) or steroid drug
- takes other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- takes more or for a longer time than directed

Sore throat warning: Severe or persistent sore throat or sore throat accompanied by high fever, headache, nausea, and vomiting may be serious. Consult doctor promptly. Do not use more than 2 days or administer to children under 3 years of age unless directed by doctor.

Do not use

- if the child has ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

3 0045-0192-04 2

LOT:

EXP:

USA
000000
000





Drug Facts (continued)

Ask a doctor before use if

- stomach bleeding warning applies to your child
- child has a history of stomach problems, such as heartburn
- child has problems or serious side effects from taking pain relievers or fever reducers
- child has not been drinking fluids
- child has lost a lot of fluid due to vomiting or diarrhea
- child has high blood pressure, heart disease, liver cirrhosis, or kidney disease
- child has asthma
- child is taking a diuretic

Ask a doctor or pharmacist before use if the child is

- under a doctor's care for any serious condition
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs
- the risk of heart attack or stroke may increase if you use more than directed or for longer than directed

Stop use and ask a doctor if

- child experiences any of the following signs of stomach bleeding
 - feels faint
 - vomits blood
 - has bloody or black stools
- has stomach pain that does not get better
- the child does not get any relief within first day (24 hours) of treatment
- fever or pain gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions

- this product does not contain directions or complete warnings for adult use
- do not give more than directed
- shake well before using
- mL = milliliter; tsp = teaspoonful
- find right dose on chart - if possible, use weight to dose; otherwise use age.
- use only enclosed dosing cup. Do not use any other dosing device.

Drug Facts (continued)

- if needed, repeat dose every 6-8 hours
- do not use more than 4 times a day
- replace original bottle cap to maintain child resistance

Dosing Chart

Weight (lb)	Age (yr)	Dose (mL or tsp)*
under 24	under 2 years	ask a doctor
24-35 lbs	2-3 years	5 mL (1 tsp)
36-47 lbs	4-5 years	7.5 mL (1½ tsp)
48-59 lbs	6-8 years	10 mL (2 tsp)
60-71 lbs	9-10 years	12.5 mL (2½ tsp)
72-95 lbs	11 years	15 mL (3 tsp)

* or as directed by a doctor

Other information

- each 5 mL (1 tsp) contains: sodium 2 mg
- store between 20-25°C (68-77°F)
- do not use if carton is opened or if carton tape or bottle wrap imprinted "SAFETY SEAL®" is broken or missing

Inactive ingredients acesulfame potassium, anhydrous citric acid, FD&C red no. 40, flavors, glycerin, polysorbate 80, pregelatinized starch, purified water, sodium benzoate, sucrose, xanthan gum

Questions or comments?
 Call 1-877-895-3665 (toll-free) or 215-273-8755 (collect)

McNeil

Distributed by:
 McNeil Consumer Healthcare
 Division of McNEIL-PPC, Inc.
 Fort Washington, PA 19034 USA
 © McNEIL-PPC, INC. 2015
www.motrin.com 00000000

Children's Motrin
 contains
Ibuprofen Oral Suspension (NSAID)
 Bubble Gum Flavor

For Ages 2 to 11 Years
Children's Motrin
 contains **Ibuprofen Oral Suspension**
 100 mg per 5 mL
 Lasts up to 8 hours
 Bubble Gum Flavor
 4 fl oz (120 ml)

3 0045-0604-04 0
 LOT:
 EXP:

Important: Read all product information before using. Keep this box for important information. This product is intended for use in children.

Drug Facts

Active ingredient (in each 5 mL = 1 teaspoonful)	Purpose
Ibuprofen 100 mg (NSAID)*	Pain reliever/fever reducer

*nonsteroidal anti-inflammatory drug

Uses temporarily:

- relieves minor aches and pains due to the common cold, flu, sore throat, headache and toothache
- reduces fever

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if your child:

- has had stomach ulcers or bleeding problems
- takes a blood thinning (anticoagulant) or steroid drug
- takes other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- takes more or for a longer time than directed

Sore throat warning: Severe or persistent sore throat or sore throat accompanied by high fever, headache, nausea, and vomiting may be serious. Consult doctor promptly. Do not use more than 2 days or administer to children under 3 years of age unless directed by doctor.

Do not use

- if the child has ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

USA
 000000
 000

63,5 mm
146 mm

NDC 50580-684-04

Children's Motrin®

contains **Ibuprofen**
Oral Suspension (NSAID)
 100 mg per 5 mL

Pain Reliever/Fever Reducer 4 fl oz (120 mL)

Bubble Gum Flavor

Important: Read all product information before using. Keep the box for important information. This product is intended for use in children.

Active ingredient (in each 5 mL = 1 teaspoonful) **Purpose**
 Ibuprofen 100 mg (NSAID) Pain reliever/fever reducer
 Nonsteroidal anti-inflammatory drug

Uses: Helps to relieve minor aches and pains due to the common cold, flu, sore throat, headache and toothache; reduces fever

Warnings:
Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: hives; facial swelling; wheezing; shock; skin redness; rash; blisters. If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if your child has had stomach ulcers or bleeding problems, takes a blood-thinning (anti-coagulant) or steroid drug, or takes other prescription or non-prescription NSAIDs (aspirin, ibuprofen, naproxen, or others). Do not use more or for a longer time than directed. **Stomach warning:** Severe or persistent sore throat or sore throat accompanied by high fever, headache, nausea, and vomiting may be serious. Consult doctor promptly. Do not use more than 2 days or administer to children under 3 years of age unless directed by doctor. Do not use with other pain relievers/fever reducers. **Weight before or after surgery:** Ask a doctor before use if a stomach bleeding warning applies to your child or if the child has a history of stomach problems, such as heartburn or if the child has problems or serious side effects from taking pain relievers or fever reducers. **Child has not been drinking:** If a child has lost a lot of fluid due to vomiting or diarrhea, or if a child has high blood pressure, heart disease, liver or kidney disease, or if a child has asthma, ask a doctor or pharmacist before use. **Child is under a doctor's care for any serious condition:** Do not use without a doctor's approval. **When using this product:** Do not give to a child who is taking any other medicine. Do not use more than directed or for longer than directed. Stop use and ask a doctor if a child experiences any of the following signs of stomach bleeding: vomit that is bloody or black stools; or if a child has pain that does not get better. **Do not use if a child does not get any relief within 1 day (24 hours) of the start of fever or pain, gets worse or lasts more than 3 days, or redness or swelling is present in the painful area, or any new symptoms appear.** Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222). **Directions:** See this product does not contain directions or complete warnings for adult use. Do not give more than directed. **Shake well before using.** mL = milliliter; tsp = 1 teaspoon; 1 tsp = 5 mL. **Do not use if you are pregnant or breastfeeding.** If possible, use weight to dose; otherwise use age. **Use only enclosed dosing cup.** Do not use any other dosing device. **Use:** Feed, repeat dose every 6-8 hours; do not use more than 4 times a day; replace original bottle cap to maintain freshness.

Weight (lb)	Age (yr)	Dose (mL or tsp)*
under 24	under 2 years	ask a doctor
24-35 lbs	2-5 years	5 mL (1 tsp)
36-47 lbs	4-5 years	7.5 mL (1½ tsp)
48-59 lbs	6-8 years	10 mL (2 tsp)
60-71 lbs	9-10 years	12.5 mL (2½ tsp)
72-95 lbs	11 years	15 mL (3 tsp)

*or as directed by a doctor

Other information: Each 5 mL (1 tsp) contains sodium 2 mg, inactive between 20-25°C (68-77°F). Do not use if bottle cap is imprinted "SAFETY SEAL" is broken or missing. **Active ingredients:** sodium ferrous succinate, sodium ferrous citrate, FDAC, red no. 40, flavo s, glycerol, poly sorbate 80, polyethylene glycol 400, purified water, sodium benzoate, sucrose, xanthan gum

Questions or comments? Call 1-877-899-5645 (toll-free) or 215-273-8755 (collect)

Manufactured by
 Mallinckrodt Pharmaceuticals, Inc.
 6140 Rockledge Drive, Suite 100
 Fort Washington, PA 19074, USA
 © Mallinckrodt, Inc. 2015
 www.childrensmotrin.com

LOT: XXXXXXXXXX

EXP: XXXXXXXXXX

(b) (4)

Reference ID: 3733010

w 3.9375"

Convenient 2-PACK!

NDC 50580-601-25 NDC 50580-601-25

For Ages 2 to 11 Years *For Ages 2 to 11 Years*

Children's Motrin® **Children's Motrin®**

contains ibuprofen
Oral Suspension (NSAID)
100 mg per 5 mL

Pain Reliever / Fever Reducer


Lasts up to 8 hours

Alcohol Free

Original Berry Flavor

Distributed by: McNeil Consumer Healthcare
Division of McNEIL-PPC, Inc.
Fort Washington, PA 19034 USA
©McNeil-PPC, INC. www.motrin.com

McN-XX-0000




3 0045-0192-08 0

h 5.6562"

h 0.7"

(b) (4)



(b) (4)

This is a representation of an electronic record that was signed electronically and this page is the manifestation of the electronic signature.

/s/

THERESA M MICHELE
04/16/2015