

Trim Safe Area Bleed Score

w 3.9375"

**Convenient 2-PACK!**

NDC 50580-601-25 For Ages 2 to 11  
**Children's Motrin®**  
contains **Ibuprofen**  
**Oral Suspension (NSAID)**  
100 mg per 5 mL (teaspoon)  
**Pain Reliever / Fever Reducer**

NDC 50580-601-25 For Ages 2 to 11  
**Children's Motrin®**  
contains **Ibuprofen**  
**Oral Suspension (NSAID)**  
100 mg per 5 mL (teaspoon)  
**Pain Reliever / Fever Reducer**

h 5.6562"

**Lasts up to 8 hours**  
**Alcohol Free**

**Lasts up to 8 hours**  
**Alcohol Free**

**4 fl oz (120 ml)** **4 fl oz (120 ml)**

Distributed by: McNeil Consumer Healthcare  
DIVISION OF MCNEIL - PPC, NC.  
FORT WASHINGTON, PA 19034 USA  
© MCN-PPC, NC. 2012 [www.motrin.com](http://www.motrin.com)  
McN-CM-0207

3 0045-0192-08 0

h 0.7"

Front

(b) (4)



**Drug Facts** (continued)

**Ask a doctor before use if**

- stomach bleeding warning applies to your child
- child has a history of stomach problems, such as heartburn
- child has problems or serious side effects from taking pain relievers or fever reducers
- child has not been drinking fluids
- child has lost a lot of fluid due to vomiting or diarrhea
- child has high blood pressure, heart disease, liver cirrhosis, or kidney disease
- child has asthma
- child is taking a diuretic

**Ask a doctor or pharmacist before use if the child is**

- under a doctor's care for any serious condition
- taking any other drug

**When using this product**

- take with food or milk if stomach upset occurs
- the risk of heart attack or stroke may increase if you use more than directed or for longer than directed

**Stop use and ask a doctor if**

- child experiences any of the following signs of stomach bleeding
  - feels faint
  - vomits blood
  - has bloody or black stools
- has stomach pain that does not get better
- the child does not get any relief within first day (24 hours) of treatment
- fever or pain gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

**Directions**

- this product does not contain directions or complete warnings for adult use
- do not give more than directed
- shake well before using
- find right dose on chart. If possible, use weight to dose; otherwise use age
- use only enclosed measuring cup
- if needed, repeat dose every 6-8 hours
- do not use more than 4 times a day

**Drug Facts** (continued)

- replace original bottle cap to maintain child resistance

**Dosing Chart**

Weight (lb)	Age (yr)	Dose (tsp or mL)
under 2 years		ask a doctor
24-35 lbs	2-3 years	1 tsp or 5 mL
36-47 lbs	4-5 years	1 ½ tsp or 7.5 mL
48-59 lbs	6-8 years	2 tsp or 10 mL
60-71 lbs	9-10 years	2 ½ tsp or 12.5 mL
72-95 lbs	11 years	3 tsp or 15 mL

**Other information**

- each teaspoon contains: sodium 2 mg
- store between 20-25°C (68-77°F)
- do not use if carton is opened or if carton tape or bottle wrap imprinted "SAFETY SEAL®" is broken or missing
- see bottom panel for lot number and expiration date

**Inactive ingredients**

acesulfame potassium, anhydrous citric acid, D&C yellow no. 10, FD&C red no. 40, flavors, glycerin, polysorbate 80, pregelatinized starch, purified water, sodium benzoate, sucrose, xanthan gum

**Questions or comments?**

Call 1-877-895-3665: weekdays 8:00 AM to 8:00 PM EST



Distributed by:  
 McNeil Consumer Healthcare  
 DIVISION OF MCNEIL-PPC, INC.  
 FORT WASHINGTON, PA 19034 USA  
 © MCN-PPC, INC. 2011  
[www.motrin.com](http://www.motrin.com)

**Important: Read all product information before using. Keep this box for important information. This product is intended for use in children.**

**Drug Facts**

**Active ingredient Purpose (in each 5 mL = 1 teaspoon)**

Ibuprofen 100 mg (NSAID)\*.....Pain reliever/  
 \*nonsteroidal anti-inflammatory drug fever reducer

**Uses temporarily:**

- relieves minor aches and pains due to the common cold, flu, sore throat, headache and toothache
- reduces fever

**Warnings**

**Allergy alert:** Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if your child:



- has had stomach ulcers or bleeding problems
- takes a blood thinning (anticoagulant) or steroid drug
- takes other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)

■ takes more or for a longer time than directed  
**Sore throat warning:** Severe or persistent sore throat or sore throat accompanied by high fever, headache, nausea, and vomiting may be serious. Consult doctor promptly. Do not use more than 2 days or administer to children under 3 years of age unless directed by doctor.

**Do not use**

- if the child has ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

USA  
 981804  
 905



(b) (4)

---

**This is a representation of an electronic record that was signed electronically and this page is the manifestation of the electronic signature.**

---

/s/

---

KAREN M MAHONEY  
12/11/2014