

Drug Facts (continued)

Ask a doctor before use if

- stomach bleeding warning applies to your child
- child has a history of stomach problems, such as heartburn
- child has problems or serious side effects from taking pain relievers or fever reducers
- child has not been drinking fluids
- child has lost a lot of fluid due to vomiting or diarrhea
- child has high blood pressure, heart disease, liver cirrhosis, or kidney disease
- child has asthma
- child is taking a diuretic

Ask a doctor or pharmacist before use if the child is

- under a doctor's care for any serious condition
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs
- the risk of heart attack or stroke may increase if you use more than directed or for longer than directed

Stop use and ask a doctor if

- child experiences any of the following signs of stomach bleeding
 - feels faint
 - vomits blood
 - has bloody or black stools
- has stomach pain that does not get better
- the child does not get any relief within first day (24 hours) of treatment
- fever or pain gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions

- this product does not contain directions or complete warnings for adult use
- do not give more than directed
- shake well before using
- mL = milliliter
- find right dose on chart. If possible, use weight to dose; otherwise use age.
- use only enclosed dosing cup. Do not use any other dosing device.

Drug Facts (continued)

- if needed, repeat dose every 6-8 hours
- do not use more than 4 times a day
- replace original bottle cap to maintain child resistance

Dosing Chart

Weight (lb)	Age (yr)	Dose (mL)*
under 24	under 2 years	ask a doctor
24-35 lbs	2-3 years	5 mL
36-47 lbs	4-5 years	7.5 mL
48-59 lbs	6-8 years	10 mL
60-71 lbs	9-10 years	12.5 mL
72-95 lbs	11 years	15 mL

* or as directed by a doctor

Other information

- each 5 mL contains: sodium 2 mg
- store between 20-25° C (68-77°F)
- do not use if carton is opened or if carton tape or bottle wrap imprinted "SAFETY SEAL" is broken or missing

Inactive ingredients acesulfame potassium, anhydrous citric acid, D&C red no. 33, FD&C blue no. 1, FD&C red no. 40, flavors, glycerin, polysorbate 80, pregelatinized starch, purified water, sodium benzoate, sucrose, xanthan gum

Questions or comments?
 Call 1-877-895-3665 (toll-free) or 215-273-8755 (collect)

Distributed by:
JOHNSON & JOHNSON CONSUMER INC.
 McNeil Consumer Healthcare Division
 Fort Washington, PA 19034 USA
 ©J&JCI 2015
www.motrin.com 00000000

Children's Motrin
 contains
 Ibuprofen Oral Suspension (NSAID)
 Grape Flavor

New Information: Read Entire Label
 NDC 50580-603-04

For Ages 2 to 11 Years

Children's Motrin
 contains Ibuprofen
 Oral Suspension (NSAID)
 100 mg per 5 mL
 Pain Reliever/Fever Reducer

Lasts up to 8 hours

Alcohol Free

Grape Flavor
 4 fl oz (120 mL)

Important: Read all product information before using. Keep this box for important information. This product is intended for use in children.

Drug Facts

Active ingredient (in each 5 mL)	Purpose
Ibuprofen 100 mg (NSAID)*	Pain reliever/fever reducer

*nonsteroidal anti-inflammatory drug

Uses temporarily:

- relieves minor aches and pains due to the common cold, flu, sore throat, headache and toothache
- reduces fever

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if your child:

- has had stomach ulcers or bleeding problems
- takes a blood thinning (anticoagulant) or steroid drug
- takes other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- takes more or for a longer time than directed

Sore throat warning: Severe or persistent sore throat or sore throat accompanied by high fever, headache, nausea, and vomiting may be serious. Consult doctor promptly. Do not use more than 2 days or administer to children under 3 years of age unless directed by doctor.

Do not use

- if the child has ever had an allergic reaction to ibuprofen or any other pain reliever/fever reducer
- right before or after heart surgery

3 0045-0603-04 3





Drug Facts (continued)

Ask a doctor before use if

- stomach bleeding warning applies to your child
- child has a history of stomach problems, such as heartburn
- child has problems or serious side effects from taking pain relievers or fever reducers
- child has not been drinking fluids
- child has lost a lot of fluid due to vomiting or diarrhea
- child has high blood pressure, heart disease, liver cirrhosis, or kidney disease
- child has asthma
- child is taking a diuretic

Ask a doctor or pharmacist before use if the child is

- under a doctor's care for any serious condition
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs
- the risk of heart attack or stroke may increase if you use more than directed or for longer than directed

Stop use and ask a doctor if

- child experiences any of the following signs of stomach bleeding
 - feels faint
 - vomits blood
 - has bloody or black stools
- has stomach pain that does not get better
- the child does not get any relief within first day (24 hours) of treatment
- fever or pain gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions

- this product does not contain directions or complete warnings for adult use
- do not give more than directed
- shake well before using
- mL = milliliter
- find right dose on chart. If possible, use weight to dose; otherwise use age.
- use only enclosed dosing cup. Do not use any other dosing device.

Drug Facts (continued)

- if needed, repeat dose every 6-8 hours
- do not use more than 4 times a day
- replace original bottle cap to maintain child resistance

Dosing Chart

Weight (lb)	Age (yr)	Dose (mL)*
under 24	under 2 years	ask a doctor
24-35 lbs	2-3 years	5 mL
36-47 lbs	4-5 years	7.5 mL
48-59 lbs	6-8 years	10 mL
60-71 lbs	9-10 years	12.5 mL
72-95 lbs	11 years	15 mL

* or as directed by a doctor

Other information

- each 5 mL contains: sodium 2 mg
- store between 20-25° C (68-77°F)
- do not use if carton is opened or if carton tape or bottle wrap imprinted "SAFETY SEAL" is broken or missing

Inactive ingredients acesulfame potassium, anhydrous citric acid, FD&C red no. 40, flavors, glycerin, polysorbate 80, pregelatinized starch, purified water, sodium benzoate, sucrose, xanthan gum

Questions or comments?
 Call 1-877-895-3665 (toll-free) or 215-273-8755 (collect)

Distributed by:
JOHNSON & JOHNSON CONSUMER INC.
 McNeil Consumer Healthcare Division
 Fort Washington, PA 19034 USA
 ©J&JCI 2015
www.motrin.com 00000000

Children's Motrin
 contains
 Ibuprofen Oral Suspension (NSAID)
 Bubble Gum Flavor

New Information: Read Entire Label
 NDC 50580-604-04

For Ages 2 to 11 Years

Children's Motrin
 contains Ibuprofen
 Oral Suspension (NSAID)
 100 mg per 5 mL
 Pain Reliever/Fever Reducer

Lasts up to 8 hours

Alcohol Free

Bubble Gum Flavor
 4 fl oz (120 mL)

Important: Read all product information before using. Keep this box for important information. This product is intended for use in children.

Drug Facts

Active ingredient (in each 5 mL)	Purpose
Ibuprofen 100 mg (NSAID)*	Pain reliever/fever reducer

*nonsteroidal anti-inflammatory drug

Uses temporarily:

- relieves minor aches and pains due to the common cold, flu, sore throat, headache and toothache
- reduces fever

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if your child:

- has had stomach ulcers or bleeding problems
- takes a blood thinning (anticoagulant) or steroid drug
- takes other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- takes more or for a longer time than directed

Sore throat warning: Severe or persistent sore throat or sore throat accompanied by high fever, headache, nausea, and vomiting may be serious. Consult doctor promptly. Do not use more than 2 days or administer to children under 3 years of age unless directed by doctor.

Do not use

- if the child has ever had an allergic reaction to ibuprofen or any other pain reliever/fever reducer
- right before or after heart surgery

3 0045-0604-04 0



NDC 50580-604-04

Important: Read all product information before using. Keep the box for important information. This product is intended for use in children.

Active ingredient (in each 5 mL): **Ibuprofen 100 mg (NSAID)**. **Purpose:** Pain reliever/fever reducer.

Warnings: Allergy alert: Ibuprofen may cause an allergic reaction, especially in people allergic to aspirin. Symptoms may include hives, facial swelling, asthma (wheezing), shock, weak or falling asleep, or dizziness. If an allergic reaction occurs, stop use and seek medical help right away. **Stomach bleeding warning:** This product contains an NSAID, which may cause stomach bleeding. The chance is higher if you or your child: has had stomach ulcers or bleeding problems; takes a blood-thinning (anticoagulant) or steroid drug; takes other drugs containing aspirin or other NSAIDs (aspirin, ibuprofen, naproxen, or others); or takes more or for a longer time than directed. **Some throat, fever, headache, nausea, and vomiting may be serious. Consult doctor promptly.** Do not use more than 2 days to relieve a fever or more than 3 days to relieve pain unless directed by doctor. **Do not use with other NSAIDs.** Do not use with other pain relievers, such as aspirin, acetaminophen, or other NSAIDs. **Do not use with other blood thinners, such as warfarin, unless directed by your doctor.** Do not use with other blood thinners, such as aspirin, acetaminophen, or other NSAIDs. **Do not use with other blood thinners, such as warfarin, unless directed by your doctor.**

Children's Motrin
 contains **Ibuprofen Oral Suspension (NSAID)**
100 mg per 5 mL 4 fl oz (120 mL)
Bubble Gum Flavor

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222). Directions: See important information about this product on the inside of this container. For adults and children 12 years of age and older, use weight to dose. For children under 12 years of age, use only dosing device. Do not use any other dosing device. If needed, repeat dose every 6-8 hours. Do not use more than 4 times a day. Use original bottle to maintain child's position.

Weight (lb)	Age (yr)	Dose (mL)*
under 24	under 2 years	ask a doctor
24-35 lbs	2-3 years	5 mL
36-47 lbs	4-5 years	7.5 mL
48-59 lbs	6-8 years	10 mL
60-71 lbs	9-10 years	12.5 mL
72-95 lbs	11 years	15 mL

*Or as directed by a doctor. Other info may be on the 5 mL container: sodium 2 mg; stores between 20-25°C (68-77°F); do not use if bottle wrap is printed "SAFETY SEAL" is broken or missing. Inactive ingredients: saccharin potassium, anhydrous citric acid, FD&C red no. 40, blue 1, glycerin, polyethylene glycol, purified water, sodium benzoate, sucrose, xanthan gum. Questions or comments? Call 1-877-895-1665 (10-14) or 215-223-4255 (collect). Distributed by: JOHNSON & JOHNSON CONSUMER INC., McNeil Consumer Healthcare Division, Fort Washington, PA, 19034 USA. ©J&JCI 2015. www.motrin.com 00000000

(b) (4)



(b) (4)

Important: Read all product information before using. Keep the box for important information. This product is intended for use in children.

Active ingredient (in each 5 mL): **Ibuprofen 100 mg (NSAID)**

Purpose: Pain reliever/fever reducer

Use: Temporarily relieve minor aches and pains due to the common cold, flu, sore throat, headache, and toothache in children.

Warnings:
 Allergy alert: Ibuprofen may cause an allergic reaction, especially in people allergic to aspirin. Symptoms may include hives, facial swelling, asthma (wheezing), shock, weakness, dizziness, or fainting. If an allergic reaction occurs, stop use and seek medical help right away. **Stomach bleeding warning:** This product contains an NSAID, which may cause serious stomach bleeding. The chance is higher if you or your child: has had stomach ulcers or bleeding problems; takes a blood thinner (an anticoagulant) or steroid drug; takes other drugs containing aspirin or other NSAIDs (aspirin, ibuprofen, naproxen, or others); or takes more or for a longer time than directed. **Some throat fever, headache, nausea, and vomiting may be serious. Consult doctor promptly.** Do not use more than 2 days to relieve fever or more than 3 days to relieve pain unless directed by doctor. **Do not use if the child has ever had an allergic reaction to ibuprofen or any other pain reliever/fever reducer, or if the child has heart disease, kidney disease, or a history of stomach problems, such as heartburn or indigestion, or if the child has had any problems or serious side effects from taking pain relievers or fever reducers.**

NDC 50580-601-20

Children's Motrin

contains **Ibuprofen**
Oral Suspension (NSAID)
100 mg per 5 mL 4 fl oz (120 mL)

Pain Reliever/Fever Reducer

Original Berry Flavor

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222). Directions on this product do not contain all restrictions or complete warnings for adults use. Do not give more than directed unless well advised by a doctor. Use only if directed on the label. If possible, use weight to dose. If weight is not given, use only the dosing chart on the label. Do not use any other dosing device. If needed, repeat dose every 6-8 hours. Do not use more than 4 times a day. Use the original bottle to maintain child's position.

Dosing Chart

Weight (lb)	Age (yr)	Dose (mL)*
under 24	under 2 years	ask a doctor
24-35 lbs	2-3 years	5 mL
36-47 lbs	4-5 years	7.5 mL
48-59 lbs	6-8 years	10 mL
60-71 lbs	9-10 years	12.5 mL
72-95 lbs	11 years	15 mL

*Or as directed by a doctor. Other information on each 5 mL contains sodium 2 mg. Store between 20° C (68°-77° F). Do not use if bottle cap is imprinted "SAFETY SEAL" is broken or missing. Inactive ingredients contain: benzalkonium chloride, hydroxyethylcellulose, polyethylene glycol, glycerin, polyethylene glycol, propylparaben, sodium benzoate, sucrose, xanthan gum. Questions or comments? Call 1-877-895-3665 (toll-free) or 215-273-8755 (local).
 Distributed by: JOHNSON & JOHNSON CONSUMER INC.
 McNeil Consumer Healthcare Division
 Fort Washington, PA 19034 USA ©J&JCI 2015 www.motrin.com 00000000

(b) (4)



Important: Read all product information before using. Keep the box for important information. This product is intended for use in children.

Active ingredient (in each 5 mL): **Ibuprofen 100 mg (NSAID)**

Purpose: Pain reliever/fever reducer

Use: Temporarily relieve minor aches and pains due to the common cold, flu, sore throat, headache, and toothache.

Warnings:
 Allergy alert: Ibuprofen may cause an allergic reaction, especially in people allergic to aspirin. Symptoms may include hives, facial swelling, asthma (wheezing), shock, weakness, dizziness, or fainting. If an allergic reaction occurs, stop use and seek medical help right away. **Stomach bleeding warning:** This product contains an NSAID, which may cause stomach bleeding. The chance is higher if you or your child: has had stomach ulcers or bleeding problems; takes a blood thinner (an anticoagulant) or steroid drug; takes another drug containing aspirin or other NSAIDs (aspirin, ibuprofen, naproxen, or others); or takes more or for a longer time than directed. **Some throat fever, headache, nausea, and vomiting may be serious. Consult doctor promptly.** Do not use more than 2 days to relieve fever or more than 3 days to relieve pain unless directed by doctor. **Do not use if the child has ever had an allergic reaction to ibuprofen or any other pain reliever/fever reducer, might be allergic to heart surgery, or if the child has a history of stomach problems, such as heartburn or if the child has problems or serious side effects from taking pain relievers or fever reducers.**

NDC 50588-184-50

Children's Motrin®

contains **Ibuprofen**
Oral Suspension (NSAID)
100 mg per 5 mL 4 fl oz (120 mL)

Pain Reliever/Fever Reducer

Non-Staining Dye-Free • Original Berry Flavor

Child has not been drinking fluids or child has lost a lot of fluid due to vomiting or diarrhea. Child has high blood pressure, heart disease, liver or kidney disease, or kidney disease. Child has asthma or is taking a diuretic. **Ask a doctor or pharmacist before use if the child is under 6 months of age or for any serious condition, or if longer than directed.** When using, this product may cause dizziness or lightheadedness. **Stop use and ask a doctor if the child experiences any of the following signs of stomach bleeding: feels faint, vomits blood, or has bloody or black stools.** **Stop use and ask a doctor if the child does not get any relief within 24 hours of treatment or if pain gets worse or lasts more than 3 days or if signs of allergy are present in the painful area.**


Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222). Directions in this product do not contain all restrictions or complete warnings for adults. Use only as directed. Do not use if child is under 6 months of age. Use only enclosed dosing cup. Do not use any other dosing device. If needed, repeat dose every 6-8 hours. Do not use more than 4 times a day in original bottle. Do not use if child is allergic to ibuprofen.

Dosing Chart

Weight (lb)	Age (yr)	Dose (mL)*
under 24	under 2 years	ask a doctor
24-35 lbs	2-3 years	5 mL
36-47 lbs	4-5 years	7.5 mL
48-59 lbs	6-8 years	10 mL
60-71 lbs	9-10 years	12.5 mL
72-95 lbs	11 years	15 mL

*Or as directed by a doctor. Other information in each 5 mL contains sodium 2 mg. Store between 20° (68-77°F). Do not use if bottle cap is imprinted "SAFETY SEAL" is broken or missing. Inactive ingredients: saccharin, sodium benzoate, sodium citrate, xanthan gum, polyethylene glycol, purified water, sodium benzoate, sucrose, xanthan gum. Questions or comments? Call 1-877-993-3665 (toll-free) or 215-271-8755 (local).
 Distributed by: **JOHNSON & JOHNSON CONSUMER INC.**
 McNeil Consumer Healthcare Division
 Fort Washington, PA 19034 USA ©J&JCI 2015 www.motrin.com
 00000000

(b) (4)



Drug Facts (continued)

Ask a doctor before use if

- stomach bleeding warning applies to your child
- child has a history of stomach problems, such as heartburn
- child has problems or serious side effects from taking pain relievers or fever reducers
- child has not been drinking fluids
- child has lost a lot of fluid due to vomiting or diarrhea
- child has high blood pressure, heart disease, liver cirrhosis, or kidney disease
- child has asthma
- child is taking a diuretic

Ask a doctor or pharmacist before use if the child is

- under a doctor's care for any serious condition
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs
- the risk of heart attack or stroke may increase if you use more than directed or for longer than directed

Stop use and ask a doctor if

- child experiences any of the following signs of stomach bleeding
 - feels faint
 - vomits blood
 - has bloody or black stools
- has stomach pain that does not get better
- the child does not get any relief within first day (24 hours) of treatment
- fever or pain gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions

- this product does not contain directions or complete warnings for adult use
- do not give more than directed
- shake well before using
- mL = milliliter
- find right dose on chart. If possible, use weight to dose; otherwise use age.
- use only enclosed dosing cup. Do not use any other dosing device.

Drug Facts (continued)

- if needed, repeat dose every 6-8 hours
- do not use more than 4 times a day
- replace original bottle cap to maintain child resistance

Dosing Chart

Weight (lb)	Age (yr)	Dose (mL)*
under 24	under 2 years	ask a doctor
24-35 lbs	2-3 years	5 mL
36-47 lbs	4-5 years	7.5 mL
48-59 lbs	6-8 years	10 mL
60-71 lbs	9-10 years	12.5 mL
72-95 lbs	11 years	15 mL

* or as directed by a doctor

Other information

- each 5 mL contains: sodium 2 mg
- store between 20-25° C (68-77°F)
- do not use if carton is opened or if carton tape or bottle wrap imprinted "SAFETY SEAL" is broken or missing

Inactive ingredients acesulfame potassium, anhydrous citric acid, D&C yellow no. 10, FD&C red no. 40, flavors, glycerin, polysorbate 80, pregelatinized starch, purified water, sodium benzoate, sucrose, xanthan gum

Questions or comments?
 Call 1-877-895-3665 (toll-free) or 215-273-8755 (collect)

Distributed by:
JOHNSON & JOHNSON CONSUMER INC.
 McNeil Consumer Healthcare Division
 Fort Washington, PA 19034 USA
 ©J&JCI 2015
www.motrin.com 00000000

FOR HOSPITAL/GOVERNMENT USE ONLY
 For Ages 2 to 11 Years

Children's Motrin®
 contains
Ibuprofen Oral Suspension (NSAID)
 Pain Reliever / Fever Reducer
 Original Berry Flavor

New Information: Read Entire Label
 NDC 50580-601-21

FOR HOSPITAL/GOVERNMENT USE ONLY
 For Ages 2 to 11 Years

Children's Motrin®
 contains *Ibuprofen Oral Suspension (NSAID)*
 100 mg per 5 mL

Pain Reliever / Fever Reducer

Original Berry Flavor

Lasts up to 8 hours

Alcohol Free

Not for Retail Sale

4 fl oz (120 mL)

Important: Read all product information before using. Keep this box for important information. This product is intended for use in children.

Drug Facts

Active ingredient (in each 5 mL)	Purpose
Ibuprofen 100 mg (NSAID)*	Pain reliever/fever reducer

*nonsteroidal anti-inflammatory drug

Uses temporarily:

- relieves minor aches and pains due to the common cold, flu, sore throat, headache and toothache
- reduces fever

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if your child:

- has had stomach ulcers or bleeding problems
- takes a blood thinning (anticoagulant) or steroid drug
- takes other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- takes more or for a longer time than directed

Sore throat warning: Severe or persistent sore throat or sore throat accompanied by high fever, headache, nausea, and vomiting may be serious. Consult doctor promptly. Do not use more than 2 days or administer to children under 3 years of age unless directed by doctor.

Do not use

- if the child has ever had an allergic reaction to ibuprofen or any other pain reliever/fever reducer
- right before or after heart surgery



Important: Read all product information before using. Keep the box for important information. This product is intended for use in children.

Active ingredient (in each 5 mL): **Ibuprofen 100 mg (NSAID)**

Purpose: Pain reliever/fever reducer

Use: Temporarily relieve minor aches and pains due to the common cold, flu, sore throat, headache, and toothache in children.

Warnings:
 Allergy alert: Ibuprofen may cause an allergic reaction, especially in people allergic to aspirin. Symptoms may include hives, facial swelling, asthma (wheezing), shock, weakness, dizziness, or fainting. If an allergic reaction occurs, stop use and seek medical help right away. **Stomach bleeding warning:** This product contains an NSAID, which may cause serious stomach bleeding. The chance is higher if your child: has had stomach ulcers or bleeding problems; takes a blood thinner (an anticoagulant) or steroid drug; takes other drugs containing aspirin or other NSAIDs (aspirin, ibuprofen, naproxen, or others); or takes more or for a longer time than directed. **Some throat fever, headache, nausea, and vomiting may be serious. Consult doctor promptly.** Do not use more than 2 days to administer to a child under 3 years of age unless directed by doctor. **Do not use with:** If child has ever had an allergic reaction to ibuprofen or any other pain reliever/fever reducer, might be sensitive to heart surgery. **Ask a doctor before use if:** child has a history of stomach problems, such as heartburn or stomach ulcers; child has a history of stomach problems, such as heartburn or stomach ulcers; child has a history of stomach problems or serious side effects from taking pain relievers or fever reducers.

FOR HOSPITAL/GOVERNMENT USE ONLY

Net for Retail Sale NDC 58580-601-21

Children's Motrin

contains **Ibuprofen Oral Suspension (NSAID)**
100 mg per 5 mL

Pain Reliever/Fever Reducer 4 fl oz (120 mL)

Original Berry Flavor

Child has not been drinking fluids; child has lost a lot of fluid due to vomiting or diarrhea; child has high blood pressure, heart disease, liver or kidney disease; child has asthma; child is taking a diuretic. **Ask a doctor or pharmacist before use if the child is under 6 months of age for any serious condition; taking any other drug; when using, child should not take with food or milk; if stomach upset occurs, heart attack or stroke may occur; if you use more than directed or for longer than directed; stop use and ask a doctor if child experiences any of the following signs of stomach bleeding: feels faint, vomit or blood, has bloody or black stools, or has stomach pain that does not get better when the child does not get any relief within 24 hours of treatment while or pain gets worse or lasts more than 3 days; or signs or symptoms present in the painful area, or any**

new symptoms appear. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. (1) 800-222-1222. Directions in this product does not contain all essential or complete warnings for adults use. **Do not give more than directed.** Shake well before using. 5 mL (1 mL for children) dose on label below. If possible, use weight to dose; otherwise use age. **Use only enclosed dosing cup.** Do not use any other dosing device. If needed, repeat dose every 6-8 hours; do not use more than 4 times a day. **Cap and original bottle up to maintain child's safety.**

Dosing Chart

Weight (lb)	Age (yr)	Dose (mL)*
under 24	under 2 years	ask a doctor
24-35 lbs	2-3 years	5 mL
36-47 lbs	4-5 years	7.5 mL
48-59 lbs	6-8 years	10 mL
60-71 lbs	9-10 years	12.5 mL
72-95 lbs	11 years	15 mL

*Or as directed by a doctor. Other information: each 5 mL contains sodium 2 mg. Store between 20°C (68°F) and 25°C (77°F). Do not use if bottle cap is imprinted "SAFETY SEAL" is broken or missing. Inactive ingredients: sodium benzoate, anhydrous citric acid, D&C yellow no. 10, FD&C white no. 40, lactose, glycerin, polyethylene glycol, propylparaben, purified water, sodium benzoate, sucrose, xanthan gum. Questions or comments? Call 1-877-895-3665 (toll-free) or 215-273-8755 (local).
 Distributed by: **JOHNSON & JOHNSON CONSUMER INC.**
 McNeil Consumer Healthcare Division
 Fort Washington, PA 19034 USA ©J&JCI 2015 www.motrin.com
 00000000

(b) (4)

Trim Safe Area Bleed Score

w 3.875"

Convenient 2-PACK!

NDC 50580-601-25 NDC 50580-601-25

For Ages 2 to 11 Years *For Ages 2 to 11 Years*

Children's Motrin® **Children's Motrin®**

contains ibuprofen
Oral Suspension (NSAID)
100 mg per 5 mL

contains ibuprofen
Oral Suspension (NSAID)
100 mg per 5 mL

Pain Reliever/Fever Reducer **Pain Reliever/Fever Reducer**

Lasts up to 8 hours **Lasts up to 8 hours**

Alcohol Free **Alcohol Free**

Original Berry Flavor *Original Berry Flavor*

h 5.65625"

Distributed by:
JOHNSON & JOHNSON CONSUMER INC.
McNeil Consumer Healthcare Division
Fort Washington, PA 19034 USA
©J&JCI 2015 www.motrin.com
McN-XX-0000



3 0045-0192-08 0

Score 0.65625" from bottom

(b) (4)



This is a representation of an electronic record that was signed electronically and this page is the manifestation of the electronic signature.

/s/

VALERIE S PRATT
06/16/2016