



(b) (4)



**Important: Read all product information before using. Keep the box for important information. This product is intended for use in children.**

**Active ingredient (in each 5 mL):** ibuprofen (NSAID) **Purpose:** Pain reliever/fever reducer

**Uses:** Temporarily relieves minor aches and pains due to the common cold, flu, sore throat, headache and toothache. Reduces fever.

**Warnings:** Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: hives; facial swelling; as hives (wheezing); shock; skin reddening; rash; itching. If an allergic reaction occurs, stop use and seek medical help right away. **Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: have had stomach ulcers or bleeding problems; take a blood-thinning (anti-coagulant) or steroid drug; take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or rofecoxib); take more or for a longer time than directed. **Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed. **Severe or persistent sore throat or sore throat** accompanied by high fever, headache, nausea, and vomiting may be serious. Consult your doctor promptly. Do not use more than 2 days or administer to children under 3 years of age unless directed by doctor. **Do not use if:** the child has ever had an allergic reaction to ibuprofen or any other pain reliever/fever reducer; right before or after heart surgery. **Ask a doctor before use if:** stomach bleeding warning applies to your child; child has a history of stomach problems, such as heartburn; child has problems or serious side effects from taking pain relievers or fever reducers; child

NDC 50580-601-20

# Children's Motrin

contains **Ibuprofen**  
**Oral Suspension (NSAID)**  
**100 mg per 5 mL** 4 fl oz (120 mL)

**Pain Reliever/Fever Reducer**

**Original Berry Flavor**

the pain of area. Any new symptoms appear. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. (800) 222-1222.

**Directions:** This product does not contain dextrose or complete warnings for adult use. Do not give more than directed. Shake well before using. Measure with the provided measuring device. If possible, use weight to dose or otherwise use age. Use only enclosed dosing cup. Do not use any other dosing device. If needed, repeat dose every 6-8 hours. Do not use more than 4 times a day. Replace original bottle cap to maintain child resistance.

**Dosing Chart**

Weight (lb)	Age (yr)	Dose (mL)*
under 24	under 2 years	ask a doctor
24-35 lbs	2-3 years	5 mL
36-47 lbs	4-5 years	7.5 mL
48-59 lbs	6-8 years	10 mL
60-71 lbs	9-10 years	12.5 mL
72-95 lbs	11 years	15 mL

\*or as directed by a doctor

**Other information:** Each 5 mL contains: sodium 2 mg; store between 20-25°C (68-77°F). **Do not use if bottle wrap imprinted "SAFETY SEAL" is broken or missing.**

**Inactive ingredients:** saccharin, potassium, anhydrous citric acid, D&C yellow no. 10, FD&C red no. 40, flavo-s, glycol polybase 80, pregelatinized starch, purified water, sodium benzoate, sucrose, xanthan gum. **Questions or comments?** Call 1-877-895-3665 (toll-free) or 215-273-8755 (no text).

Distributed by: **JOHNSON & JOHNSON CONSUMER INC.**  
McKee Consumer Healthcare Division  
Fort Washington, PA 19034 USA ©J&JCI 2016 [www.motrin.com](http://www.motrin.com)  
00000000

(b) (4)

# Children's Motrin<sup>®</sup>

contains  
**Ibuprofen Oral Suspension (NSAID)**

## Original Berry Flavor

---

**Drug Facts (continued)**

**Ask a doctor before use if**

- stomach bleeding warning applies to your child
- child has a history of stomach problems, such as heartburn
- child has problems or serious side effects from taking pain relievers or fever reducers
- child has not been drinking fluids
- child has lost a lot of fluid due to vomiting or diarrhea
- child has high blood pressure, heart disease, liver cirrhosis, kidney disease, or had a stroke
- child has asthma
- child is taking a diuretic

**Ask a doctor or pharmacist before use if the child is**

- under a doctor's care for any serious condition
- taking any other drug

**When using this product**

- take with food or milk if stomach upset occurs

**Stop use and ask a doctor if**

- child experiences any of the following signs of stomach bleeding:
  - feels faint
  - vomits blood
  - has bloody or black stools
- has stomach pain that does not get better
- child has symptoms of heart problems or stroke:
  - chest pain
  - trouble breathing
  - weakness in one part or side of body
  - slurred speech
  - leg swelling
- the child does not get any relief within first day (24 hours) of treatment
- fever or pain gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

**Directions**

- this product does not contain directions or complete warnings for adult use
- do not give more than directed

**Drug Facts (continued)**

- shake well before using
- mL = milliliter
- find right dose on chart. If possible, use weight to dose; otherwise use age.
- use only enclosed dosing cup. Do not use any other dosing device.
- if needed, repeat dose every **6-8 hours**
- do not use more than **4 times a day**
- replace original bottle cap to maintain child resistance

**Dosing Chart**

Weight (lb)	Age (yr)	Dose (mL)*
under 24	under 2 years	ask a doctor
24-35 lbs	2-3 years	5 mL
36-47 lbs	4-5 years	7.5 mL
48-59 lbs	6-8 years	10 mL
60-71 lbs	9-10 years	12.5 mL
72-95 lbs	11 years	15 mL

\* or as directed by a doctor

**Other information**

- each 5 mL contains: sodium 2 mg
- store between 20-25° C (68-77° F)
- do not use if carton is opened or if carton tape or bottle wrap imprinted "SAFETY SEAL" is broken or missing

**Inactive ingredients** acesulfame potassium, anhydrous citric acid, D&C yellow no. 10, FD&C red no. 40, flavors, glycerin, polysorbate 80, pregelatinized starch, purified water, sodium benzoate, sucrose, xanthan gum

**Questions or comments?**  
Call **1-877-895-3665** (toll-free) or **215-273-8755** (collect)

Distributed by:  
**JOHNSON & JOHNSON CONSUMER INC.**  
McNeil Consumer Healthcare Division  
Fort Washington, PA 19034 USA  
©J&JCI 2016  
[www.motrin.com](http://www.motrin.com) 00000000

**Important: Read all product information before using. Keep this box for important information. This product is intended for use in children.**

**Drug Facts**

Active ingredient (in each 5 mL)	Purpose
ibuprofen 100 mg (NSAID)*.....Pain reliever/fever reducer	
*nonsteroidal anti-inflammatory drug	

**Uses temporarily:**

- relieves minor aches and pains due to the common cold, flu, sore throat, headache and toothache
- reduces fever

**Warnings**

**Allergy alert:** ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- asthma (wheezing)
- skin reddening
- facial swelling
- shock
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if your child:

- has had stomach ulcers or bleeding problems
- takes a blood thinning (anticoagulant) or site old drug
- takes other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- takes more or for a longer time than directed

**Heart attack and stroke warning** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

**Sore throat warning:** Severe or persistent sore throat or sore throat accompanied by high fever, headache, nausea, and vomiting may be serious. Consult doctor promptly. Do not use more than 2 days or administer to children under 3 years of age unless directed by doctor.

**Do not use**

- if the child has ever had an allergic reaction to ibuprofen or any other pain reliever/fever reducer
- right before or after heart surgery

*See New Warnings*  
NDC 50580-601-20

**For Ages 2 to 11 Years**

# Children's Motrin<sup>®</sup>

contains **Ibuprofen Oral Suspension (NSAID)**  
**100 mg per 5 mL**

**Pain Reliever /Fever Reducer**

**Lasts up to 8 hours**

**Alcohol Free**

**Original Berry Flavor**

**4 fl oz (120 mL)**

3 0045- 0192- 04 2

LOT:  
EXP:





**Children's Motrin**  
**DYE - FREE**  
contains *Ibuprofen Oral Suspension (NSAID)*  
Non-Staining  
*Original Berry Flavor*

**Drug Facts (continued)**

**Ask a doctor before use if**

- stomach bleeding warning applies to your child
- child has a history of stomach problems, such as heartburn
- child has problems or serious side effects from taking pain relievers or fever reducers
- child has not been drinking fluids
- child has lost a lot of fluid due to vomiting or diarrhea
- child has high blood pressure, heart disease, liver cirrhosis, kidney disease, or had a stroke
- child has asthma
- child is taking a diuretic

**Ask a doctor or pharmacist before use if the child is**

- under a doctor's care for any serious condition
- taking any other drug

**When using this product**

- take with food or milk if stomach upset occurs

**Stop use and ask a doctor if**

- child experiences any of the following signs of stomach bleeding:
  - feels faint
  - vomits blood
  - has bloody or black stools
- has stomach pain that does not get better
- child has symptoms of heart problems or stroke:
  - chest pain
  - trouble breathing
  - weakness in one part or side of body
  - slurred speech
  - leg swelling
- the child does not get any relief within first day (24 hours) of treatment
- fever or pain gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

**Directions**

- this product does not contain directions or complete warnings for adult use
- do not give more than directed

**Drug Facts (continued)**

- shake well before using
- mL = milliliter
- find right dose on chart. If possible, use weight to dose; otherwise use age.
- use only enclosed dosing cup. Do not use any other dosing device.
- if needed, repeat dose every **6-8 hours**
- do not use more than **4 times a day**
- replace original bottle cap to maintain child resistance

**Dosing Chart**

Weight (lb)	Age (yr)	Dose (mL)*
under 24	under 2 years	ask a doctor
24-35 lbs	2-3 years	5 mL
36-47 lbs	4-5 years	7.5 mL
48-59 lbs	6-8 years	10 mL
60-71 lbs	9-10 years	12.5 mL
72-95 lbs	11 years	15 mL

\* or as directed by a doctor

**Other information**

- each 5 mL contains: sodium 2 mg
- store between 20-25°C (68-77°F)
- do not use if carton is opened or if carton tape or bottle wrap imprinted "SAFETY SEAL®" is broken or missing

**Inactive ingredients** acesulfame potassium, anhydrous citric acid, flavors, glycerin, polysorbate 80, pregelatinized starch, purified water, sodium benzoate, sucrose, xanthan gum

**Questions or comments?**  
Call 1-877-895-3665 (toll-free) or 215-273-8755 (collect)

Distributed by:  
**JOHNSON & JOHNSON CONSUMER INC.**  
McNeil Consumer Healthcare Division  
Fort Washington, PA 19034 USA  
© J&JCI 2016  
[www.motrin.com](http://www.motrin.com) 00000000

*See New Warnings*  
NDC 50580-184-50

**For Ages 2 to 11 Years**



**Children's Motrin**  
contains *Ibuprofen Oral Suspension (NSAID)*  
**100 mg per 5 mL**  
**Pain Reliever / Fever Reducer**

**DYE - FREE**

**Lasts up to 8 hours**

**Alcohol Free**

*Non-Staining Original Berry Flavor*

**4 fl oz (120 mL)**

3 0045-0184-04 7

LOT:  
EXP:

**Important: Read all product information before using. Keep this box for important information. This product is intended for use in children.**

**Drug Facts**

Active ingredient (in each 5 mL)	Purpose
ibuprofen 100 mg (NSAID)*.....Pain reliever/fever reducer	
*nonsteroidal anti-inflammatory drug	

**Uses** temporarily:

- relieves minor aches and pains due to the common cold, flu, sore throat, headache and toothache
- reduces fever

**Warnings**

**Allergy alert:** ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- asthma (wheezing)
- skin reddening
- facial swelling
- shock
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if your child:

- has had stomach ulcers or bleeding problems
- takes a blood thinning (anticoagulant) or steroid drug
- takes other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- takes more or for a longer time than directed

**Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

**Sore throat warning:** Severe or persistent sore throat or sore throat accompanied by high fever, headache, nausea, and vomiting may be serious. Consult doctor promptly. Do not use more than 2 days or administer to children under 3 years of age unless directed by doctor.

**Do not use**

- if the child has ever had an allergic reaction to ibuprofen or any other pain reliever/fever reducer
- right before or after heart surgery

NDC 50580-604-04

**Important: Read all product information before using. Keep the box for important information. This product is intended for use in children.**

**Active ingredient (in each 5 mL):** ibuprofen (NSAID) 100 mg

**Purpose:** Pain reliever/fever reducer

**Warnings:** Do not use if your child has had stomach ulcers or bleeding, or if they are taking blood thinning (anti-coagulant) or steroid drugs. Do not use if your child has had a heart attack or stroke, or if they have a history of stomach problems, such as heartburn. Do not use if your child has had a severe allergic reaction to ibuprofen or any other pain reliever/fever reducer. Do not use if your child has had a severe allergic reaction to any other NSAID. Do not use if your child has had a severe allergic reaction to any other NSAID. Do not use if your child has had a severe allergic reaction to any other NSAID.

**Directions:** See the enclosed dosing information. Do not use any other dosing device. If needed, repeat dose every 6-8 hours. Do not use more than 4 times a day. Replace original bottle cap to maintain child resistance.

**Other information:** Each 5 mL contains sodium 2 mg. Store between 20-25°C (68-77°F). Do not use if bottle wrap imprinted "SAFETY SEAL" is broken or missing. Inactive ingredients: acacia, croscarmellose sodium, hydroxypropyl methylcellulose, polyethylene glycol 400, polyethylene glycol 600, polyethylene glycol 80, pregelatinized starch, purified water, sodium benzoate, sucrose, xanthan gum.

**Questions or comments?** Call 1-877-895-3665 (toll-free) or 215-273-8755 (collect).

**Distributed by:** JOHNSON & JOHNSON CONSUMER INC.  
New Jersey Consumer Healthcare Division  
Fort Washington, PA 19034 USA © J&JCI 2016

**www.motrin.com**  
00000000

**Children's Motrin**  
contains **Ibuprofen Oral Suspension (NSAID)**  
**100 mg per 5 mL** 4 fl oz (120 mL)  
**Pain Reliever/Fever Reducer**  
**Bubble Gum Flavor**

**Warnings:** Do not use if your child has had stomach ulcers or bleeding, or if they are taking blood thinning (anti-coagulant) or steroid drugs. Do not use if your child has had a heart attack or stroke, or if they have a history of stomach problems, such as heartburn. Do not use if your child has had a severe allergic reaction to ibuprofen or any other pain reliever/fever reducer. Do not use if your child has had a severe allergic reaction to any other NSAID. Do not use if your child has had a severe allergic reaction to any other NSAID.

**Directions:** See the enclosed dosing information. Do not use any other dosing device. If needed, repeat dose every 6-8 hours. Do not use more than 4 times a day. Replace original bottle cap to maintain child resistance.

**Other information:** Each 5 mL contains sodium 2 mg. Store between 20-25°C (68-77°F). Do not use if bottle wrap imprinted "SAFETY SEAL" is broken or missing. Inactive ingredients: acacia, croscarmellose sodium, hydroxypropyl methylcellulose, polyethylene glycol 400, polyethylene glycol 600, polyethylene glycol 80, pregelatinized starch, purified water, sodium benzoate, sucrose, xanthan gum.

**Questions or comments?** Call 1-877-895-3665 (toll-free) or 215-273-8755 (collect).

**Distributed by:** JOHNSON & JOHNSON CONSUMER INC.  
New Jersey Consumer Healthcare Division  
Fort Washington, PA 19034 USA © J&JCI 2016

**www.motrin.com**  
00000000

(b) (4)

# Children's Motrin®

contains  
**Ibuprofen Oral Suspension (NSAID)**

## Bubble Gum Flavor

**Drug Facts (continued)**

**Ask a doctor before use if**

- stomach bleeding warning applies to your child
- child has a history of stomach problems, such as heartburn
- child has problems or serious side effects from taking pain relievers or fever reducers
- child has not been drinking fluids
- child has lost a lot of fluid due to vomiting or diarrhea
- child has high blood pressure, heart disease, liver cirrhosis, kidney disease, or had a stroke
- child has asthma
- child is taking a diuretic

**Ask a doctor or pharmacist before use if the child is**

- under a doctor's care for any serious condition
- taking any other drug

**When using this product**

- take with food or milk if stomach upset occurs

**Stop use and ask a doctor if**

- child experiences any of the following signs of stomach bleeding:
  - feels faint
  - vomits blood
  - has bloody or black stools
- has stomach pain that does not get better
- child has symptoms of heart problems or stroke:
  - chest pain
  - trouble breathing
  - weakness in one part or side of body
  - slurred speech
  - leg swelling
- the child does not get any relief within first day (24 hours) of treatment
- fever or pain gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

**Directions**

- this product does not contain directions or complete warnings for adult use
- do not give more than directed →

**Drug Facts (continued)**

- shake well before using
- mL = milliliter
- find right dose on chart. If possible, use weight to dose; otherwise use age.
- use only enclosed dosing cup. Do not use any other dosing device.
- if needed, repeat dose every **6-8 hours**
- do not use more than **4 times a day**
- replace original bottle cap to maintain child resistance

**Dosing Chart**

Weight (lb)	Age (yr)	Dose (mL)*
under 24	under 2 years	ask a doctor
24-35 lbs	2-3 years	5 mL
36-47 lbs	4-5 years	7.5 mL
48-59 lbs	6-8 years	10 mL
60-71 lbs	9-10 years	12.5 mL
72-95 lbs	11 years	15 mL

\* or as directed by a doctor

**Other information**

- each 5 mL contains: sodium 2 mg
- store between 20-25° C (68-77° F)
- do not use if carton is opened or if carton tape or bottle wrap imprinted "SAFETY SEAL®" is broken or missing

**Inactive ingredients** acesulfame potassium, anhydrous citric acid, FD&C red no. 40, flavors, glycerin, polysorbate 80, pregelatinized starch, purified water, sodium benzoate, sucrose, xanthan gum

**Questions or comments?**  
Call **1-877-895-3665** (toll-free) or **215-273-8755** (collect)

Distributed by:  
**JOHNSON & JOHNSON CONSUMER INC.**  
McNeil Consumer Healthcare Division  
Fort Washington, PA 19034 USA  
©J&JCI 2016  
[www.motrin.com](http://www.motrin.com) 00000000

**See New Warnings**

NDC 50580-604-04

**For Ages 2 to 11 Years**

# Children's Motrin®

contains **Ibuprofen Oral Suspension (NSAID)**  
**100 mg per 5 mL**

**Pain Reliever / Fever Reducer**

**Lasts up to 8 hours**

**Alcohol Free**



**Bubble Gum Flavor**

**4 fl oz (120 mL)**



3 0045-0604-408



LOT: \_\_\_\_\_

EXP: \_\_\_\_\_

Important: Read all product information before using. Keep this box for important information. This product is intended for use in children.

**Drug Facts**

Active ingredient (in each 5 mL)	Purpose
bupropion 100 mg (NSAID)*.....Pain reliever/fever reducer	
*nonsteroidal anti-inflammatory drug	

**Uses temporarily:**

- relieves minor aches and pains due to the common cold, flu, sore throat, headache and toothache
- reduces fever

**Warnings**

**Allergy alert:** bupropion may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- asthma (wheezing)
- skin reddening
- facial swelling
- shock
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if your child:

- has had stomach ulcers or bleeding problems
- takes a blood thinning (anticoagulant) or antiplatelet drug
- takes other drugs containing prescription or nonprescription NSAIDs (aspirin, bupropion, naproxen, or others)
- takes more or for a longer time than directed

**Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

**Sore throat warning:** Severe or persistent sore throat or sore throat accompanied by high fever, headache, nausea, and vomiting may be serious. Consult doctor promptly. Do not use more than 2 days or administer to children under 3 years of age unless directed by doctor.

**Do not use**

- if the child has ever had an allergic reaction to bupropion or any other pain reliever/fever reducer
- right before or after heart surgery →

**Important: Read all product information before using. Keep the box for important information. This product is intended for use in children.**

**Active ingredient (in each 5 mL):** ibuprofen (NSAID)  
Purpose: Pain reliever/fever reducer

**Warnings:** Allergic alert: ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: hives, facial swelling, rash, wheezing, shock, skin reddening, dizziness, fainting, or loss of consciousness. If an allergic reaction occurs, stop use and seek medical help right away. **Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you have had stomach ulcers or bleeding problems, take blood thinning (anti-coagulant) or steroid drugs, or take aspirin, ibuprofen, or other NSAIDs (aspirin, ibuprofen, naproxen, or rofecoxib) for a longer time than directed. **Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed. **Severe or persistent sore throat or sore mouth:** accompanied by high fever, headache, nausea, and vomiting may be serious. Consult your doctor promptly. Do not use more than 2 days or administer to children under 3 years of age unless directed by doctor. **Do not use** if the child has ever had an allergic reaction to ibuprofen or any other pain reliever/fever reducer, right before or after heart surgery. **Ask a doctor before use if** stomach bleeding warning applies to your child, if the child has a history of stomach problems, such as heartburn, or if the child has had a serious side effect from taking pain relievers or fever reducers.

NDC 50580-603-04

# Children's Motrin

contains **Ibuprofen Oral Suspension (NSAID)**  
**100 mg per 5 mL**      4 fl oz (120 mL)

**Pain Reliever/Fever Reducer**

**Grape Flavor**

Keep out of reach of children. If any new symptoms appear, keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (800) 222-1222.

**Directions:** This product does not contain dextrose or complete warnings for adult use. Do not give to someone who is allergic to aspirin. Shake well before using. Measure with the provided measuring device. If possible, use weight to dose; otherwise use age. Use only enclosed dosing cup. Do not use any other dosing device. If needed, repeat dose every 6-8 hours. Do not use more than 4 times a day. Replace original bottle cap to maintain child resistance.

**Dosing Chart**

Weight (lb)	Age (yr)	Dose (mL)*
under 24	under 2 years	ask a doctor
24-35 lbs	2-3 years	5 mL
36-47 lbs	4-5 years	7.5 mL
48-59 lbs	6-8 years	10 mL
60-71 lbs	9-10 years	12.5 mL
72-95 lbs	11 years	15 mL

\* or as directed by a doctor

**Other information:** Each 5 mL contains sodium 2 mg. Store between 20-25°C (68-77°F). Do not use if bottle wrap imprinted SAFETY SEAL® is broken or missing. Inactive ingredients: acetylsalicylic acid, potassium, anhydrous citric acid, D&C red no. 33, FD&C blue no. 1, FD&C red no. 40, iron oxide yellow, polyethylene glycol 400, pregelatinized starch, purified water, sodium benzoate, sucrose, xanthan gum. Questions or comments? Call 1-877-895-3665 (toll-free) or 215-275-8755 (direct).

Distributed by **JOHNSON & JOHNSON CONSUMER INC.**  
Medical Consumer Healthcare Division  
Ft. Washington, PA 19034 USA ©J&J 2016 [www.motrin.com](http://www.motrin.com)  
00000000

(b) (4)

# Children's Motrin®

contains  
**Ibuprofen Oral Suspension (NSAID)**

## Grape Flavor

**Drug Facts (continued)**

**Ask a doctor before use if**

- stomach bleeding warning applies to your child
- child has a history of stomach problems, such as heartburn
- child has problems or serious side effects from taking pain relievers or fever reducers
- child has not been drinking fluids
- child has lost a lot of fluid due to vomiting or diarrhea
- child has high blood pressure, heart disease, liver cirrhosis, kidney disease, or had a stroke
- child has asthma
- child is taking a diuretic

**Ask a doctor or pharmacist before use if the child is**

- under a doctor's care for any serious condition
- taking any other drug

**When using this product**

- take with food or milk if stomach upset occurs

**Stop use and ask a doctor if**

- child experiences any of the following signs of stomach bleeding:
  - feels faint
  - vomits blood
  - has bloody or black stools
- has stomach pain that does not get better
- child has symptoms of heart problems or stroke:
  - chest pain
  - trouble breathing
  - weakness in one part or side of body
  - slurred speech
  - leg swelling
- the child does not get any relief within first day (24 hours) of treatment
- fever or pain gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

**Directions**

- this product does not contain directions or complete warnings for adult use
- do not give more than directed

**Drug Facts (continued)**

- shake well before using
- mL = milliliter
- find right dose on chart. If possible, use weight to dose; otherwise use age.
- use only enclosed dosing cup. Do not use any other dosing device.
- if needed, repeat dose every **6-8 hours**
- do not use more than **4 times a day**
- replace original bottle cap to maintain child resistance

**Dosing Chart**

Weight (lb)	Age (yr)	Dose (mL)*
under 24	under 2 years	ask a doctor
24-35 lbs	2-3 years	5 mL
36-47 lbs	4-5 years	7.5 mL
48-59 lbs	6-8 years	10 mL
60-71 lbs	9-10 years	12.5 mL
72-95 lbs	11 years	15 mL

\* or as directed by a doctor

**Other information**

- each 5 mL contains: sodium 2 mg
- store between 20-25° C (68-77° F)
- do not use if carton is opened or if carton tape or bottle wrap imprinted "SAFETY SEAL®" is broken or missing

**Inactive ingredients** acesulfame potassium, anhydrous citric acid, D&C red no. 33, FD&C blue no. 1, FD&C red no. 40, flavors, glycerin, polysorbate 80, pregelatinized starch, purified water, sodium benzoate, sucrose, xanthan gum

**Questions or comments?**  
Call **1-877-895-3665** (toll-free) or **215-273-8755** (collect)

Distributed by:  
**JOHNSON & JOHNSON CONSUMER INC.**  
McNeil Consumer Healthcare Division  
Fort Washington, PA 19034 USA  
©J&JCI 2016  
[www.motrin.com](http://www.motrin.com) 00000000

*See New Warnings*  
NDC 50580-603-04

**For Ages 2 to 11 Years**

# Children's Motrin®

contains **Ibuprofen Oral Suspension (NSAID)**  
**100 mg per 5 mL**

**Pain Reliever / Fever Reducer**

**Lasts up to 8 hours**

**Alcohol Free**



**Grape Flavor**  
**4 fl oz (120 mL)**

**Important: Read all product information before using. Keep this box for important information. This product is intended for use in children.**

Drug Facts	Purpose
<p><b>Active ingredient (in each 5 mL)</b></p> <p>ibuprofen 100 mg (NSAID)*.....Pain reliever/fever reducer</p> <p>*nonsteroidal anti-inflammatory drug</p>	
<p><b>Uses temporarily:</b></p> <ul style="list-style-type: none"> <li>■ relieves minor aches and pains due to the common cold, flu, sore throat, headache and toothache</li> <li>■ reduces fever</li> </ul>	
<p><b>Warnings</b></p> <p><b>Allergy alert:</b> ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:</p> <ul style="list-style-type: none"> <li style="width: 50%;">■ hives</li> <li style="width: 50%;">■ facial swelling</li> <li style="width: 50%;">■ asthma (wheezing)</li> <li style="width: 50%;">■ shock</li> <li style="width: 50%;">■ skin reddening</li> <li style="width: 50%;">■ rash</li> <li style="width: 50%;">■ blisters</li> </ul> <p>If an allergic reaction occurs, stop use and seek medical help right away.</p> <p><b>Stomach bleeding warning:</b> This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if your child:</p> <ul style="list-style-type: none"> <li>■ has had stomach ulcers or bleeding problems</li> <li>■ takes a blood thinning (anticoagulant) or antiplatelet drug</li> <li>■ takes other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)</li> <li>■ takes more or for a longer time than directed</li> </ul> <p><b>Heart attack and stroke warning:</b> NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.</p> <p><b>Sore throat warning:</b> Severe or persistent sore throat or sore throat accompanied by high fever, headache, nausea, and vomiting may be serious. Consult doctor promptly. Do not use more than 2 days or administer to children under 3 years of age unless directed by doctor.</p>	
<p><b>Do not use</b></p> <ul style="list-style-type: none"> <li>■ if the child has ever had an allergic reaction to ibuprofen or any other pain reliever/fever reducer</li> <li>■ right before or after heart surgery</li> </ul>	



3 0045-0603-40 1



LOT: \_\_\_\_\_

EXP: \_\_\_\_\_

**Important: Read all product information before using. Keep the box for important information. This product is intended for use in children.**

**Active ingredient (in each 5 mL)**      **Purpose**  
 ibuprofen 100 mg (NSAID)      Pain reliever/fever reducer

**Uses:** Temporarily relieves minor aches and pains due to the common cold, flu, sore throat, headache and toothache. Reduces fever.

**Warnings:**  
**Allergic alert:** Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: hives, facial swelling, rash, wheezing, shock, skin reddening, dizziness, fainting, loss of consciousness. If an allergic reaction occurs, stop use and seek medical help right away. **Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: have had stomach ulcers or bleeding; take a blood-thinning (anti-coagulant) or steroid drug; take over-the-counter aspirin or other NSAIDs (aspirin, ibuprofen, naproxen, or rofecoxib); take more or for a longer time than directed. **Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed. **Severe or persistent sore throat or sore throat** accompanied by high fever, headache, nausea, and vomiting may be serious. Consult your doctor promptly. Do not use more than 2 days or administer to children under 3 years of age unless directed by doctor. **Do not use if:** you have ever had an allergic reaction to ibuprofen or any other pain reliever/fever reducer; right before or after heart surgery. **Ask a doctor before use if:** stomach bleeding warning applies to you; you have a history of stomach problems, such as heartburn; you have had stomach problems or serious side effects from taking pain relievers or fever reducers; you have not been drinking fluids; your child has lost a lot of fluid due to vomiting or diarrhea; your child has high blood pressure, heart disease, liver or kidney disease, or had a stroke; your child has asthma; your child is taking a diuretic. **Ask a doctor or pharmacist before use if:** the child is under a doctor's care for any serious condition; taking any other drug; when using this product; taking with food or milk if stomach upset occurs. **Stop use and ask a doctor if:** you experience any of the following signs of stomach bleeding: vomits blood; has bloody or black stools; has stomach pain that does not get better; child has symptoms of heart problems or stroke; chest pain; labored breathing; weakness in one part of the body; slurred speech; leg swelling; the child does not get any relief within the first day (24 hours) of treatment; fever or pain gets worse or lasts more than 3 days; redness or swelling is present in the pain relief area; any new symptoms appear. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222).

**Directions:**  
 This product does not contain dextrose or complete warnings for adult use. Do not give to someone who is allergic to ibuprofen before using. Measure the right dose on the label below. If possible, use weight to dose; otherwise use age. Use only enclosed dosing cup. Do not use any other dosing device. If needed, repeat dose every 6-8 hours. Do not use more than 4 times a day. Replace original bottle cap to maintain child resistance.

**Dosing Chart**

Weight (lb)	Age (yr)	Dose (mL)*
under 24	under 2 years	ask a doctor
24-35 lbs	2-3 years	5 mL
36-47 lbs	4-5 years	7.5 mL
48-59 lbs	6-8 years	10 mL
60-71 lbs	9-10 years	12.5 mL
72-95 lbs	11 years	15 mL

**Other information:** Each 5 mL contains: sodium 2 mg. Store between 20-25°C (68-77°F). Do not use if bottle wrap imprinted "SAFETY SEAL" is broken or missing. **Inactive ingredients:** acacia, lactose, polyethylene glycol, anhydrous citric acid, D&C yellow no. 10, FD&C red no. 40, flavor, glycerin, polyethylene glycol 80, pregelatinized starch, purified water, sodium benzoate, sucrose, xanthan gum. **Questions or comments?** Call 1-877-895-3665 (Toll-Free) or 215-273-8755 (local).

Distributed by: **JOHNSON & JOHNSON CONSUMER INC.**  
 McNeil Consumer Healthcare Division  
 Fort Washington, PA 19034 USA ©J&JCI 2016 [www.motrin.com](http://www.motrin.com)  
 00000000

**FOR HOSPITAL/GOVERNMENT USE ONLY**  
 Not for Retail Sale      NDC 50580-601-21

**Children's Motrin**  
 contains **Ibuprofen Oral Suspension (NSAID)**  
**100 mg per 5 mL**  
**Pain Reliever/Fever Reducer      4 fl oz (120 mL)**  
**Original Berry Flavor**

(b) (4)

Reference ID: 4069064

**Drug Facts (continued)**

**Ask a doctor before use if**

- stomach bleeding warning applies to your child
- child has a history of stomach problems, such as heartburn
- child has problems or serious side effects from taking pain relievers or fever reducers
- child has not been drinking fluids
- child has lost a lot of fluid due to vomiting or diarrhea
- child has high blood pressure, heart disease, liver cirrhosis, kidney disease, or had a stroke
- child has asthma
- child is taking a diuretic

**Ask a doctor or pharmacist before use if the child is**

- under a doctor's care for any serious condition
- taking any other drug

**When using this product**

- take with food or milk if stomach upset occurs

**Stop use and ask a doctor if**

- child experiences any of the following signs of stomach bleeding:
  - feels faint
  - vomits blood
  - has bloody or black stools
- has stomach pain that does not get better
- child has symptoms of heart problems or stroke:
  - chest pain
  - trouble breathing
  - weakness in one part or side of body
  - slurred speech
  - leg swelling
- the child does not get any relief within first day (24 hours) of treatment
- fever or pain gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

**Directions**

- this product does not contain directions or complete warnings for adult use
- do not give more than directed

**Drug Facts (continued)**

- shake well before using
- mL = milliliter
- find right dose on chart. If possible, use weight to dose; otherwise use age.
- use only enclosed dosing cup. Do not use any other dosing device.
- if needed, repeat dose every **6-8 hours**
- do not use more than **4 times a day**
- replace original bottle cap to maintain child resistance

**Dosing Chart**

Weight (lb)	Age (yr)	Dose (mL)*
under 24	under 2 years	ask a doctor
24-35 lbs	2-3 years	5 mL
36-47 lbs	4-5 years	7.5 mL
48-59 lbs	6-8 years	10 mL
60-71 lbs	9-10 years	12.5 mL
72-95 lbs	11 years	15 mL

\* or as directed by a doctor

**Other information**

- each 5 mL contains: sodium 2 mg
- store between 20-25° C (68-77° F)
- do not use if carton is opened or if carton tape or bottle wrap imprinted "SAFETY SEAL®" is broken or missing

**Inactive ingredients** acesulfame potassium, anhydrous citric acid, D&C yellow no. 10, FD&C red no. 40, flavors, glycerin, polysorbate 80, pregelatinized starch, purified water, sodium benzoate, sucrose, xanthan gum

**Questions or comments?**  
Call 1-877-895-3665 (toll-free) or 215-273-8755 (collect)

Distributed by:  
**JOHNSON & JOHNSON CONSUMER INC.**  
McNeil Consumer Healthcare Division  
Fort Washington, PA 19034 USA  
©J&JCI 2016  
[www.motrin.com](http://www.motrin.com) 00000000

**FOR HOSPITAL/GOVERNMENT USE ONLY**  
For Ages 2 to 11 Years

**Children's Motrin®**  
contains **Ibuprofen Oral Suspension (NSAID)**  
Pain Reliever / Fever Reducer  
Original Berry Flavor

*See New Warnings*  
NDC 50580-601-21

**FOR HOSPITAL/GOVERNMENT USE ONLY**  
For Ages 2 to 11 Years

**Children's Motrin®**  
contains **Ibuprofen Oral Suspension (NSAID)**  
100 mg per 5 mL  
Pain Reliever / Fever Reducer  
Original Berry Flavor

**Lasts up to 8 hours**  
Alcohol Free

**4 fl oz (120 mL)**

50580 60121 1

LOT:  
EXP:

**Important: Read all product information before using. Keep this box for important information. This product is intended for use in children.**

**Drug Facts**

Active ingredient (in each 5 mL)	Purpose
ibuprofen 100 mg (NSAID)*.....Pain reliever/fever reducer	
*nonsteroidal anti-inflammatory drug	

**Uses temporarily:**

- relieves minor aches and pains due to the common cold, flu, sore throat, headache and toothache
- reduces fever

**Warnings**

**Allergy alert:** ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if your child:

- has had stomach ulcers or bleeding problems
- takes a blood thinning (anticoagulant) or site old drug
- takes other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- takes more or for a longer time than directed

**Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

**Sore throat warning:** Severe or persistent sore throat or sore throat accompanied by high fever, headache, nausea, and vomiting may be serious. Consult doctor promptly. Do not use more than 2 days or administer to children under 3 years of age unless directed by doctor.

**Do not use**

- if the child has ever had an allergic reaction to ibuprofen or any other pain reliever/fever reducer
- right before or after heart surgery

(b) (4)



(b) (4)