 Boehringer Ingelheim
Copy for NDA Submission
Product: Zantac 75®
Component: 75 mg 1ct Pouch
Date: July 2014

100%

FRONT




BACK



200%





FRONT




BACK

(b) (4)


 Boehringer Ingelheim
Copy for NDA Submission
Product: Zantac 75[®]
Component: 75 mg 4ct Blister Foil Label
Date: July 2014



100%



200%

 Boehringer Ingelheim
Copy for NDA Submission
Product: Zantac 75[®]
Component: 75 mg 4ct Blister Carton
Date: January 2015

Drug Facts (continued)

Inactive ingredients
hypromellose, magnesium stearate, microcrystalline cellulose, synthetic red iron oxide, titanium dioxide, triacetin

Questions? call 1-888-285-9159 (English/Spanish) M-F, 8:30-5 EST, or visit www.zantac.com

6 81421 03002 1



Drug Facts

Active ingredient (in each tablet)
Ranitidine 75 mg (as ranitidine hydrochloride 84 mg).....Acid reducer

Purpose
Acid reducer

Uses
relieves heartburn associated with acid indigestion and sour stomach
prevents heartburn associated with acid indigestion and sour stomach
brought on by eating or drinking certain foods and beverages

Warnings
Do not use if you are allergic to ranitidine or other acid reducers
See your doctor if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition.

Ask a doctor before use if you have
had heartburn over 3 months. This may be a sign of a more serious condition.
heartburn with lightheadedness, sweating or dizziness
chest pain or shoulder pain with shortness of breath, sweating, pain spreading to arms, neck or shoulders, or lightheadedness
frequent chest pain
unexplained weight loss
nausea or vomiting
stomach pain

Other information
do not use if individual blister unit is open or torn
store at 20°-25°C (68°-77°F)
avoid excessive heat or humidity
this product is sodium and sugar free

Directions
adults and children 12 years and over:
to relieve symptoms, swallow 1 tablet with a glass of water
to prevent symptoms, swallow 1 tablet with a glass of water 30 to 60 minutes before eating food or drinking beverages that cause heartburn
can be used up to twice daily (do not take more than 2 tablets in 24 hours)
children under 12 years: ask a doctor

Boehringer Ingelheim
Division of Bristol-Myers Squibb
Ingelheim Pharmaceuticals, Inc.
Cape May, NJ 08204
Copyright © 2015, Boehringer
Ingelheim Pharmaceuticals, Inc.
All rights reserved.
Made in Spain. Manufacturer of this item.

Drug Facts (continued)

Stop use and ask a doctor if:
your heartburn continues or worsens
you need to take this product for more than 14 days

If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

LOT

EXP

Zantac
Ranitidine Tablets 75 mg /Acid Reducer

75

ZANTAC does not supply store brands

Read the directions, consumer information leaflet and warnings before use. Keep the carton. It contains important information.

Zantac
Ranitidine Tablets 75 mg /Acid Reducer

75

**PREVENTS & RELIEVES
HEARTBURN** associated with
acid indigestion and sour stomach

**4 TABLETS
(4 DOSES)**



P011057-005



502


090320944/4
CT6118B

LOT

EXP



(b) (4)


 Boehringer Ingelheim
Copy for NDA Submission
Product: Zantac 75[®]
Component: 75 mg 10ct Blister Foil Label
Date: July 2014

Zantac 75[®] <small>RANITIDINE TABLETS 75 mg ACID REDUCER Boehringer Ingelheim Pharmaceuticals, Inc. Ridgefield, CT 06877</small>	Zantac 75[®] <small>RANITIDINE TABLETS 75 mg ACID REDUCER Boehringer Ingelheim Pharmaceuticals, Inc. Ridgefield, CT 06877</small>	Zantac 75[®] <small>RANITIDINE TABLETS 75 mg ACID REDUCER Boehringer Ingelheim Pharmaceuticals, Inc. Ridgefield, CT 06877</small>	Zantac 75[®] <small>RANITIDINE TABLETS 75 mg ACID REDUCER Boehringer Ingelheim Pharmaceuticals, Inc. Ridgefield, CT 06877</small>
Zantac 75[®] <small>RANITIDINE TABLETS 75 mg ACID REDUCER Boehringer Ingelheim Pharmaceuticals, Inc. Ridgefield, CT 06877</small>	Zantac 75[®] <small>RANITIDINE TABLETS 75 mg ACID REDUCER Boehringer Ingelheim Pharmaceuticals, Inc. Ridgefield, CT 06877</small> P011056-005 10009704 TA/202/US/3 PUSH THROUGH PRINTED SIDE.	Zantac 75[®] <small>RANITIDINE TABLETS 75 mg ACID REDUCER Boehringer Ingelheim Pharmaceuticals, Inc. Ridgefield, CT 06877</small>	Zantac 75[®] <small>RANITIDINE TABLETS 75 mg ACID REDUCER Boehringer Ingelheim Pharmaceuticals, Inc. Ridgefield, CT 06877</small>
Zantac 75[®] <small>RANITIDINE TABLETS 75 mg ACID REDUCER Boehringer Ingelheim Pharmaceuticals, Inc. Ridgefield, CT 06877</small>	Zantac 75[®] <small>RANITIDINE TABLETS 75 mg ACID REDUCER Boehringer Ingelheim Pharmaceuticals, Inc. Ridgefield, CT 06877</small>	Zantac 75[®] <small>RANITIDINE TABLETS 75 mg ACID REDUCER Boehringer Ingelheim Pharmaceuticals, Inc. Ridgefield, CT 06877</small>	Zantac 75[®] <small>RANITIDINE TABLETS 75 mg ACID REDUCER Boehringer Ingelheim Pharmaceuticals, Inc. Ridgefield, CT 06877</small>

100%

Zantac 75[®] <small>RANITIDINE TABLETS 75 mg ACID REDUCER Boehringer Ingelheim Pharmaceuticals, Inc. Ridgefield, CT 06877</small>	Zantac 75[®] <small>RANITIDINE TABLETS 75 mg ACID REDUCER Boehringer Ingelheim Pharmaceuticals, Inc. Ridgefield, CT 06877</small>	Zantac 75[®] <small>RANITIDINE TABLETS 75 mg ACID REDUCER Boehringer Ingelheim Pharmaceuticals, Inc. Ridgefield, CT 06877</small>	Zantac 75[®] <small>RANITIDINE TABLETS 75 mg ACID REDUCER Boehringer Ingelheim Pharmaceuticals, Inc. Ridgefield, CT 06877</small>
Zantac 75[®] <small>RANITIDINE TABLETS 75 mg ACID REDUCER Boehringer Ingelheim Pharmaceuticals, Inc. Ridgefield, CT 06877</small>	Zantac 75[®] <small>RANITIDINE TABLETS 75 mg ACID REDUCER Boehringer Ingelheim Pharmaceuticals, Inc. Ridgefield, CT 06877</small> P011056-005 10009704 TA/202/US/3 PUSH THROUGH PRINTED SIDE.	Zantac 75[®] <small>RANITIDINE TABLETS 75 mg ACID REDUCER Boehringer Ingelheim Pharmaceuticals, Inc. Ridgefield, CT 06877</small>	Zantac 75[®] <small>RANITIDINE TABLETS 75 mg ACID REDUCER Boehringer Ingelheim Pharmaceuticals, Inc. Ridgefield, CT 06877</small>
Zantac 75[®] <small>RANITIDINE TABLETS 75 mg ACID REDUCER Boehringer Ingelheim Pharmaceuticals, Inc. Ridgefield, CT 06877</small>	Zantac 75[®] <small>RANITIDINE TABLETS 75 mg ACID REDUCER Boehringer Ingelheim Pharmaceuticals, Inc. Ridgefield, CT 06877</small>	Zantac 75[®] <small>RANITIDINE TABLETS 75 mg ACID REDUCER Boehringer Ingelheim Pharmaceuticals, Inc. Ridgefield, CT 06877</small>	Zantac 75[®] <small>RANITIDINE TABLETS 75 mg ACID REDUCER Boehringer Ingelheim Pharmaceuticals, Inc. Ridgefield, CT 06877</small>

200%

 Boehringer Ingelheim	
Copy for NDA Submission	
Product:	Zantac 75[®]
Component:	75 mg 10ct Blister Carton
Date:	January 2015

LOT
EXP

Drug Facts (continued)

M-F, 8:30-5 EST, or visit www.zantacac.com
 Questions? call 1-888-285-9159 (English/Spanish)
 Boehringer Ingelheim Consumer Health Care
 Products Division of Boehringer Ingelheim
 Pharmaceuticals, Inc., Ridgefield, CT 06877
 Copyright © 2015, Boehringer Ingelheim
 Pharmaceuticals, Inc. All rights reserved.
 Product of Spain. Manufactured in Mexico.

Drug Facts
 Ranitidine 75 mg (as ranitidine hydrochloride 84 mg).....Acid reducer

Uses
 • relieves heartburn associated with acid indigestion and sour stomach
 • prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain foods and beverages

Warnings
 • if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.
 • with other acid reducers

Do not use
 • if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.

Allergy alert: Do not use if you are allergic to ranitidine or other acid reducers

Ask a doctor before use if you have
 • had heartburn over 3 months. This may be a sign of a more serious condition.
 • heartburn with lightheadedness, sweating or dizziness
 • chest pain or shoulder pain with shortness of breath, sweating, pain spreading to arms, neck or shoulders, or lightheadedness
 • frequent chest pain • frequent wheezing, particularly with heartburn • unexplained weight loss
 • nausea or vomiting • stomach pain

Stop use and ask a doctor if • your heartburn continues or worsens • you need to take this product for more than 14 days

If pregnant or breast-feeding, ask a health professional before use.
 Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions
 • adults and children 12 years and over:
 • to relieve symptoms, swallow 1 tablet with a glass of water
 • to prevent symptoms, swallow 1 tablet with a glass of water **30 to 60 minutes before** eating food or drinking beverages
 • can be used up to twice daily (do not take more than 2 tablets in 24 hours)
 • children under 12 years: ask a doctor

Other information
 • do not use if individual blister unit is open or torn
 • avoid excessive heat or humidity
 • store at 20°-25°C (68°-77°F)
 • this product is sodium and sugar free

Inactive ingredients hypromellose, magnesium stearate, microcrystalline cellulose, synthetic red iron oxide, titanium dioxide, triacetin

81421 03003 8

ZANTAC does not supply store brands.

Read the directions,
consumer information leaflet and warnings before use.
Keep the carton. It contains important information.

090320945/4
CT16120A

PO11060-006



Zantac
Ranitidine Tablets 75 mg / Acid Reducer

75

Zantac

Ranitidine Tablets 75 mg / Acid Reducer

75



PREVENTS & RELIEVES
HEARTBURN

associated with
acid indigestion and sour stomach

10


TABLETS
(10 DOSES)



503



(b) (4)

 Boehringer Ingelheim
Copy for NDA Submission
Product: Zantac 75®
Component: 75 mg 60ct Bottle Label
Date: July 2014

Warnings Do not use • if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor. • with other acid reducers **Ask a doctor before use if you have** • had heartburn over 3 months. This may be a sign of a more serious condition. • heartburn with **lightheadedness, sweating or dizziness** • chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness • frequent **chest pain** • frequent wheezing, particularly with heartburn • unexplained weight loss • nausea or vomiting • stomach pain. **Stop use and ask a doctor if** • your heartburn continues or worsens • you need to take this product for more than 14 days. **If pregnant or breast-feeding**, ask a health professional before use. **Keep out of reach of children**. In case of overdose, get medical help or contact a Poison Control Center right away.

Bottle sealed with printed foil under cap. Do not use if foil is open or torn.

Zantac
RANITIDINE TABLETS
75 mg / ACID REDUCER

60
TABLETS
(60 DOSES)

**PREVENTS & RELIEVES
HEARTBURN** associated
with acid indigestion and
sour stomach

Directions • adults and children 12 years and over: • to relieve symptoms, swallow 1 tablet with a glass of water • to prevent symptoms, swallow 1 tablet with a glass of water **30 to 60 minutes before** eating food or drinking beverages that cause heartburn • can be used up to twice daily (do not take more than 2 tablets in 24 hours) • children under 12 years: ask a doctor. **Questions? call 1-888-285-9159** (English/Spanish) M – F, 8:30 – 5 EST. Or visit www.zantacotc.com Copyright © 2014, Boehringer Ingelheim Pharmaceuticals, Inc., Ridgefield, CT. All rights reserved. 10009595 4/14 001 P033123-001

LOT
EXP

090330988/1

70

100%

Warnings Do not use • if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor. • with other acid reducers **Ask a doctor before use if you have** • had heartburn over 3 months. This may be a sign of a more serious condition. • heartburn with **lightheadedness, sweating or dizziness** • chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness • frequent **chest pain** • frequent wheezing, particularly with heartburn • unexplained weight loss • nausea or vomiting • stomach pain. **Stop use and ask a doctor if** • your heartburn continues or worsens • you need to take this product for more than 14 days. **If pregnant or breast-feeding**, ask a health professional before use. **Keep out of reach of children**. In case of overdose, get medical help or contact a Poison Control Center right away.

Bottle sealed with printed foil under cap. Do not use if foil is open or torn.

Zantac
RANITIDINE TABLETS
75 mg / ACID REDUCER

60
TABLETS
(60 DOSES)

**PREVENTS & RELIEVES
HEARTBURN** associated
with acid indigestion and
sour stomach

Directions • adults and children 12 years and over: • to relieve symptoms, swallow 1 tablet with a glass of water • to prevent symptoms, swallow 1 tablet with a glass of water **30 to 60 minutes before** eating food or drinking beverages that cause heartburn • can be used up to twice daily (do not take more than 2 tablets in 24 hours) • children under 12 years: ask a doctor. **Questions? call 1-888-285-9159** (English/Spanish) M – F, 8:30 – 5 EST. Or visit www.zantacotc.com Copyright © 2014, Boehringer Ingelheim Pharmaceuticals, Inc., Ridgefield, CT. All rights reserved. 10009595 4/14 001 P033123-001


LOT
EXP

090330988/1

70

200%



 Boehringer Ingelheim
Copy for NDA Submission
Product: Zantac 75®
Component: 75 mg 60ct Bottle Carton
Date: January 2015



Zantac 75
Ranitidine Tablets 75 mg / Acid Reducer

60 TABLETS (60 DOSES)

PREVENTS & RELIEVES
HEARTBURN associated with
acid indigestion and sour stomach



090320949/5
CT6123A
P011066-007

Zantac 75
Ranitidine Tablets 75 mg / Acid Reducer



508

Zantac 75
Ranitidine Tablets 75 mg / Acid Reducer

60 TABLETS (60 DOSES)



Drug Facts

Active ingredient (in each tablet) Ranitidine 75 mg (as ranitidine hydrochloride 84 mg)	Purpose Acid reducer
Uses	
<ul style="list-style-type: none"> • relieves heartburn associated with acid indigestion and sour stomach • prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain foods and beverages 	
Warnings	
Allergy alert: Do not use if you are allergic to ranitidine or other acid reducers	
Do not use	
<ul style="list-style-type: none"> • if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor. • with other acid reducers 	
Ask a doctor before use if you have	
<ul style="list-style-type: none"> • had heartburn over 3 months. This may be a sign of a more serious condition. • heartburn with lightheadedness, sweating or dizziness • chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness • frequent chest pain • frequent wheezing, particularly with heartburn • unexplained weight loss • nausea or vomiting • stomach pain 	
Stop use and ask a doctor if • your heartburn continues or worsens • you need to take this product for more than 14 days	
If pregnant or breast-feeding, ask a health professional before use.	
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.	
Directions	
<ul style="list-style-type: none"> • adults and children 12 years and over: <ul style="list-style-type: none"> • to relieve symptoms, swallow 1 tablet with a glass of water • to prevent symptoms, swallow 1 tablet with a glass of water 30 to 60 minutes before eating food or drinking beverages that cause heartburn • can be used up to twice daily (do not take more than 2 tablets in 24 hours) • children under 12 years: ask a doctor 	
Other information	
<ul style="list-style-type: none"> • do not use if printed foil under bottle cap is open or torn • avoid excessive heat or humidity 	<ul style="list-style-type: none"> • store at 20°-25°C (68°-77°F) • this product is sodium and sugar free

Zantac 75
Ranitidine Tablets 75 mg / Acid Reducer

60 TABLETS (60 DOSES)



LOT
EXP

ZANTAC does not supply store brands

6 81421 03006 9



Drug Facts (continued)

Inactive ingredients hypromellose, magnesium stearate, microcrystalline cellulose, synthetic red iron oxide, titanium dioxide, triacetin

Questions? call 1-888-285-9159 (English/Spanish) M - F, 8:30 - 5 EST, or visit www.zantacotc.com

Read the directions, consumer information leaflet and warnings before use. Keep the carton. It contains important information.

 **Boehringer Ingelheim** Boehringer Ingelheim Consumer Health Care Products
 Division of Boehringer Ingelheim Pharmaceuticals, Inc., Ridgefield, CT 06877
 Copyright © 2015, Boehringer Ingelheim Pharmaceuticals, Inc. All rights reserved.
 Product of Spain. Manufactured in Mexico.

LOT
EXP

090320949/5
CT6123A
P011066-007


Zantac 75
Ranitidine Tablets 75 mg / Acid Reducer




508




(b) (4)

 Boehringer Ingelheim
Copy for NDA Submission
Product: Zantac 75®
Component: 75 mg 80ct Bottle Label
Date: July 2014




100%



200%



 Boehringer Ingelheim
Copy for NDA Submission
Product: Zantac 75[®]
Component: 75 mg 80ct Bottle Carton
Date: January 2015



Zantac 75[®]
Ranitidine Tablets 75 mg / Acid Reducer

80 TABLETS (80 DOSES)

PREVENTS & RELIEVES HEARTBURN
associated with acid indigestion and sour stomach



090320950/5
CT6125A
P011088-007

Zantac 75[®]
Ranitidine Tablets 75 mg / Acid Reducer



5 5

Zantac 75[®]
Ranitidine Tablets 75 mg / Acid Reducer

80 TABLETS (80 DOSES)



6 81421 03007 6

Drug Facts

Active ingredient (in each tablet)	Purpose
Ranitidine 75 mg (as ranitidine hydrochloride 84 mg)	Acid reducer

Uses

- relieves heartburn associated with acid indigestion and sour stomach
- prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain foods and beverages

Warnings

Allergy alert: Do not use if you are allergic to ranitidine or other acid reducers

Do not use

- if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.
- with other acid reducers

Ask a doctor before use if you have

- had heartburn over 3 months. This may be a sign of a more serious condition.
- heartburn with **lightheadedness, sweating or dizziness**
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
- frequent **chest pain** • frequent wheezing, particularly with heartburn • unexplained weight loss
- nausea or vomiting • stomach pain

Stop use and ask a doctor if • your heartburn continues or worsens • you need to take this product for more than 14 days

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- adults and children 12 years and over:
 - to **relieve** symptoms, swallow 1 tablet with a glass of water
 - to **prevent** symptoms, swallow 1 tablet with a glass of water **30 to 60 minutes before** eating food or drinking beverages that cause heartburn
 - can be used up to twice daily (do not take more than 2 tablets in 24 hours)
- children under 12 years: ask a doctor

Other information

- do not use if printed foil under bottle cap is open or torn
- store at 20°-25°C (68°-77°F)
- avoid excessive heat or humidity
- this product is sodium and sugar free

Zantac 75[®]
Ranitidine Tablets 75 mg / Acid Reducer

80 TABLETS (80 DOSES)

LOT
EXP

ZANTAC does not supply store brands

Drug Facts (continued)

Inactive ingredients hypromellose, magnesium stearate, microcrystalline cellulose, synthetic red iron oxide, titanium dioxide, triacetin

Questions? call 1-888-285-9159 (English/Spanish) M-F, 8:30-5 EST, or visit www.zantacotc.com


Read the directions, consumer information leaflet and warnings before use. Keep the carton. It contains important information.

 **Boehringer Ingelheim** Boehringer Ingelheim Consumer Health Care Products
 Division of Boehringer Ingelheim Pharmaceuticals, Inc., Ridgefield, CT 06877
 Copyright © 2015, Boehringer Ingelheim Pharmaceuticals, Inc. All rights reserved.
 Product of Spain. Manufactured in Mexico.





(b) (4)

 Boehringer Ingelheim
Copy for NDA Submission
Product: Zantac 75[®]
Component: 75 mg Consumer Leaflet
Date: July 2014




What you should know about

Zantac

RANITIDINE TABLETS 75 mg 75[®]
ACID REDUCER

(Please read all of this information before taking Zantac 75. Save this leaflet for future reference.)

- Zantac 75 contains 75 mg of ranitidine (as ranitidine hydrochloride, 84 mg), a medicine that doctors have prescribed more than 200 million times worldwide.
- Zantac 75 has been used safely and effectively for years as an over-the-counter medication.
- Zantac 75 tablets are sodium and sugar free.



- Zantac 75 relieves and prevents heartburn associated with acid indigestion and sour stomach.

Certain foods or beverages, and even lying down to sleep, can cause heartburn associated with acid indigestion or sour stomach. It is normal for the stomach to produce acid, especially after consuming food or beverages.

However, acid in the wrong place, such as the esophagus, or too much acid, can cause burning pain and discomfort.

- To **relieve** symptoms, swallow 1 tablet with a glass of water.
- To **prevent** symptoms, swallow 1 tablet with a glass of water **30 to 60 minutes before** eating food or drinking beverages that cause heartburn.

This medicine can be used up to twice daily (do not take more than 2 tablets in 24 hours).

Zantac 75 should not be given to children under 12 years old unless directed by a doctor.

- **Allergy alert:** Do not use if you are allergic to ranitidine or other acid reducers
- Zantac 75 reduces the production of stomach acid. This is what makes Zantac 75 different from antacids, which neutralize the acid already in your stomach. Antacids do not reduce the production of acid.
- Do not lie flat or bend over soon after eating
- Do not eat late at night, or just before bedtime
- Certain foods or drinks are more likely to cause heartburn, such as rich, spicy, fatty, and fried foods, chocolate, caffeine, alcohol, even some fruits and vegetables
- Eat slowly and do not eat big meals
- If you are overweight, lose weight
- If you smoke, quit smoking
- Raise the head of your bed
- Wear loose fitting clothing around your stomach

090340239/5

What is Zantac 75?

Excellent Safety Record

What symptoms does Zantac 75 relieve and prevent?

How should I take Zantac 75?

How does Zantac 75 work?



Additional tips for managing heartburn

Clinical studies prove Zantac 75 is effective in relieving ...

... and preventing heartburn.

When should I see a doctor?

In clinical studies, Zantac 75 was significantly better than placebo pills.



- **Do not use**
 - if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.
 - with other acid reducers
- **Ask a doctor before use if you have**
 - had heartburn over 3 months. This may be a sign of a more serious condition.
 - heartburn with **lightheadedness, sweating or dizziness**
 - chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
 - frequent **chest pain**
 - frequent wheezing, particularly with heartburn
 - unexplained weight loss
 - nausea or vomiting
 - stomach pain
- **Stop use and ask a doctor if**
 - your heartburn continues or worsens
 - you need to take this product for more than 14 days
- **If pregnant or breast-feeding**, ask a healthcare professional before use.
- **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

Questions? call 1-888-285-9159 (English/Spanish) M - F, 8:30 - 5 EST, or visit www.zantacotc.com

BOTTLES: Bottle is sealed with printed foil under cap. Do not use if foil is open or torn.

BLISTERS: Do not use if the individual blister unit is open or torn.

Boehringer Ingelheim Consumer Health Care Products Division of Boehringer Ingelheim Pharmaceuticals, Inc. Ridgefield, CT 06877 Copyright © 2014, Boehringer Ingelheim Pharmaceuticals, Inc. All rights reserved.

P011212-005
10009709
04/14 003

090340239/5