



| Job Information | Color Legend |
|---|---|
| Date 6/27/03 PASS #3 Component # UK24326-1 Drawing # 3919 Rev 3 | |
| Identification Production Packaging Dept. Five / Farms Madison, NJ 07940 Phone: (973) 660-5595/5153 Fax: (973) 660-6048 | |
| Vendor Identification Production Packaging Dept. Linda Hill 6.27.3 | Printer Notes PRINTER TO REPLACE FPO UPC'S WITH HI-RES DIGITAL ART ALSO PRINTER TO SUPPLY COLOR MATCHPRINT FOR COLOR APPROVAL PRIOR TO PRESS RUN. |

| DRUG FACTS TEXT DEFINED | TYPE SIZE |
|---------------------------------------|---------------|
| • DRUG FACTS Title | 9 pt |
| • DRUG FACTS CONTINUED | 8 pt |
| • HEADINGS | 8 pt |
| • SUBHEADS/BODY TEXT | 6 pt |
| • LEADING | 6.5 pt |
| • # OF CHARACTERS PER INCH | <39 |
| • BULLETS | 5 pt |
| • SPACE AFTER BULLETED SECTION | 2 ems |
| • BARLINES, HAIRLINES | 1.5 pt, .5 pt |
| • SPACE BETWEEN HAIRLINES AND BOX END | 2 spaces |

| DEPARTMENT | REVIEWED BY | | DATE | ✓ with Changes |
|---------------------|-------------|-----------|------|--------------------------|
| | Print Name | Signature | | |
| Graphic Artist | | | | <input type="checkbox"/> |
| Packaging Engineer | | | | <input type="checkbox"/> |
| Labeling Compliance | | | | <input type="checkbox"/> |
| Medical | | | | <input type="checkbox"/> |
| Regulatory Affairs | | | | <input type="checkbox"/> |
| Marketing | | | | <input type="checkbox"/> |
| Legal | | | | <input type="checkbox"/> |
| Trademark | | | | <input type="checkbox"/> |

| RELEASED | |
|-----------|------------|
| (1) _____ | Date _____ |
| (2) _____ | Date _____ |

DIGITAL MECHANICAL
COLOR REPRESENTATION MAY BE OFF
FOLLOW COLOR LEGEND FOR ACTUAL COLOR.



| Job Information | Color Legend |
|---|--|
| Date 6/27/03 PASS #3 Component # UK24323-1 Drawing # 3919 Rev 3 | |
| Identification Wyeth Consumer Healthcare Production Packaging Dept. Five / Farms Madison, NJ 07940 Phone: (973) 660-5595/5153 Fax: (973) 660-6048 | |
| Vendor Identification Production Packaging Dept. Linda Hill SC: 6.27.3 | Printer Notes PRINTER TO REPLACE FPO UPC'S WITH HI-RES DIGITAL ART ALSO PRINTER TO SUPPLY COLOR MATCHPRINT FOR COLOR APPROVAL PRIOR TO PRESS RUN. |

| DRUG FACTS TEXT DEFINED | TYPE SIZE |
|---------------------------------------|---------------|
| • DRUG FACTS Title | 9 pt |
| • DRUG FACTS CONTINUED | 8 pt |
| • HEADINGS | 8 pt |
| • SUBHEADS/BODY TEXT | 6 pt |
| • LEADING | 6.5 pt |
| • # OF CHARACTERS PER INCH | <39 |
| • BULLETS | 5 pt |
| • SPACE AFTER BULLETED SECTION | 2 ems |
| • BARLINES, HAIRLINES | 1.5 pt, .5 pt |
| • SPACE BETWEEN HAIRLINES AND BOX END | 2 spaces |

| DEPARTMENT | REVIEWED BY | | DATE | ✓ with Changes |
|---------------------|-------------|-----------|------|--------------------------|
| | Print Name | Signature | | |
| Graphic Artist | | | | <input type="checkbox"/> |
| Packaging Engineer | | | | <input type="checkbox"/> |
| Labeling Compliance | | | | <input type="checkbox"/> |
| Medical | | | | <input type="checkbox"/> |
| Regulatory Affairs | | | | <input type="checkbox"/> |
| Marketing | | | | <input type="checkbox"/> |
| Legal | | | | <input type="checkbox"/> |
| Trademark | | | | <input type="checkbox"/> |

RELEASED

(1) _____ Date _____

(2) _____ Date _____

DIGITAL MECHANICAL
COLOR REPRESENTATION MAY BE OFF
FOLLOW COLOR LEGEND FOR ACTUAL COLOR.

Children's Advil[®]
Suspension

Do Not Use if plastic wrap imprinted with "CHILDREN'S ADVIL SAFETY SEALED" is broken or missing.

Children's Advil[®]
Suspension

Ibuprofen 100 mg >> Fever Reducer/Pain Reliever
per 5 mL

Fever
Aches & Pains

Lasts Up To
8
Hours

Now BLUE RASPBERRY!
Blue Raspberry Flavored Liquid
for ages 2-11 years

4 FL OZ (120 mL) Alcohol Free

Wyeth[®]
Wyeth Consumer Healthcare
Madison, NJ 07940
Made in USA
U.S. Patent No 4,788,220

F.P.O.

3 0573-0174-30 5

UK24321-1

FPO 128 Code #243211

Children's Advil[®]
Suspension

Drug Facts (continued)
Ask a doctor or pharmacist before use if the child is

- under a doctor's care for any serious condition
- taking any other product that contains ibuprofen, or any other pain reliever/fever reducer
- taking a prescription product for anticoagulation (blood thinning)
- taking any other drug

When using this product give with food or milk if stomach upset occurs
Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away.
- fever or pain gets worse or lasts more than 3 days
- the child does not get any relief within first day (24 hours) of treatment
- stomach pain or upset gets worse or lasts
- redness or swelling is present in the painful area
- any new symptoms appear

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not give more than directed
- shake well before using
- find right dose on chart below. If possible, use weight to dose; otherwise use age.
- repeat dose every 6-8 hours, if needed
- do not use more than 4 times a day
- measure only with the blue dosing cup provided.

Blue dosing cup to be used with Children's Advil Suspension only. Do not use with other products. Dose lines account for product remaining in cup due to thickness of suspension.

| Weight (lb) | Age (yr) | Dose (tsp) |
|-------------|------------|--------------|
| under 24 lb | under 2 yr | ask a doctor |
| 24-35 lb | 2-3 yr | 1 tsp |
| 36-47 lb | 4-5 yr | 1 1/2 tsp |
| 48-59 lb | 6-8 yr | 2 tsp |
| 60-71 lb | 9-10 yr | 2 1/2 tsp |
| 72-95 lb | 11 yr | 3 tsp |

Children's Advil[®]
Suspension

Drug Facts (continued)
Other information

- one dose lasts 6-8 hours
- store at 20-25°C (68-77°F)
- see bottom of box for lot number and expiration date

Inactive ingredients
carboxymethylcellulose sodium, citric acid, edetate disodium, FD&C blue no. 1, flavors, glycerin, microcrystalline cellulose, polysorbate 80, purified water, sodium benzoate, sodium citrate, sorbitol solution, sucrose, xanthan gum

Questions or comments?
Call weekdays from 9am to 5pm EST at **1-800-88-ADVIL** or ask your pharmacist, doctor or health care professional

Long Lasting: Up to 8 hours of fever relief.
Great Convenience: Long duration means fewer doses needed per day.
Alcohol free

Important: Read all product information before using. Keep this box for important information. This product is intended for use in children ages 2 years - 11 years.

Drug Facts
Active ingredients **Purposes**
(in each 5 mL)
Ibuprofen 100 mg.....Fever reducer/Pain reliever

Uses temporarily:

- reduces fever
- relieves minor aches and pains due to the common cold, flu, sore throat, headaches and toothaches

Warnings
Allergy alert: Ibuprofen may cause a severe allergic reaction which may include:

- hives
- asthma (wheezing)
- facial swelling
- shock

Stomach bleeding warning: Taking more than recommended may cause stomach bleeding.
Sore throat warning: Severe or persistent sore throat or sore throat accompanied by high fever, headache, nausea, and vomiting may be serious. Consult doctor promptly. Do not use more than 2 days or administer to children under 3 years of age unless directed by doctor.
Do not use if the child has ever had an allergic reaction to any other fever reducer/pain reliever

Ask a doctor before use if the child has

- not been drinking fluids
- ulcers
- bleeding problems
- lost a lot of fluid due to continued vomiting or diarrhea
- problems or serious side effects from taking fever reducers or pain relievers
- stomach problems that last or come back, such as heartburn, upset stomach or pain
- high blood pressure, heart or kidney disease or is taking a diuretic

| Job Information | Color Legend |
|--|---|
| Date 7/14/03 PASS #3 Component # UK24321-1 Drawing # 3919 Rev 3 | |
| Identification Production Packaging Dept. Five / Farms Madison, NJ 07940 Phone: (973) 660-5595/5153 Fax: (973) 660-6048 | Printer Notes PRINTER TO REPLACE FPO UPC'S WITH HI-RES DIGITAL ART ALSO PRINTER TO SUPPLY COLOR MATCHPRINT FOR COLOR APPROVAL PRIOR TO PRESS RUN. |
| Vendor Identification | |

| DRUG FACTS TEXT DEFINED | TYPE SIZE |
|---------------------------------------|---------------|
| • DRUG FACTS Title | 9 pt |
| • DRUG FACTS CONTINUED | 8 pt |
| • HEADINGS | 8 pt |
| • SUBHEADS/BODY TEXT | 6 pt |
| • LEADING | 6.5 pt |
| • # OF CHARACTERS PER INCH | <39 |
| • BULLETS | 5 pt |
| • SPACE AFTER BULLETED SECTION | 2 ems |
| • BARLINES, HAIRLINES | 1.5 pt, .5 pt |
| • SPACE BETWEEN HAIRLINES AND BOX END | 2 spaces |


| DEPARTMENT | REVIEWED BY | | DATE | ✓ with Changes |
|---------------------|-------------|-----------|------|--------------------------|
| | Print Name | Signature | | |
| Graphic Artist | | | | <input type="checkbox"/> |
| Packaging Engineer | | | | <input type="checkbox"/> |
| Labeling Compliance | | | | <input type="checkbox"/> |
| Medical | | | | <input type="checkbox"/> |
| Regulatory Affairs | | | | <input type="checkbox"/> |
| Marketing | | | | <input type="checkbox"/> |
| Legal | | | | <input type="checkbox"/> |
| Trademark | | | | <input type="checkbox"/> |

RELEASED

(1) _____ Date _____

(2) _____ Date _____

DIGITAL MECHANICAL
COLOR REPRESENTATION MAY BE OFF
FOLLOW COLOR LEGEND FOR ACTUAL COLOR.



| DRUG FACTS TEXT DEFINED | TYPE SIZE |
|--------------------------------|-----------|
| • HEADINGS | 4.5 pt |
| • SUBHEADS/BODY TEXT | 4.5 pt |
| • LEADING | 5 pt |
| • BULLETS | 3.5 |
| • SPACE AFTER BULLETED SECTION | 2 ems |
| • WARNING BOX LINE, HAIRLINES | .5 |

| Job Information | Color Legend |
|--|---|
| Date 12/17/03 PASS #3 Component # U0171-30-N2 Drawing # 5691 Rev.2 | |
| Identification Wyeth Consumer Healthcare | |
| Production Packaging Dept. Five / Farms Madison, NJ 07940 Phone: (973) 660-5595/5153 Fax: (973) 660-6048 | |
| Vendor Identification | Printer Notes |
| PPD: Cheng Lin | PRINTER TO REPLACE FPO UPC'S WITH HI-RES DIGITAL ART ALSO PRINTER TO SUPPLY COLOR MATCHPRINT FOR COLOR APPROVAL PRIOR TO PRESS RUN. |

| DEPARTMENT | REVIEWED BY | | DATE | ✓ with Changes |
|---------------------|-------------|-----------|------|--------------------------|
| | Print Name | Signature | | |
| Graphic Artist | | | | <input type="checkbox"/> |
| Packaging Engineer | | | | <input type="checkbox"/> |
| Labeling Compliance | | | | <input type="checkbox"/> |
| Medical | | | | <input type="checkbox"/> |
| Regulatory Affairs | | | | <input type="checkbox"/> |
| Marketing | | | | <input type="checkbox"/> |
| Legal | | | | <input type="checkbox"/> |
| Trademark | | | | <input type="checkbox"/> |

RELEASED

(1) _____ Date _____

(2) _____ Date _____

**DIGITAL MECHANICAL
COLOR REPRESENTATION MAY BE OFF
FOLLOW COLOR LEGEND FOR ACTUAL COLOR.**

**UNVARNISHED
UNPRINTED
AREA FOR LOT
& EXP.**

Warnings
Allergy alert: Ibuprofen may cause a severe allergic reaction which may include:
 ■ hives ■ asthma (wheezing) ■ facial swelling ■ shock
Stomach bleeding warning: Taking more than recommended may cause stomach bleeding.
Sore throat warning: Severe or persistent sore throat or sore throat accompanied by high fever, headache, nausea, and vomiting may be serious. Consult doctor promptly. Do not use more than 2 days or administer to children under 3 years of age unless directed by doctor.
Do not use if the child has ever had an allergic reaction to any other fever reducer/pain reliever.
Ask a doctor before use if the child has
 ■ not been drinking fluids ■ ulcers ■ bleeding problems
 ■ lost a lot of fluid due to continued vomiting or diarrhea
 ■ problems or serious side effects from taking fever reducers or pain relievers
 ■ stomach problems that last or come back, such as heartburn, upset stomach or pain
 ■ high blood pressure, heart or kidney disease or is taking a diuretic
Ask a doctor or pharmacist before use if the child is
 ■ under a doctor's care for any serious condition
 ■ taking any other product that contains ibuprofen, or any other pain reliever/fever reducer
 ■ taking a prescription product for anticoagulation (blood thinning)
 ■ taking any other drug
When using this product give with food or milk if upset stomach occurs
Stop use and ask a doctor if
 ■ an allergic reaction occurs. Seek medical help right away.
 ■ fever or pain gets worse or lasts more than 3 days
 ■ the child does not get any relief within first day (24 hours) of treatment
 ■ stomach pain or upset gets worse or lasts
 ■ redness or swelling is present in the painful area
 ■ any new symptoms appear

Do Not Use if plastic wrap imprinted with "CHILDREN'S ADVIL SAFETY SEALED" is broken or missing.

Children's Advil Suspension
Fever Reducer/Pain Reliever

Active ingredient (in each 5 mL).....Ibuprofen 100 mg
Uses temporarily:
 ■ reduces fever
 ■ relieves minor aches and pains due to the common cold, flu, sore throat, headaches and toothaches

One Dose Lasts 6-8 Hours


Grape Flavored Liquid
4 FL OZ (120 mL)

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.
Directions
 ■ do not give more than directed ■ shake well before using
 ■ find right dose on chart below. If possible, use weight to dose; otherwise use age.
 ■ repeat dose every 6-8 hours, if needed ■ do not use more than 4 times a day
 ■ measure only with the blue dosing cup provided. Blue dosing cup to be used with Children's Advil Suspension only. Do not use with other products. Dose lines account for product remaining in cup due to thickness of suspension.

| Weight (lb) | Age (yr) | Dose (tsp) |
|-------------|------------|--------------|
| under 24 lb | under 2 yr | ask a doctor |
| 24-35 lb | 2-3 yr | 1 tsp |
| 36-47 lb | 4-5 yr | 1 1/2 tsp |
| 48-59 lb | 6-8 yr | 2 tsp |
| 60-71 lb | 9-10 yr | 2 1/2 tsp |
| 72-95 lb | 11 yr | 3 tsp |

One Dose Lasts 6-8 Hours

Store at 20-25°C (68-77°F)
Questions or comments? Call toll-free 1-800-88-ADVIL or ask your pharmacist, doctor or health care professional
 Wyeth Consumer Healthcare, Madison, NJ 07940 Made in USA
 U.S. Patent No. 4,788,220 U0171-30-N2



100%

| DRUG FACTS TEXT DEFINED | TYPE SIZE |
|--------------------------------|-----------|
| • HEADINGS | 4.5 pt |
| • SUBHEADS/BODY TEXT | 4.5 pt |
| • LEADING | 5 pt |
| • BULLETS | 3.5 |
| • SPACE AFTER BULLETED SECTION | 2 ems |
| • WARNING BOX LINE, HAIRLINES | .5 |

| Job Information | Color Legend |
|--|---|
| Date 12/16/03 PASS #4 Component # U0170-30-N2 Drawing # 5691 Rev2 | |
| Identification Wyeth Consumer Healthcare Production Packaging Dept. Five / Farms Madison, NJ 07940 Phone: (973) 660-5595/5153 Fax: (973) 660-6048 | Printer Notes PRINTER TO REPLACE FPO UPC'S WITH HI-RES DIGITAL ART ALSO PRINTER TO SUPPLY COLOR MATCHPRINT FOR COLOR APPROVAL PRIOR TO PRESS RUN. |
| Vendor Identification Production Packaging Dept. Cheng Lin SC: 7.1.3 | |

Warnings
Allergy alert: Ibuprofen may cause a severe allergic reaction which may include:
 ■ hives ■ asthma (wheezing) ■ facial swelling ■ shock
Stomach bleeding warning: Taking more than recommended may cause stomach bleeding.
Sore throat warning: Severe or persistent sore throat or sore throat accompanied by high fever, headache, nausea, and vomiting may be serious. Consult doctor promptly. Do not use more than 2 days or administer to children under 3 years of age unless directed by doctor.
Do not use if the child has ever had an allergic reaction to any other fever reducer/pain reliever.
Ask a doctor before use if the child has
 ■ not been drinking fluids ■ ulcers ■ bleeding problems
 ■ lost a lot of fluid due to continued vomiting or diarrhea
 ■ problems or serious side effects from taking fever reducers or pain relievers
 ■ stomach problems that last or come back, such as heartburn, upset stomach or pain
 ■ high blood pressure, heart or kidney disease or is taking a diuretic
Ask a doctor or pharmacist before use if the child is
 ■ under a doctor's care for any serious condition
 ■ taking any other product that contains ibuprofen, or any other pain reliever/fever reducer
 ■ taking a prescription product for anticoagulation (blood thinning)
 ■ taking any other drug
When using this product give with food or milk if upset stomach occurs
Stop use and ask a doctor if
 ■ an allergic reaction occurs. Seek medical help right away.
 ■ fever or pain gets worse or lasts more than 3 days
 ■ the child does not get any relief within first day (24 hours) of treatment
 ■ stomach pain or upset gets worse or lasts
 ■ redness or swelling is present in the painful area
 ■ any new symptoms appear

Do Not Use if plastic wrap imprinted with "CHILDREN'S ADVIL SAFETY SEALED" is broken or missing.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions
 ■ do not give more than directed ■ shake well before using
 ■ find right dose on chart below. If possible, use weight to dose; otherwise use age.
 ■ repeat dose every 6-8 hours, if needed ■ do not use more than 4 times a day
 ■ measure only with the blue dosing cup provided. Blue dosing cup to be used with Children's Advil Suspension only. Do not use with other products. Dose lines account for product remaining in cup due to thickness of suspension.

Dosing Chart

| Weight (lb) | Age (yr) | Dose (tsp) |
|-------------|------------|--------------|
| under 24 lb | under 2 yr | ask a doctor |
| 24-35 lb | 2-3 yr | 1 tsp |
| 36-47 lb | 4-5 yr | 1 1/2 tsp |
| 48-59 lb | 6-8 yr | 2 tsp |
| 60-71 lb | 9-10 yr | 2 1/2 tsp |
| 72-95 lb | 11 yr | 3 tsp |

One Dose Lasts 6-8 Hours
 Store at 20-25°C (68-77°F)
Questions or comments? Call toll-free 1-800-88-ADVIL or ask your pharmacist, doctor or health care professional
 Wyeth Consumer Healthcare, Madison, NJ 07940 Made in USA
 U.S. Patent No. 4,788,220 U0170-30-N2

Children's Advil Suspension
 Fever Reducer/Pain Reliever

Active ingredient (in each 5 mL).....Ibuprofen 100 mg
Uses temporarily:
 ■ reduces fever
 ■ relieves minor aches and pains due to the common cold, flu, sore throat, headaches and toothaches

Fruit Flavored Liquid
 4 FL OZ (120 mL)

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions
 ■ do not give more than directed ■ shake well before using
 ■ find right dose on chart below. If possible, use weight to dose; otherwise use age.
 ■ repeat dose every 6-8 hours, if needed ■ do not use more than 4 times a day
 ■ measure only with the blue dosing cup provided. Blue dosing cup to be used with Children's Advil Suspension only. Do not use with other products. Dose lines account for product remaining in cup due to thickness of suspension.

Dosing Chart

| Weight (lb) | Age (yr) | Dose (tsp) |
|-------------|------------|--------------|
| under 24 lb | under 2 yr | ask a doctor |
| 24-35 lb | 2-3 yr | 1 tsp |
| 36-47 lb | 4-5 yr | 1 1/2 tsp |
| 48-59 lb | 6-8 yr | 2 tsp |
| 60-71 lb | 9-10 yr | 2 1/2 tsp |
| 72-95 lb | 11 yr | 3 tsp |

One Dose Lasts 6-8 Hours
 Store at 20-25°C (68-77°F)
Questions or comments? Call toll-free 1-800-88-ADVIL or ask your pharmacist, doctor or health care professional
 Wyeth Consumer Healthcare, Madison, NJ 07940 Made in USA
 U.S. Patent No. 4,788,220 U0170-30-N2

150%


| DEPARTMENT | REVIEWED BY | | DATE | ✓ with Changes |
|---------------------|-------------|-----------|------|--------------------------|
| | Print Name | Signature | | |
| Graphic Artist | | | | <input type="checkbox"/> |
| Packaging Engineer | | | | <input type="checkbox"/> |
| Labeling Compliance | | | | <input type="checkbox"/> |
| Medical | | | | <input type="checkbox"/> |
| Regulatory Affairs | | | | <input type="checkbox"/> |
| Marketing | | | | <input type="checkbox"/> |
| Legal | | | | <input type="checkbox"/> |
| Trademark | | | | <input type="checkbox"/> |

RELEASED

(1) _____ Date _____

(2) _____ Date _____

DIGITAL MECHANICAL
 COLOR REPRESENTATION MAY BE OFF
 FOLLOW COLOR LEGEND FOR ACTUAL COLOR.



100%

| DRUG FACTS TEXT DEFINED | TYPE SIZE |
|--------------------------------|-----------|
| • HEADINGS | 4.5 pt |
| • SUBHEADS/BODY TEXT | 4.5 pt |
| • LEADING | 5 pt |
| • BULLETS | 3.5 |
| • SPACE AFTER BULLETED SECTION | 2 ems |
| • WARNING BOX LINE, HAIRLINES | .5 |

| Job Information | Color Legend |
|---|---|
| Date 12/18/03 PASS #1 Component # U0174-30-N2 Drawing # 5691 Rev.2 | |
| Identification | Printer Notes |
| Wyeth Consumer Healthcare Production Packaging Dept. Five / Farms Madison, NJ 07940 Phone: (973) 660-5595/5153 Fax: (973) 660-6048 | PRINTER TO REPLACE FPO UPC'S WITH HI-RES DIGITAL ART ALSO PRINTER TO SUPPLY COLOR MATCHPRINT FOR COLOR APPROVAL PRIOR TO PRESS RUN. |
| Vendor Identification | |
| Production Packaging Dept. Cheng Lin SC: 7.15.3 | |

| DEPARTMENT | REVIEWED BY | | DATE | ✓ with Changes |
|---------------------|-------------|-----------|------|--------------------------|
| | Print Name | Signature | | |
| Graphic Artist | | | | <input type="checkbox"/> |
| Packaging Engineer | | | | <input type="checkbox"/> |
| Labeling Compliance | | | | <input type="checkbox"/> |
| Medical | | | | <input type="checkbox"/> |
| Regulatory Affairs | | | | <input type="checkbox"/> |
| Marketing | | | | <input type="checkbox"/> |
| Legal | | | | <input type="checkbox"/> |
| Trademark | | | | <input type="checkbox"/> |

RELEASED

(1) _____ Date _____

(2) _____ Date _____

DIGITAL MECHANICAL
COLOR REPRESENTATION MAY BE OFF
FOLLOW COLOR LEGEND FOR ACTUAL COLOR.

150%

Warnings
Allergy alert: Ibuprofen may cause a severe allergic reaction which may include:
 ■ hives ■ asthma (wheezing) ■ facial swelling ■ shock
Stomach bleeding warning: Taking more than recommended may cause stomach bleeding.
Sore throat warning: Severe or persistent sore throat or sore throat accompanied by high fever, headache, nausea, and vomiting may be serious. Consult doctor promptly. Do not use more than 2 days or administer to children under 3 years of age unless directed by doctor.
Do not use if the child has ever had an allergic reaction to any other fever reducer/pain reliever.
Ask a doctor before use if the child has
 ■ not been drinking fluids ■ ulcers ■ bleeding problems
 ■ lost a lot of fluid due to continued vomiting or diarrhea
 ■ problems or serious side effects from taking fever reducers or pain relievers
 ■ stomach problems that last or come back, such as heartburn, upset stomach or pain
 ■ high blood pressure, heart or kidney disease or is taking a diuretic
Ask a doctor or pharmacist before use if the child is
 ■ under a doctor's care for any serious condition
 ■ taking any other product that contains ibuprofen, or any other pain reliever/fever reducer
 ■ taking a prescription product for anticoagulation (blood thinning)
 ■ taking any other drug
When using this product give with food or milk if upset stomach occurs
Stop use and ask a doctor if
 ■ an allergic reaction occurs. Seek medical help right away.
 ■ fever or pain gets worse or lasts more than 3 days
 ■ the child does not get any relief within first day (24 hours) of treatment
 ■ stomach pain or upset gets worse or lasts
 ■ redness or swelling is present in the painful area
 ■ any new symptoms appear

Do Not Use if plastic wrap imprinted with "CHILDREN'S ADVIL SAFETY SEALED" is broken or missing.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions
 ■ do not give more than directed ■ shake well before using
 ■ find right dose on chart below. If possible, use weight to dose; otherwise use age.
 ■ repeat dose every 6-8 hours, if needed ■ do not use more than 4 times a day
 ■ measure only with the blue dosing cup provided. Blue dosing cup to be used with Children's Advil Suspension only. Do not use with other products. Dose lines account for product remaining in cup due to thickness of suspension.

| Weight (lb) | Age (yr) | Dose (tsp) |
|-------------|------------|--------------|
| under 24 lb | under 2 yr | ask a doctor |
| 24-35 lb | 2-3 yr | 1 tsp |
| 36-47 lb | 4-5 yr | 1 1/2 tsp |
| 48-59 lb | 6-8 yr | 2 tsp |
| 60-71 lb | 9-10 yr | 2 1/2 tsp |
| 72-95 lb | 11 yr | 3 tsp |

One Dose Lasts 6-8 Hours

Store at 20-25°C (68-77°F)
Questions or comments? Call toll-free 1-800-88-ADVIL or ask your pharmacist, doctor or health care professional
 Wyeth Consumer Healthcare, Madison, NJ 07940 Made in USA
 U.S. Patent No. 4,788,220

Children's Advil Suspension
Fever Reducer/Pain Reliever

Active ingredient (in each 5 mL).....Ibuprofen 100 mg
Uses temporarily:
 ■ reduces fever
 ■ relieves minor aches and pains due to the common cold, flu, sore throat, headaches and toothaches

Blue Raspberry Flavored Liquid
4 FL OZ (120 mL)

U0174-30-N2