




PRE-PRINTED TEXT. NOTE: LOT/EXP (OR EQUIVALENT) TO BE PRINTED PER MARKET REQUIREMENTS AT A MINIMUM OF 8 pt. DATE OF MANUFACTURE CAN BE ADDED AS A THIRD LINE IF NEEDED.

UNPRINTED AREA FOR DATE & LOT NUMBER UNVARNISHED

3/16" X 3/16" ARRAY ALPHA NUMERIC DATA MATRIX CODE (ECC 200) PRINTED IN BLACK OR DARK BLUE ON WHITE FIELD WITH WHITE 1/8" QUIET ZONE ON ALL 4 SIDES. DATA MATRIX CODE = 080542

PRODUCTION (NOT COLOR ACCURATE)



DRUG FACTS TEXT DEFINED	TYPE SIZE
• DRUG FACTS TITLE	9 pt
• DRUG FACTS CONTINUED	8 pt
• HEADINGS	8 pt
• SUBHEADINGS/BODY TEXT	6 pt
• LEADING	6.5 pt
• # OF CHARACTERS PER INCH	<39
• BULLETS	5 pt
• SPACE BEFORE BULLET	2 ems
• BARLINES, HAIRLINES	1.5 pt, .5 pt
• SPACE BETWEEN HAIRLINES AND BOX END	2 spaces

090177e189bb1af1\Final\Final On: 27-Sep-2016 17:53

090177e189bb1af5\Final\Final On: 27-Sep-2016 18:02

Do Not Use if breakable ring on bottle cap is separated or missing.

**Children's
Advil®
Suspension**

Ibuprofen Oral Suspension
100 mg per 5 mL

Pain Reliever/Fever Reducer (NSAID)

White Grape-Flavored Liquid
4 FL OZ (120 mL)

Active ingredient (in each 5 mL): Ibuprofen 100 mg

Use temporarily reduce fever, relieve minor aches and pains due to the common cold, flu, sore throat, headaches and toothaches

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: hives, facial swelling, asthma, wheezing, shock, skin redness, rash, blisters. If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warnings: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if the child has had stomach ulcers or bleeding problems, takes a blood thinner (warfarin/coumadin) or stored drug, takes other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) takes more or for a longer time than directed.

Heart attack and stroke warnings: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Sore throat warning: Severe or persistent sore throat or sore throat accompanied by high fever, headache, nausea, and vomiting may be serious. Consult a doctor promptly. Do not use more than 2 days or administer to children under 3 years of age unless directed by a doctor.

Do not use if the child has ever had an allergic reaction to any other fever reducer/pain reliever right before or after heart surgery.

Ask a doctor before use if stomach bleeding warning applies to the child; child has a history of stomach problems, such as heartburn; child has high blood pressure, heart disease, liver disease, kidney disease, asthma, or had a stroke; child has not been drinking fluids; child has lost a lot of fluid due to vomiting or diarrhea; child is taking a diuretic.

Ask a doctor or pharmacist before use if the child is under a doctor's care for any serious condition; taking any other drug.

When using this product take with food or milk if upset stomach occurs

Stop use and ask a doctor if child experiences any of the following signs of stomach bleeding: feels faint, vomits blood, has bloody or black stools, or has stomach pain that does not get better; child has symptoms of heart problems or stroke: chest pain, trouble breathing, weakness in one part or side of body, slurred speech, leg weakness; the child does not get any relief within first day (24 hours) of treatment; fever or pain gets worse or lasts more than 3 days; redness or swelling is present in the painful area; any new symptoms appear.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Each 5 mL contains: sodium 3 mg. Store at 20-25°C (68-77°F).

Directions: This product does not contain directions or complete warnings for adult use. Do not give more than directed. Shake well before using. mL = milliliter. Find right dose on chart below. If possible, use weight to dose, otherwise use age. Repeat dose every 6-8 hours, if needed.

Do not use more than 4 times a day. Measure only with the dosing cap provided. Dosing cap to be used with Children's Advil Suspension only. Do not use with other products. These lines account for product remaining in cap due to thickness of suspension.


Dosing Chart		
Weight (lb)	Age (yr)	Dose (mL)
under 24 lb	under 2 yr	ask a doctor
24-35 lb	2-3 yr	5 mL
36-47 lb	4-5 yr	7.5 mL
48-59 lb	6-8 yr	10 mL
60-71 lb	9-10 yr	12.5 mL
72-95 lb	11 yr	15 mL

Questions or comments? call 1-800-88-ADAIL.
 Pfizer, Madison, NJ 07940 USA © 2016 Pfizer Inc.
 For most recent product information, visit www.Advil.com PAA080543

3/16" X 3/16" DATA MATRIX CODE (ECC200),
 PRINTED IN BLACK OR DARK BLUE ON UNPRINTED
 WHITE BACKGROUND WITH A MINIMUM OF 1/8"
 QUIET ZONE ALL 4 SIDES.
DATA MATRIX CODE = 080543

UNVARNISHED
 UNPRINTED AREA
 FOR LOT & EXP.

PRODUCTION (NOT COLOR ACCURATE)



DRUG FACTS TEXT DEFINED	TYPE SIZE
• HEADINGS	4.5 pt
• SUBHEADINGS/BODY TEXT	4.5 pt
• LEADING	4.57 pt
• BULLETS	N/A
• SPACE BEFORE BULLET	N/A
• WARNING BOX LINE, HAIRLINES	.5 pt

Children's Advil[®] Suspension

Packaged with Tamper-Evident bottle cap. Do Not Use if breakable ring is separated or missing.

Children's Advil[®] Suspension

Children's Advil[®] Suspension

Ibuprofen Oral Suspension
100 mg per 5 mL
Pain Reliever/Fever Reducer (NSAID)

Fever
Aches & Pains
Lasts Up To 8 Hours

Grape-Flavored Liquid
for ages 2-11 years

4 FL OZ (120 mL) Alcohol-Free

Drug Facts (continued)

- child has not been drinking fluids
- child has lost a lot of fluid due to vomiting or diarrhea
- child is taking a diuretic

Ask a doctor or pharmacist before use if the child is

- under a doctor's care for any serious condition
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- child experiences any of the following signs of stomach bleeding:
 - feels faint
 - vomits blood
 - has bloody or black stools
 - has stomach pain that does not get better
- child has symptoms of heart problems or stroke:
 - chest pain
 - trouble breathing
 - weakness in one part or side of body
 - slurred speech
 - leg swelling
- the child does not get any relief within first day (24 hours) of treatment
- fever or pain gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- this product does not contain directions or complete warnings for adult use
- do not give more than directed
- shake well before using
- mL = milliliter
- find right dose on chart. If possible, use weight to dose; otherwise use age.
- repeat dose every 6-8 hours, if needed
- do not use more than 4 times a day
- measure only with the dosing cup provided. Dosing cup to be used with Children's Advil Suspension only. Do not use with other products. Dose lines account for product remaining in cup due to thickness of suspension.

Weight (lb)	Age (yr)	Dose (mL)
under 24 lb	under 2 yr	ask a doctor
24-35 lb	2-3 yr	5 mL
36-47 lb	4-5 yr	7.5 mL
48-59 lb	6-8 yr	10 mL
60-71 lb	9-10 yr	12.5 mL
72-95 lb	11 yr	15 mL

Drug Facts (continued)

Other information

- each 5 mL contains: sodium 3 mg
- one dose lasts 6-8 hours
- store at 20-25°C (68-77°F)
- see bottom of box for lot number and expiration date

Inactive ingredients

acetic acid, artificial flavor, butylated hydroxytoluene, carboxymethylcellulose sodium, citric acid monohydrate, edetate disodium, FD&C blue no. 1, FD&C red no. 40, glycerin, microcrystalline cellulose, polysorbate 80, propylene glycol, purified water, sodium benzoate, sorbitol solution, sucrose, xanthan gum

Questions or comments?
call toll-free 1-800-88-ADVIL or ask your pharmacist, doctor or health care professional

Long Lasting: Up to 8 hours of fever reduction
Great Convenience: Long duration means fewer doses needed per day.
Alcohol-Free

For most recent product information, visit www.Advil.com
Pfizer, Madison, NJ 07940 USA © 2016 Pfizer Inc.

F.P.O.

3 0573 0171 30 4

PAA082055

LOT:
EXP:

PRE-PRINTED TEXT. NOTE: LOT/EXP (OR EQUIVALENT) TO BE PRINTED PER MARKET REQUIREMENTS AT A MINIMUM OF 8 pt. DATE OF MANUFACTURE CAN BE ADDED AS A THIRD LINE IF NEEDED.


UNPRINTED AREA FOR DATE & LOT NUMBER UNVARNISHED

3/16" X 3/16" ARRAY ALPHA NUMERIC DATA MATRIX CODE (ECC 200) PRINTED IN BLACK OR DARK BLUE ON WHITE FIELD WITH WHITE 1/8" QUIET ZONE ON ALL 4 SIDES.
DATA MATRIX CODE = 082055

PRODUCTION (NOT COLOR ACCURATE) (b) (4)

DRUG FACTS TEXT DEFINED	TYPE SIZE
• DRUG FACTS TITLE	9 pt
• DRUG FACTS CONTINUED	8 pt
• HEADINGS	8 pt
• SUBHEADINGS/BODY TEXT	6 pt
• LEADING	6.5 pt
• # OF CHARACTERS PER INCH	<39
• BULLETS	5 pt
• SPACE BEFORE BULLET	2 ems
• BARLINES, HAIRLINES	1.5 pt, .5 pt
• SPACE BETWEEN HAIRLINES AND BOX END	2 spaces

090177e18ac656de\Final\Final On: 08-Dec-2016 04:04




3/16" X 3/16" DATA MATRIX CODE (ECC 200),
 PRINTED IN BLACK OR DARK BLUE ON
 UNPRINTED WHITE BACKGROUND WITH A
 MINIMUM OF 1/8" QUIET ZONE ALL 4 SIDES.
DATA MATRIX CODE = 082056

UNVARNISHED
 UNPRINTED AREA
 FOR LOT & EXP.

PRODUCTION

(NOT COLOR ACCURATE)

(b) (4)



DRUG FACTS TEXT DEFINED	TYPE SIZE
• HEADINGS	4.5 pt
• SUBHEADINGS/BODY TEXT	4.5 pt
• LEADING	4.6 pt
• BULLETS	N/A
• SPACE BEFORE BULLET	N/A
• WARNING BOX LINE, HAIRLINES	.5 pt

Children's Advil[®] Suspension

Packaged with Tamper-Evident bottle cap. Do Not Use if breakable ring is separated or missing.

SUGAR-FREE

Children's Advil[®] Suspension

Ibuprofen Oral Suspension
100 mg per 5 mL
Pain Reliever/Fever Reducer (NSAID)

Fever
Aches & Pains
Lasts Up To **8** Hours

DYE-FREE BERRY
Berry-Flavored Liquid for ages 2-11 years

4 FL OZ (120 mL) Alcohol-Free

Drug Facts (continued)

- child has not been drinking fluids
- child has lost a lot of fluid due to vomiting or diarrhea
- child is taking a diuretic

Ask a doctor or pharmacist before use if the child is

- under a doctor's care for any serious condition
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- child experiences any of the following signs of stomach bleeding:
 - feels faint
 - vomits blood
 - has bloody or black stools
 - has stomach pain that does not get better
- child has symptoms of heart problems or stroke:
 - chest pain
 - trouble breathing
 - weakness in one part or side of body
 - slurred speech
 - leg swelling
- the child does not get any relief within first day (24 hours) of treatment
- fever or pain gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- this product does not contain directions or complete warnings for adult use
- do not give more than directed
- shake well before using
- mL = milliliter
- find right dose on chart. If possible, use weight to dose; otherwise use age.
- repeat dose every 6-8 hours, if needed
- do not use more than 4 times a day
- measure only with the dosing cup provided. Dosing cup to be used with Children's Advil Suspension only. Do not use with other products. Dose lines account for product remaining in cup due to thickness of suspension.

Dosing Chart		
Weight (lb)	Age (yr)	Dose (mL)
under 24 lb	under 2 yr	ask a doctor
24-35 lb	2-3 yr	5 mL
36-47 lb	4-5 yr	7.5 mL
48-59 lb	6-8 yr	10 mL
60-71 lb	9-10 yr	12.5 mL
72-85 lb	11 yr	15 mL

Other information

- each 5 mL contains: sodium 10 mg
- one dose lasts 6-8 hours
- store at 20-25°C (68-77°F)
- see bottom of box for lot number and expiration date

Inactive ingredients

artificial flavor, carboxymethylcellulose sodium, citric acid monohydrate, edetate disodium, glycerin, microcrystalline cellulose, polysorbate 80, propylene glycol, purified water, sodium benzoate, sodium citrate, sorbitol solution, sucralose, xanthan gum

Questions or comments?
call toll-free 1-800-88-ADVIL or ask your pharmacist, doctor or health care professional

Long Lasting: Up to 8 hours of fever reduction
Great Convenience: Long duration means fewer doses needed per day.
Alcohol-Free

For most recent product information, visit www.Advill.com
Pfizer, Madison, NJ 07940 USA © 2016 Pfizer Inc.

Important: Read all product information before using. Keep this box for important information. This product is intended for use in children ages 2 years - 11 years.

Drug Facts

Active ingredient (in each 5 mL) Purpose
Ibuprofen 100 mg (NSAID)* ...Fever reducer/Pain reliever
*nonsteroidal anti-inflammatory drug

Uses temporarily:

- reduces fever
- relieves minor aches and pains due to the common cold, flu, sore throat, headaches and toothaches

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if the child

- has had stomach ulcers or bleeding problems
- takes a blood thinning (anticoagulant) or steroid drug
- takes other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]
- takes more or for a longer time than directed

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Sore throat warning: Severe or persistent sore throat or sore throat accompanied by high fever, headache, nausea, and vomiting may be serious. Consult doctor promptly. Do not use more than 2 days or administer to children under 3 years of age unless directed by doctor.

Do not use

- if the child has ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- stomach bleeding warning applies to the child
- child has problems or serious side effects from taking pain relievers or fever reducers
- child has a history of stomach problems, such as heartburn
- child has high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke

GLUE AREA UNVARNISHED

PAA081707

LOT:
EXP:

PRE-PRINTED TEXT. NOTE: LOT/EXP (OR EQUIVALENT) TO BE PRINTED PER MARKET REQUIREMENTS AT A MINIMUM OF 8 pt. DATE OF MANUFACTURE CAN BE ADDED AS A THIRD LINE IF NEEDED.

UNPRINTED AREA FOR DATE & LOT NUMBER UNVARNISHED

3/16" X 3/16" ARRAY ALPHA NUMERIC DATA MATRIX CODE (ECC 200) PRINTED IN BLACK OR DARK BLUE ON WHITE FIELD WITH WHITE 1/8" QUIET ZONE ON ALL 4 SIDES.
DATA MATRIX CODE = 081707


PRODUCTION (NOT COLOR ACCURATE)

(b) (4)

DRUG FACTS TEXT DEFINED	TYPE SIZE
• DRUG FACTS TITLE	9 pt
• DRUG FACTS CONTINUED	8 pt
• HEADINGS	8 pt
• SUBHEADINGS/BODY TEXT	6 pt
• LEADING	6.5 pt
• # OF CHARACTERS PER INCH	<39
• BULLETS	5 pt
• SPACE BEFORE BULLET	2 ems
• BARLINES, HAIRLINES	1.5 pt, .5 pt
• SPACE BETWEEN HAIRLINES AND BOX END	2 spaces

090177e18ac63d8e\Final\Final On: 08-Dec-2016 04:00

090177e18ac64b01Final\Final On: 08-Dec-2016 04:00




3/16" X 3/16" DATA MATRIX CODE (ECC 200),
 PRINTED IN BLACK OR DARK BLUE ON
 UNPRINTED WHITE BACKGROUND WITH A
 MINIMUM OF 1/8" QUIET ZONE ALL 4 SIDES.
DATA MATRIX CODE = 081708

UNVARNISHED
 UNPRINTED AREA
 FOR LOT & EXP.

PRODUCTION

(NOT COLOR ACCURATE)

(b) (4)



DRUG FACTS TEXT DEFINED	TYPE SIZE
• HEADINGS	4.5 pt
• SUBHEADINGS/BODY TEXT	4.5 pt
• LEADING	4.6 pt
• BULLETS	N/A
• SPACE BEFORE BULLET	N/A
• WARNING BOX LINE, HAIRLINES	.5 pt

PRE-PRINTED TEXT. NOTE: LOT/EXP (OR EQUIVALENT) TO BE PRINTED PER MARKET REQUIREMENTS AT A MINIMUM OF 8 pt. DATE OF MANUFACTURE CAN BE ADDED AS A THIRD LINE IF NEEDED.


UNRRRNIEEDARBEA/F0BR
DATE&LOTNUMBER
UNRRRISBBBD

3/16" X 3/16" ARRAY ALPHA NUMERIC DATA MATRIX CODE (ECC 200) PRINTED IN BLACK OR DARK BLUE ON WHITE FIELD WITH WHITE 1/8" QUIET ZONE ON ALL 4 SIDES. DATA MATRIX CODE = 082853

PRODUCTION (NOT COLOR ACCURATE) (b) (4)

DRUG FACTS TEXT DEFINED	TYPE SIZE
• DRUG FACTS TITLE	9 pt
• DRUG FACTS CONTINUED	8 pt
• HEADINGS	8 pt
• SUBHEADINGS/BODY TEXT	6 pt
• LEADING	6.5 pt
• # OF CHARACTERS PER INCH	<39
• BULLETS	5 pt
• SPACE BEFORE BULLET	2 ems
• BARLINES, HAIRLINES	1.5 pt, .5 pt
• SPACE BETWEEN HAIRLINES AND BOX END	2 spaces

090177e18ac64b04\Final\Final On: 08-Dec-2016 04:05



3/16" X 3/16" DATA MATRIX CODE (ECC 200),
 PRINTED IN BLACK OR DARK BLUE ON
 UNPRINTED WHITE BACKGROUND WITH A
 MINIMUM OF 1/8" QUIET ZONE ALL 4 SIDES.
DATA MATRIX CODE = 082054


UNVARNISHED
 UNPRINTED AREA
 FOR LOT & EXP.

PRODUCTION

(NOT COLOR ACCURATE)

(b) (4)

DRUG FACTS TEXT DEFINED	TYPE SIZE
• HEADINGS	4.5 pt
• SUBHEADINGS/BODY TEXT	4.5 pt
• LEADING	4.6 pt
• BULLETS	N/A
• SPACE BEFORE BULLET	N/A
• WARNING BOX LINE, HAIRLINES	.5 pt




090177e18ac64b05\Final\Final On: 08-Dec-2016 04:04

PRODUCTION (NOT COLOR ACCURATE) (b) (4)

DRUG FACTS TEXT DEFINED	TYPE SIZE
• DRUG FACTS TITLE	9 pt
• DRUG FACTS CONTINUED	8 pt
• HEADINGS	8 pt
• SUBHEADINGS/BODY TEXT	6 pt
• LEADING	6.5 pt
• # OF CHARACTERS PER INCH	<39
• BULLETS	5 pt
• SPACE BEFORE BULLET	2 ems
• BARLINES, HAIRLINES	1.5 pt, .5 pt
• SPACE BETWEEN HAIRLINES AND BOX END	2 spaces

090177e18ac64b06\Final\Final On: 08-Dec-2016 04:03



3/16" X 3/16" DATA MATRIX CODE (ECC200),
 PRINTED IN BLACK OR DARK BLUE ON
 UNPRINTED WHITE BACKGROUND WITH A
 MINIMUM OF 1/8" QUIET ZONE ALL 4 SIDES.
DATA MATRIX CODE = 082590

UNVARNISHED
 UNPRINTED AREA
 FOR LOT & EXP.


PRODUCTION

(NOT COLOR ACCURATE)

(b) (4)




DRUG FACTS TEXT DEFINED	TYPE SIZE
• HEADINGS	4.5 pt
• SUBHEADINGS/BODY TEXT	4.5 pt
• LEADING	4.6 pt
• BULLETS	N/A
• SPACE BEFORE BULLET	N/A
• WARNING BOX LINE, HAIRLINES	.5 pt



PRODUCTION (NOT COLOR ACCURATE) (b) (4)

DRUG FACTS TEXT DEFINED	TYPE SIZE
• DRUG FACTS TITLE	9 pt
• DRUG FACTS CONTINUED	8 pt
• HEADINGS	8 pt
• SUBHEADINGS/BODY TEXT	6 pt
• LEADING	6.5 pt
• # OF CHARACTERS PER INCH	<39
• BULLETS	5 pt
• SPACE BEFORE BULLET	2 ems
• BARLINES, HAIRLINES	1.5 pt, .5 pt
• SPACE BETWEEN HAIRLINES AND BOX END	2 spaces




3/16" X 3/16" DATA MATRIX CODE (ECC 200),
 PRINTED IN BLACK OR DARK BLUE ON
 UNPRINTED WHITE BACKGROUND WITH A
 MINIMUM OF 1/8" QUIET ZONE ALL 4 SIDES.
DATA MATRIX CODE = 082585

UNVARNISHED
 UNPRINTED AREA
 FOR LOT & EXP.

PRODUCTION

(NOT COLOR ACCURATE)

(b) (4)



DRUG FACTS TEXT DEFINED	TYPE SIZE
• HEADINGS	4.5 pt
• SUBHEADINGS/BODY TEXT	4.5 pt
• LEADING	4.6 pt
• BULLETS	N/A
• SPACE BEFORE BULLET	N/A
• WARNING BOX LINE, HAIRLINES	.5 pt

This is a representation of an electronic record that was signed electronically and this page is the manifestation of the electronic signature.

/s/

VALERIE S PRATT
03/23/2017