



PAA166901

DRUG FACTS TEXT DEFINED	TYPE SIZE
• DRUG FACTS TITLE	9 pt
• DRUG FACTS CONTINUED	8 pt
• HEADINGS	8 pt
• SUBHEADINGS/BODY TEXT	6 pt
• LEADING	6.5 pt
• # OF CHARACTERS PER INCH	<39
• BULLETS	5 pt
• SPACE BEFORE BULLET	2 ems
• BARLINES, HAIRLINES	1.5 pt, .5 pt
• SPACE BETWEEN HAIRLINES AND BOX END	2 spaces

Base Label Page 5

Datos sobre el fármaco (continuación)

Cuadro de dosis		
Peso (libras)	Edad (años)	Dosis (mL)
menos de 24 libras	menos de 2 años	consulte a un médico
24 a 35 libras	2 a 3 años	5 mL
36 a 47 libras	4 a 5 años	7.5 mL
48 a 59 libras	6 a 8 años	10 mL
60 a 71 libras	9 a 10 años	12.5 mL
72 a 95 libras	11 años	15 mL

Información adicional

- cada 5 mL contienen: **3 mg de sodio**
- una dosis dura entre 6 y 8 horas
- consérvelo entre 20 y 25 °C (68 a 77 °F)
- consulte el número de lote y la fecha de vencimiento en la base de la caja

Ingredientes inactivos
saborizante artificial, carboximetilcelulosa sódica, ácido cítrico monohidratado, edetato disódico, FD&C rojo n.º 40, glicerina, celulosa microcristalina, polisorbato 80, agua purificada, benzoato de sodio, solución de sorbitol, sacarosa, goma xantana

¿Preguntas o comentarios?
llame gratuitamente al **1-800-88-ADVIL** o consulte a su farmacéutico, médico o profesional de atención de la salud

Base Label Page 5

Insert Back Page 2

Datos sobre el fármaco (continuación)

Advertencias

Alerta de alergia: El ibuprofeno puede provocar una reacción alérgica intensa, especialmente en personas alérgicas a la aspirina. Los síntomas pueden incluir:

- urticaria
- asma (sibilancia)
- enrojecimiento de la piel
- erupción cutánea
- hinchazón facial
- choque
- ampollas.

Si se produce una reacción alérgica, interrumpa su uso y busque ayuda médica de inmediato.

Advertencia de sangrado estomacal: Este producto contiene un AINE, que puede provocar sangrado estomacal intenso. Las probabilidades son mayores si el niño:

- ha tenido problemas de úlceras o sangrado en el estómago
- usa un anticoagulante o un corticoesteroide
- usa otros fármacos de venta con o sin receta que contienen AINE [aspirina, ibuprofeno, naproxeno u otros]
- toma este producto en dosis mayores o durante un período más prolongado que lo indicado.

Advertencia de ataque cardíaco y accidente cerebrovascular: Los AINE, exceptuando la aspirina, aumentan el riesgo de ataque cardíaco, infarto y accidentes cerebrovasculares. Estos pueden ser fatales. El riesgo es más alto si usted usa más de lo que se receta, o por un período más largo del que se receta.

Advertencia de dolor de garganta: El dolor de garganta intenso o persistente, o dolor de garganta acompañado de fiebre alta, dolor de cabeza, náuseas y vómitos pueden ser graves. Consulte a un médico sin demora. No use este producto durante más de 2 días ni lo administre a niños menores de 3 años a menos que lo indique un médico.

Datos sobre el fármaco (continuación)

No lo use

- si el niño ha presentado alguna vez una reacción alérgica a cualquier otro analgésico-antifebril
- inmediatamente antes o después de una cirugía de corazón.

Consulte a un médico antes de usarlo si

- la advertencia de sangrado estomacal es aplicable al niño
- el niño tiene problemas o efectos secundarios graves por tomar analgésicos o antifebriles
- el niño tiene antecedentes de problemas estomacales, como acidez;
- el niño tiene presión arterial alta, enfermedad cardíaca, cirrosis hepática, enfermedad renal, asma, o tuvo un accidente cerebrovascular;
- el niño no ha estado tomando líquidos;
- el niño ha perdido gran cantidad de líquido debido a vómitos o diarrea,
- el niño está tomando un diurético.

Consulte a un médico o farmacéutico antes de usarlo si el niño

- recibe cuidados médicos por cualquier afección grave
- está usando algún otro fármaco.

Al usar este producto

- adminístrelo con alimentos o leche si se produce malestar estomacal

Interrumpa su uso y consulte a un médico si

- el niño presenta alguno de los siguientes signos de sangrado estomacal:
 - siente que va a desmayarse,
 - vomita sangre
 - tiene heces con sangre o de color negro
 - tiene dolor estomacal que no mejora

Insert Front Page 4

Datos sobre el fármaco (continuación)

- niño(a) tiene síntomas de problemas cardíacos o accidente cerebrovascular:
- dolor de pecho
- debilidad en una parte o lado del cuerpo
- habla ininteligible o afectada
- hinchazón de piernas
- el niño no presenta ninguna mejoría dentro del primer día (24 horas) de tratamiento;
- la fiebre o el dolor empeoran o duran más de 3 días
- se presenta enrojecimiento o hinchazón en el área dolorida
- aparece algún síntoma nuevo.

Manténgalo fuera del alcance de los niños. En caso de sobredosis, obtenga ayuda médica o comuníquese con un centro de toxicología de inmediato.

Instrucciones

- este producto no contiene instrucciones ni advertencias completas para su uso en adultos
- no administre cantidades mayores de las indicadas
- agite bien el frasco antes de usarlo
- mL = mililitro
- consulte la dosis correcta en el cuadro. Si es posible, use el peso como referencia para la dosis; como alternativa, use la edad.
- si es necesario, repita la dosis cada **6 a 8 horas**
- no lo use más de **4 veces por día**
- mida únicamente con el vaso dosificador suministrado. El vaso dosificador debe utilizarse exclusivamente con Children's Advil Suspension/Suspension. No lo use con otros productos. Las líneas que marcan las dosis toman en cuenta el producto que queda adherido al vaso debido a la viscosidad de la suspensión.

Drug Facts (continued)

Other information

- each 5 mL contains: **sodium 3 mg**
- one dose lasts 6-8 hours
- store at 20-25°C (68-77°F)
- see bottom of box for lot number and expiration date

Inactive ingredients
artificial flavor, carboxymethylcellulose sodium, citric acid monohydrate, edetate disodium, FD&C red no. 40, glycerin, microcrystalline cellulose, polysorbate 80, purified water, sodium benzoate, sorbitol solution, sucrose, xanthan gum

Questions or comments?
call toll-free **1-800-88-ADVIL** or ask your pharmacist, doctor or health care professional

Importante: Lea toda la información sobre el producto antes de usarlo. Conserve esta caja para información importante. Este producto está pensado para ser administrado a niños de 2 a 11 años.

Datos sobre el fármaco

Ingrediente activo (por cada 5 mL)	Finalidad
ibuprofeno 100 mg (AINE)*	Antifebril/analgésico
	*antiinflamatorio no esteroideo

Usos: temporalmente: ■ baja la fiebre

- alivia dolores y malestares leves causados por el resfriado común, la gripe, el dolor de garganta, el dolor de cabeza y el dolor de muelas

For Drug Facts in Spanish, LIFT HERE
Para obtener datos sobre el fármaco en español, LEVANTE AQUÍ

PAA166902

Insert Front Page 1

PAA166902

DRUG FACTS TEXT DEFINED	TYPE SIZE
• DRUG FACTS TITLE	9 pt
• DRUG FACTS CONTINUED	8 pt
• HEADINGS	8 pt
• SUBHEADINGS/BODY TEXT	6 pt
• LEADING	6.5 pt
• # OF CHARACTERS PER INCH	<39
• BULLETS	5 pt
• SPACE BEFORE BULLET	2 ems
• BARLINES, HAIRLINES	1.5 pt, .5 pt
• SPACE BETWEEN HAIRLINES AND BOX END	2 spaces

TOP LABEL

**Children's
Advil**
Suspension/Suspensión
Ibuprofen Oral Suspension
100 mg per 5 mL
Pain Reliever/Fever Reducer (NSAID)
Ibuprofeno suspensión oral
100 mg por cada 5 mL
Analgésico/antifebril (AINE)

**Fruit-Flavored Liquid
Líquido con sabor a fruta
4 FL OZ (120 mL)**

Do Not Use if plastic wrap imprinted with "SAFETY SEAL" is broken or missing. No lo use si el empaque plástico con la leyenda "SAFETY SEAL" está roto o falta.

Active ingredient
(for each 5 mL).....Ibuprofen 100 mg
Use: temporarily reduces fever, relieves minor aches and pains due to the common cold, flu, sore throat, headache and toothaches

Warnings
Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: hives, facial swelling, asthma (wheezing), shock, skin reddening, rash, h itams. If an allergic reaction occurs, stop use and seek medical help right away.
Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if the child has had stomach ulcers or bleeding problems, takes a blood thinning (anticoagulant) or steroid drug, takes other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others), takes more or for a longer time than directed.

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed. **Sore throat warning:** Severe or persistent sore throat or sore throat accompanied by high fever, headache, nausea, and vomiting may be serious. Consult doctor promptly. Do not use more than 2 days or administer to children under 3 years of age unless directed by doctor.
Do not use if the child has ever had an allergic reaction to any other pain reliever/fever reducer, right before or after heart surgery.
Ask a doctor before use if stomach bleeding warning applies to the child, child has a history of stomach problems, such as heartburn; child has high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke; child has not been drinking fluids; child has lost a lot of fluid due to vomiting or diarrhea; child is taking a diuretic.
Ask a doctor or pharmacist before use if the child is under a doctor's care for any serious condition, taking any other drug.
When using this product take with food or milk if stomach upset occurs.

Stop use and ask a doctor if child experiences any of the following signs of stomach bleeding: feels faint, vomits blood, has bloody or black stools, or has stomach pain that does not get better; child has symptoms of heart problems or stroke: chest pain, trouble breathing, weakness in one part or side of body, slurred speech, leg swelling, the child does not get any relief within first day (24 hours) of treatment; fever or pain gets worse or lasts more than 3 days; redness or swelling is present in the painful area; any new symptoms appear.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Each 5 mL contains acetaminophen 3 mg. Store at 20-25° C (68-77° F).

PAA166835

BASE LABEL

Directions This product does not contain directions or complete warnings for adult use. Do not give more than directed. Shake well before using. mL = milliliter. Find right dose on chart. If possible, use weight to dose; otherwise use age. Repeat dose every 6-8 hours, if needed. Do not use more than 4 times a day. Measure only with the dosing cup provided. Dosing cup to be used with Children's Advil Suspension only. Do not use with other products. Dose lines account for product remaining in cup due to thickness of suspension.

Dosing Chart

Weight (lb)	Age (yr)	Dose (mL)
Under 24 lb	Under 2 yr	Ask a doctor
24-35 lb	2-3 yr	5 mL
36-47 lb	4-5 yr	7.5 mL
48-59 lb	6-8 yr	10 mL
60-71 lb	9-10 yr	12.5 mL
72-85 lb	11 yr	15 mL

Questions or comments? Call 1-800-88-ADWIL. Distributed by: GSK Consumer Healthcare, Warren, NJ 07059. For most recent product information, visit www.Advil.com.

Ingredientes activos
(por cada 5 mL).....Ibuprofeno 100 mg
Usar: baja temporalmente la fiebre y alivia dolores y molestias leves causados por el resaca común, la gripe, el dolor de garganta, el dolor de cabeza y el dolor de muelas.

Advertencias
Alerta de alergia: El ibuprofeno puede provocar una reacción alérgica intensa, especialmente en personas alérgicas a la aspirina. Los síntomas pueden incluir: eritema, hinchazón facial, asma (sibilancias), choque anafiláctico, erupción cutánea y ampollas. Si se produce una reacción alérgica, interrumpa su uso y busque ayuda médica de inmediato.
Advertencia de sangrado estomacal: Este producto contiene un AINE, que puede provocar sangrado estomacal intenso. Las probabilidades son mayores si el niño(a) ha tenido problemas de úlcera o sangrado en el estómago, usa un anticoagulante o un corticosteroide, usa otros fármacos de venta con o sin receta que contienen AINES (aspirina, ibuprofeno, naproxeno u otros), toma este producto en dosis mayores o durante un periodo más prolongado que lo indicado.
Advertencia de Ataque Cardíaco y Accidente Cerebrovascular: Los AINES, exceptando la aspirina, aumentan el riesgo de ataque cardíaco, infarto y accidentes cerebrovasculares. Estos pueden ser fatales. El riesgo es más alto si usted usa más de lo que se receta, o por un periodo más largo del que se receta.
Advertencia de dolor de garganta: El dolor de garganta intenso o persistente, o dolor de garganta acompañado de fiebre alta, dolor de cabeza, náuseas y vómitos pueden ser graves. Consulte a un médico sin demora. No use este producto durante más de 2 días ni lo administre a niños menores de 3 años a menos que lo indique un médico.
No lo use si el niño ha presentado alguna vez una reacción alérgica a cualquier otro analgésico/antifebril; inmediatamente antes o después de una cirugía de corazón.
Consulte a un médico antes de usarlo si la advertencia de sangrado estomacal es aplicable al niño; el niño tiene antecedentes de problemas estomacales, como acidez; el niño tiene presión arterial alta, enfermedad cardíaca, cirrosis hepática, enfermedad renal, asma, o tuvo un accidente cerebrovascular; el niño no ha estado tomando líquidos; el niño ha perdido gran cantidad de líquido debido a vómitos o diarrea, o el niño está tomando un diurético.
Consulte a un médico o farmacéutico antes de usarlo si el niño recibe cuidados médicos por cualquier afección grave; si está usando algún otro fármaco.

INSIDE TOP LABEL

Al usar este producto, adminístrelo o con alimentos o leche si se produce malestar estomacal. Interrumpa su uso y consulte a un médico si el niño presenta alguno de los siguientes signos de sangrado estomacal: siente que va a desmayarse, vomita sangre, tiene heces con sangre o de color negro, o tiene dolor estomacal que no mejora; niño(a) tiene síntomas de problemas cardíacos o accidente cerebrovascular: dolor de pecho, problemas para respirar; hinchazón de piernas: el niño no presenta ninguna mejoría dentro del primer día (24 horas) de tratamiento; la fiebre o el dolor empeoran o duran más de 3 días; se presenta enrojecimiento o hinchazón en el área del dolor; o aparecen algún síntoma nuevo.
Manténgalo fuera del alcance de los niños. En caso de sobredosis, obtenga ayuda médica o comuníquese con un centro de toxicología de inmediato.
Cada 5 mL contienen 3 mg de sodio.
Consérvelo entre 20 y 25° C (68 a 77° F).
Instrucciones: Este producto no contiene instrucciones ni advertencias completas para su uso en adultos.
No administre cantidades mayores de las indicadas.

Agita bien el frasco antes de usarlo. mL = mililitro. Consulte la dosis correcta en el cuadro. Si es posible, use el peso como referencia para la dosis; como alternativa, use la edad. Si es necesario, repita la dosis cada 6 a 8 horas. No lo use más de 4 veces por día. Mida únicamente con el vaso dosificador suministrado. El vaso dosificador debe utilizarse exclusivamente con Children's Advil Suspension. No lo use con otros productos. Las líneas que marcan las dosis toman en cuenta el producto que queda adherido al vaso debido a la viscosidad de la suspensión.

Peso (libras)	Edad (años)	Dosis (mL)
menos de 24 libras	menos de 2 años	consulta a un médico
24 a 35 libras	2 a 3 años	5 mL
36 a 47 libras	4 a 5 años	7.5 mL
48 a 59 libras	6 a 8 años	10 mL
60 a 71 libras	9 a 10 años	12.5 mL
72 a 85 libras	11 años	15 mL

¿Preguntas o comentarios? Llame al 1-800-88-ADWIL. Distribuido por: GSK Consumer Healthcare, Warren, NJ 07059. Para ver la información más reciente sobre el producto, visite www.Advil.com.

PAA166835

DRUG FACTS TEXT DEFINED	TYPE SIZE
• HEADINGS	4.5 pt
• SUBHEADINGS/BODY TEXT	4.5 pt
• LEADING	4.75 pt
• BULLETS	N/A
• SPACE BEFORE BULLET	N/A
• WARNING BOX LINE, HAIRLINES	.5 pt



PAA166837

NON MODIFIED DRUG FACTS TEXT DEFINED	TYPE SIZE
• DRUG FACTS TITLE	9 pt
• DRUG FACTS CONTINUED	8 pt
• HEADINGS	8 pt
• SUBHEADINGS/BODY TEXT	6 pt
• LEADING	6.5 pt
• # OF CHARACTERS PER INCH	<39
• BULLETS	5 pt
• SPACE BEFORE BULLET	2 ems
• BARLINES, HAIRLINES	1.5 pt, .5 pt
• SPACE BETWEEN HAIRLINES AND BOX END	2 spaces

Do Not Use if plastic wrap imprinted with "SAFETY SEAL" is broken or missing.

Children's Advil Suspension
 Ibuprofen Oral Suspension
 100 mg per 5 mL
 Pain Reliever/Fever Reducer (NSAID)
 Grape-Flavored Liquid
 4 FL OZ (120 mL)

Active ingredient (in each 5 mL).....Ibuprofen 100 mg
 (Ibuprofen temporarily reduces fever, relieves minor aches and pains due to the common cold, flu, sore throat, headaches and toothaches)

Warnings
Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include hives, facial swelling, asthma/wheezing, shock, skin redness, rash, dizziness. If an allergic reaction occurs, stop use and seek medical help right away.
Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if the child has had stomach ulcers or bleeding problems, taken a blood thinning (anticoagulant) or steroid drug, takes other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others), takes more or for a longer time than directed.
Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.
Sore throat warning: Severe or persistent sore throat or sore throat accompanied by high fever, headache, nausea, and vomiting may be serious. Consult a doctor promptly. Do not use more than 2 days or administer to children under 3 years of age unless directed by a doctor. Do not use if the child has ever had an allergic reaction to any other fever reducer/pain reliever, right before or after heart surgery.
Ask a doctor before use if stomach bleeding warning applies to the child: child has a history of stomach problems, such as heartburn; child has high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke; child has not been drinking fluids; child has lost a lot of fluid due to vomiting or diarrhea; child is taking a diuretic. Ask a doctor or pharmacist before use if the child is under a doctor's care for any serious condition; taking any other drug.
 When using this product take with food or milk if upset stomach occurs

Stop use and ask a doctor if child experiences any of the following signs of stomach bleeding: feels faint, vomits blood, has bloody or black stools, or has stomach pain that does not get better; child has symptoms of heart problems or stroke: chest pain, trouble breathing, weakness in one part or side of body, slurred speech, leg swelling; the child does not get any relief within first day (24 hours) of treatment; fever or pain gets worse or lasts more than 3 days; redness or swelling is present in the painful area; any new symptoms appear.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Each 5 mL contains: sodium 3 mg. Store at 20-25°C (68-77°F).
Directions: This product does not contain directions or complete warnings for adult use. Do not give more than directed. Shake well before using.
 mL = mL (see Table right dose on chart). If possible, use weight to dose; otherwise use age.
 Repeat dose every 6-8 hours, if needed. Do not use more than 4 times a day. Measure only with the dosing cup provided. Dosing cup to be used with Children's Advil Suspension only. Do not use with other products. Dose lines account for product remaining in cup due to thickness of suspension.

Dosing Chart		
Weight (lb)	Age (yr)	Dose (mL)
under 28 lb	under 2 yr	ask a doctor
28-35 lb	2-3 yr	5 mL
36-47 lb	4-5 yr	7.5 mL
48-59 lb	6-8 yr	10 mL
60-71 lb	9-10 yr	12.5 mL
72-85 lb	11 yr	15 mL

Questions or comments? call 1-800-48-ADVIL.
 Dist. by GSK CH, Warren, NJ 07059 ©2021 GSK or its licensor
 For most recent product information, visit www.Advil.com

PAA166830

LOT:

EXP:

UNVARNISHED UNPRINTED AREA
 FOR LOT AND EXP. DATE.
 (PRE-PRINTED LOT INFORMATION AS EXAMPLE,
 SHOULD MATCH MARKET REQUIREMENTS)

PAA166830

DRUG FACTS TEXT DEFINED	TYPE SIZE
• HEADINGS	4.5 pt
• SUBHEADINGS/BODY TEXT	4.5 pt
• LEADING	4.75 pt
• BULLETS	N/A
• SPACE BEFORE BULLET	N/A
• WARNING BOX LINE, HAIRLINES	.5 pt



PAA166841

DRUG FACTS TEXT DEFINED	TYPE SIZE
• DRUG FACTS TITLE	9 pt
• DRUG FACTS CONTINUED	8 pt
• HEADINGS	8 pt
• SUBHEADINGS/BODY TEXT	6 pt
• LEADING	6.5 pt
• # OF CHARACTERS PER INCH	<39
• BULLETS	5 pt
• SPACE BEFORE BULLET	2 ems
• BARLINES, HAIRLINES	1.5 pt, .5 pt
• SPACE BETWEEN HAIRLINES AND BOX END	2 spaces

Do Not Use if plastic wrap imprinted with "SAFETY SEAL" is broken or missing.

**Children's
Advil**
Suspension

Ibuprofen Oral Suspension
100 mg per 5 mL
Pain Reliever/Fever Reducer (NSAID)

White Grape-Flavored Liquid
4 FL OZ (120 mL)

Active ingredient (in each 5 mL): Ibuprofen 100 mg

Use temporarily to reduce fever, relieve minor aches and pains due to the common cold, flu, sore throat, headaches and toothaches

Warnings
Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: hives, facial swelling, asthma (wheezing), shock, skin redness, rash, blisters. If an allergic reaction occurs, stop use and seek medical help right away.
Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if the child has had stomach ulcers or bleeding problems, takes a blood thinning (anticoagulant) or stomach drug, takes other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others), takes more or for a longer time than directed.
Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.
Sore throat warning: Severe or persistent sore throat or sore throat accompanied by high fever, headache, nausea, and vomiting may be serious. Consult doctor promptly. Do not use more than 2 days or administer to children under 3 years of age unless directed by doctor.
 Do not use if the child has ever had an allergic reaction to any other pain reliever/fever reducer; right before or after heart surgery.
 Ask a doctor before use if stomach bleeding warning applies to the child, child has a history of stomach problems, such as heartburn; child has high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke; child has not been drinking fluids; child has lost a lot of fluid due to vomiting or diarrhea; child is taking a diuretic.
 Ask a doctor or pharmacist before use if the child is under a doctor's care for any serious condition, taking any other drug.

When using this product take with food or in AM if stomach upset occurs
 Stop use and ask a doctor if child experiences any of the following signs of stomach bleeding: feels faint, vomit to blood, has bloody or black stools, or has stomach pain that does not get better; child has symptoms of heart problems or stroke: chest pain, trouble breathing, weakness in one part or side of body, slurred speech, leg swelling; the child does not get any relief within first day (24 hours) of treatment; fever or pain gets worse or lasts more than 3 days; redness or swelling is present in the painful area; any new symptoms appear.
 Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.
 Each 5 mL contains sodium 3 mg. Store at 20-25° C (68-77° F).
Directions: This product does not contain directions or complete warnings for adult use. Do not give more than directed. Shake well before using. mL = mL fl. oz. Find right dose on chart. If possible, use weight to dose otherwise use age. Repeat dose every 6-8 hours, if needed. Do not use more than 4 times a day. Measure only with the dosing cup provided. Dosing cup to be used with Children's Advil Suspension only. Do not use with other products. Dose less account for product remaining in cup due to thickness of suspension.

Dosing Chart		
Weight (lb)	Age (yr)	Dose (mL)
Under 24 lb	Under 2 yr	Ask a doctor
24-35 lb	2-3 yr	5 mL
36-47 lb	4-5 yr	7.5 mL
48-59 lb	6-8 yr	10 mL
60-71 lb	9-10 yr	12.5 mL
72-85 lb	11 yr	15 mL

Questions or comments? call 1-800-468-ADVIL.
 Dist. by: GSK CH, Warren, NJ 07059 ©2021 GSK or licensee
 For most recent product information, visit www.Achtel.com

LOT: PAA166834

EXP: [REDACTED]

UNVARNISHED UNPRINTED AREA
 FOR LOT AND EXP. DATE.
 (PRE-PRINTED LOT INFORMATION AS EXAMPLE,
 SHOULD MATCH MARKET REQUIREMENTS)

PAA166834

DRUG FACTS TEXT DEFINED	TYPE SIZE
• HEADINGS	4.5 pt
• SUBHEADINGS/BODY TEXT	4.5 pt
• LEADING	4.57 pt
• BULLETS	N/A
• SPACE BEFORE BULLET	N/A
• WARNING BOX LINE, HAIRLINES	.5 pt

Children's Advil Suspension

Do Not Use if plastic wrap imprinted with "SAFETY SEAL" is broken or missing.

Drug Facts (continued)

- child has lost a lot of fluid due to vomiting or diarrhea
- child is taking a diuretic

Ask a doctor or pharmacist before use if the child is

- under a doctor's care for any serious condition
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- child experiences any of the following signs of stomach bleeding:
 - feels faint
 - vomits blood
 - has bloody or black stools
 - has stomach pain that does not get better
- child has symptoms of heart problems or stroke:
 - chest pain
 - trouble breathing
 - weakness in one part or side of body
 - slurred speech
 - leg swelling
- the child does not get any relief within first day (24 hours) of treatment
- fever or pain gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- this product does not contain directions or complete warnings for adult use
- do not give more than directed
- shake well before using
- mL = milliliter
- find right dose on chart. If possible, use weight to dose; otherwise use age.
- repeat dose every 8-8 hours, if needed
- do not use more than 4 times a day
- measure only with the dosing cup provided. Dosing cup to be used with Children's Advil Suspension only. Do not use with other products. Dose lines account for product remaining in cup due to thickness of suspension.

Weight (lb)	Age (yr)	Dose (mL)
Under 24 lb	Under 2 yr	Ask a doctor
24-35 lb	2-3 yr	5 mL
36-47 lb	4-5 yr	7.5 mL
48-59 lb	6-8 yr	10 mL
60-71 lb	9-10 yr	12.5 mL
72-95 lb	11 yr	15 mL

Children's Advil Suspension

Drug Facts (continued)

Other information

- each 5 mL contains: sodium 3 mg
- one dose lasts 6-8 hours
- store at 20-25°C (68-77°F)
- see bottom of box for lot number and expiration date

Inactive ingredients

carboxymethylcellulose sodium, citric acid monohydrate, edetate disodium, FD&C red no. 40, glycerin, microcrystalline cellulose, natural and artificial flavor, polysorbate 80, propylene glycol, purified water, sodium benzoate, sorbitol solution, sucrose, xanthan gum

Questions or comments?
call toll-free 1-800-88-ADVIL or ask your pharmacist, doctor or health care professional

Long Lasting: Up to 8 hours of fever reduction.
Great Convenience: Long duration means fewer doses needed per day.
Alcohol-Free

For most recent product information, visit www.Advil.com
Distributed by: GSK Consumer Healthcare
Warren, NJ 07059
Trademarks are owned by or licensed to the GSK group of companies. ©2021 GSK

Children's Advil Suspension

Ibuprofen Oral Suspension
100 mg per 5 mL
Pain Reliever/Fever Reducer (NSAID)

Fever
Aches & Pains
Lasts Up To
8
Hours

Bubble Gum-Flavored Liquid
for ages 2-11 years

4 FL OZ (120 mL) Alcohol-Free

Important: Read all product information before using. Keep this box for important information. This product is intended for use in children ages 2 years - 11 years.

Drug Facts

Active ingredient (in each 5 mL) Purpose
Ibuprofen 100 mg (NSAID)*Fever reducer/Pain reliever
*nonsteroidal anti-inflammatory drug

Uses temporarily:

- reduces fever
- relieves minor aches and pains due to the common cold, flu, sore throat, headaches and toothaches

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if the child:

- has had stomach ulcers or bleeding problems
- takes a blood thinning (anticoagulant) or steroid drug
- takes other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- takes more or for a longer time than directed

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Sore throat warning: Severe or persistent sore throat or sore throat accompanied by high fever, headache, nausea, and vomiting may be serious. Consult doctor promptly. Do not use more than 2 days or administer to children under 3 years of age unless directed by doctor.

Do not use

- if the child has ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- stomach bleeding warning applies to the child
- child has problems or serious side effects from taking pain relievers or fever reducers
- child has a history of stomach problems, such as heartburn
- child has high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- child has not been drinking fluids

PAA166839

PAA166839

DRUG FACTS TEXT DEFINED	TYPE SIZE
• DRUG FACTS TITLE	9 pt
• DRUG FACTS CONTINUED	8 pt
• HEADINGS	8 pt
• SUBHEADINGS/BODY TEXT	6 pt
• LEADING	6.5 pt
• # OF CHARACTERS PER INCH	<39
• BULLETS	5 pt
• SPACE BEFORE BULLET	2 ems
• BARLINES, HAIRLINES	1.5 pt, .5 pt
• SPACE BETWEEN HAIRLINES AND BOX END	2 spaces

Do Not Use if plastic wrap imprinted with "SAFETY SEAL" is broken or missing.

Children's Advil Suspension
 Ibuprofen Oral Suspension
 100 mg per 5 mL
 Pain Reliever/Fever Reducer (NSAID)

Bubble Gum-Flavored Liquid
 4 FL OZ (120 mL)

Act as directed (in each 5 mL) Ibuprofen 100 mg
 Use temporarily reduces fever, or eases minor aches and pains due to the common cold, flu, sore throat, headache and toothaches

Warnings
Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include hives, facial swelling, asthma (wheezing), shock, skin reddening, rash, blisters. If an allergic reaction occurs, stop use and seek medical help right away.
Stomach bleeding warning: The product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if the child has had stomach ulcers or bleeding problems, takes a blood thinning (anticoagulant) or steroid drug, takes other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others), takes more or for a longer time than directed.
Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.
Severe throat warning: Severe or persistent sore throat or sore throat accompanied by high fever, headache, nausea, and vomiting may be serious. Consult doctor promptly. Do not use more than 2 days or administer to child under 3 years of age unless directed by doctor.
Do not use if the child has ever had an allergic reaction to any other pain reliever/fever reducer, right before or after heart surgery.
Ask a doctor before use: If stomach bleeding warning applies to the child, child has a history of stomach problems, such as heartburn; child has high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke; child has not been drinking fluids; child has lost a lot of fluid due to vomiting or diarrhea; child is taking a diuretic.
Ask a doctor or pharmacist before use if the child is under a doctor's care for any serious condition; taking any other drug.

When using this product take with food or milk if stomach upset occurs.
Stop use and ask a doctor if child experiences any of the following signs of stomach bleeding: feels faint, vomits blood, has bloody or black stools, or has stomach pain that does not get better; child has symptoms of heart problems or stroke: chest pain, trouble breathing, weakness in one part or side of body, slurred speech, leg swelling, the child does not get any relief within first day (24 hours) of treatment, fever or pain gets worse or lasts more than 3 days; redness or swelling is present in the mouth area, any new symptoms appear.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.
 Each 5 mL contains sodium 3 mg. Store at 20-25°C (68-77°F).
Directions: This product does not contain directions or complete warnings for adult use. Do not give more than directed. Shake well before using. mL = milliliter. Find right dose on chart. If possible, use weight to dose; otherwise use age. Repeat dose every 6-8 hours, if needed. Do not use more than 4 times a day. Measure only with the dosing cup provided. Dosing cup to be used for Children's Advil Suspension only. Do not use with other products. Dose lines account for product remaining in cup due to thickness of suspension.

Dosing Chart		
Weight (lb)	Age (yr)	Dose (mL)
Under 24 lb	Under 2 yr	Ask a doctor
24-35 lb	2-3 yr	5 mL
36-47 lb	4-5 yr	7.5 mL
48-59 lb	6-8 yr	10 mL
60-71 lb	9-10 yr	12.5 mL
72-85 lb	11 yr	15 mL


LOT: PAA166832
EXP: [REDACTED]

For questions or comments, call 1-800-39-ADAIL. Mfg. by GSK (U.S.) Warner, NJ 07090 02021 GSK or its owner. For most recent product information, visit www.Advil.com

UNVARNISHED UNPRINTED AREA
 FOR LOT AND EXP. DATE.
 (PRE-PRINTED LOT INFORMATION AS EXAMPLE,
 SHOULD MATCH MARKET REQUIREMENTS)

PAA166832

DRUG FACTS TEXT DEFINED	TYPE SIZE
• HEADINGS	4.5 pt
• SUBHEADINGS/BODY TEXT	4.5 pt
• LEADING	4.6 pt
• BULLETS	N/A
• SPACE BEFORE BULLET	N/A
• WARNING BOX LINE, HAIRLINES	.5 pt



DATA MATRIX CODE MUST HAVE 12 CELLS MINIMUM (SIZE AT LEAST 0.25 MM EACH) WITH A WHITE QUIET ZONE OF MINIMUM 1 MM

CODING AREA TO BE PRINTED IN A DARK COLOR AND UNVARNISHED

PAA166840

DRUG FACTS TEXT DEFINED	TYPE SIZE
• DRUG FACTS TITLE	9 pt
• DRUG FACTS CONTINUED	8 pt
• HEADINGS	8 pt
• SUBHEADINGS/BODY TEXT	6 pt
• LEADING	6.5 pt
• # OF CHARACTERS PER INCH	<39
• BULLETS	5 pt
• SPACE BEFORE BULLET	2 ems
• BARLINES, HAIRLINES	1.5 pt, .5 pt
• SPACE BETWEEN HAIRLINES AND BOX END	2 spaces

Do Not Use if plastic wrap imprinted with "SAFETY SEAL" is broken or missing.

**Children's
Advil
Suspension**

Ibuprofen Oral Suspension
100 mg per 5 mL

Pain Reliever/Fever Reducer (NSAID)

Sugar-Free Dye-Free
Berry-Flavored Liquid
4 FL OZ (120 mL)

Active ingredient (in each 5 mL): Ibuprofen 100 mg

Effect: temporarily reduces fever, relieves minor aches and pains due to the common cold, flu, sore throat, headaches and toothaches

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include hives, facial swelling, asthma (wheezing), shock, skin redness, rash, blisters. If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if the child has had stomach ulcers or bleeding problems, takes a blood thinning (anticoagulant) or steroid drug, takes other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others), takes more or for a longer time than directed. Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Sore throat warning: Severe or persistent sore throat or sore throat accompanied by high fever, headache, nausea, and vomiting may be serious. Consult doctor promptly. Do not use more than 2 days or administer to children under 3 years of age unless directed by doctor.

Do not use if the child has ever had an allergic reaction to any other pain reliever/fever reducer, right before or after heart surgery

Ask a doctor before use if stomach bleeding warning applies to the child, child has a history of stomach problems, asthma/heartburn; child has high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke; child has not been drinking fluids; child has lost a lot of fluid due to vomiting or diarrhea; child is taking a diuretic.

Ask a doctor or pharmacist before use if the child is under a doctor's care for any serious condition; taking any other drug.

When using this product take with food or milk if stomach upset occurs. Stop use and ask a doctor if child experiences any of the following signs of stomach bleeding: feels faint, vomits blood, has bloody or black stools, or has stomach pain that does not get better; child has symptoms of heart problems or stroke: chest pain, trouble breathing, weakness in one part or side of body, slurred speech, leg swelling, the child does not get any relief within first day (24 hours) of treatment; fever or pain gets worse or lasts more than 3 days; redness or swelling is present in the painful area; any new symptoms appear. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Each 5 mL contains sodium 10 mg. Store at 20-25°C (68-77°F).

Directions: This product does not contain directions or complete warnings for adult use. Do not give more than directed. Shake well before using. mL = mL for. Find right dose on chart. If possible, use weight to dose; otherwise use age. Repeat dose every 6-8 hours, if needed. Do not use more than 4 times a day. Measure only with the dosing cup provided. Dosing cup to be used with Children's Advil Suspension only. Do not use with other products. Dose line account for product remaining in cup due to thickness of suspension.

Dosing Chart		
Weight (lb)	Age (yr)	Dose (mL)
Under 24 lb	Under 2 yr	Ask a doctor
24-35 lb	2-3 yr	5 mL
36-47 lb	4-5 yr	7.5 mL
48-59 lb	6-8 yr	10 mL
60-71 lb	9-10 yr	12.5 mL
72-95 lb	11 yr	15 mL

Questions or comments? call 1-800-88-ADVIL
 Dist. by: SUN-ON, Warren, NJ 07059 ©2021 SUN, or licensee
 For most recent product information, visit www.Advil.com.

LOT: PAA166833

EXP: XXXXXXXXXX

UNVARNISHED UNPRINTED AREA FOR LOT AND EXP. DATE. (PRE-PRINTED LOT INFORMATION AS EXAMPLE, SHOULD MATCH MARKET REQUIREMENTS)

PAA166833

DRUG FACTS TEXT DEFINED	TYPE SIZE
• HEADINGS	4.5 pt
• SUBHEADINGS/BODY TEXT	4.5 pt
• LEADING	4.57 pt
• BULLETS	N/A
• SPACE BEFORE BULLET	N/A
• WARNING BOX LINE, HAIRLINES	.5 pt



PAA166838

DRUG FACTS TEXT DEFINED	TYPE SIZE
• DRUG FACTS TITLE	9 pt
• DRUG FACTS CONTINUED	8 pt
• HEADINGS	8 pt
• SUBHEADINGS/BODY TEXT	6 pt
• LEADING	6.5 pt
• # OF CHARACTERS PER INCH	<39
• BULLETS	5 pt
• SPACE BEFORE BULLET	2 ems
• BARLINES, HAIRLINES	1.5 pt, .5 pt
• SPACE BETWEEN HAIRLINES AND BOX END	2 spaces

Do Not Use if plastic wrap imprinted with "SAFETY SEAL" is broken or missing.

**Children's
 Advil**
 Suspension

Ibuprofen Oral Suspension
 100 mg per 5 mL

Pain Reliever/Fever Reducer (NSAID)

Dye-Free / Alcohol-Free
 Blue Raspberry-Flavored Liquid
 4 FL OZ (120 mL)

Active ingredient (in each 5 mL): Ibuprofen 100 mg
 (See temporarily reduces fever, relieves minor aches and pains due to the common cold, flu, sore throat, headaches and toothaches)

Warnings
Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: hives, facial swelling, asthma (wheezing), shock, skin redness, rash, hives. If an allergic reaction occurs, stop use and seek medical help right away.
Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if the child has had stomach ulcers or bleeding problems, takes a blood thinning (anticoagulant) or steroid drug, takes other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others), takes more or for a longer time than directed.
Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.
Sore throat warning: Severe or persistent sore throat or sore throat accompanied by high fever, headache, nausea, and vomiting may be serious. Consult a doctor promptly. Do not use more than 2 days or administer to children under 3 years of age unless directed by a doctor.
Do not use if the child has ever had an allergic reaction to any other fever reducer/pain reliever, right before or after heart surgery
Ask a doctor before use if stomach bleeding warning applies to the child; child has a history of stomach problems, such as heartburn; child has high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke; child has not been drinking fluids; child has lost a lot of fluid due to vomiting or diarrhea; child is taking a diuretic
Ask a doctor or pharmacist before use if the child is under a doctor's care for any serious condition; taking any other drug

When using this product take with food or milk if upset stomach occurs
 Stop use and call a doctor if child experiences any of the following signs of stomach bleeding: pale faint, vomits blood, has bloody or black stools, or has stomach pain that does not get better; child has symptoms of heart problems or stroke: chest pain, trouble breathing, weakness in one part or side of body, slurred speech, big swelling; the child does not get any relief within first day (24 hours) of treatment; fever or pain gets worse or lasts more than 3 days; redness or swelling is present in the painful area; any new symptoms appear
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Each 5 mL contains sodium 10 mg. Store at 20-25°C (68-77°F).
Directions: This product does not contain directions or complete warnings for adult use. Do not give more than directed. Shake well before using. mL = milliliter. Find right dose on chart. If possible, use weight to dose; otherwise use age. Repeat dose every 6-8 hours, if needed. Do not use more than 4 times a day. Measure only with the dosing cup provided. Dosing cup to be used with Children's Advil Suspension only. Do not use with other products. Dose lines account for product remaining in cup due to thickness of suspension.

Dosing Chart		
Weight (lb)	Age (yr)	Dose (mL)
under 24 lb	under 2 yr	ask a doctor
24-35 lb	2-3 yr	5 mL
36-47 lb	4-5 yr	7.5 mL
48-59 lb	6-8 yr	10 mL
60-71 lb	9-10 yr	12.5 mL
72-85 lb	11 yr	15 mL

Quantity or comments? call 1-800-89-ADVIL
 Dist. by: GSK CK, Warren, NJ 07059 ©2021 GSK or licensee
 For most recent product information, visit www.Advil.com

PAA166831
 LOT:
 EXP:

UNVARNISHED UNPRINTED AREA
 FOR LOT AND EXP. DATE.
 (PRE-PRINTED LOT INFORMATION AS EXAMPLE,
 SHOULD MATCH MARKET REQUIREMENTS)

PAA166831

DRUG FACTS TEXT DEFINED	TYPE SIZE
• HEADINGS	4.5 pt
• SUBHEADINGS/BODY TEXT	4.5 pt
• LEADING	4.5 pt
• BULLETS	N/A
• SPACE BEFORE BULLET	N/A
• WARNING BOX LINE, HAIRLINES	.5 pt

This is a representation of an electronic record that was signed electronically. Following this are manifestations of any and all electronic signatures for this electronic record.

/s/

NUSHIN F TODD
09/20/2022 09:12:00 AM