

Drug Facts (continued)

Warnings
Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin.

Symptoms may include: hives facial swelling skin redness rash blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you or child:

- has had stomach ulcers or bleeding problems
- takes a blood thinning (anticoagulant) or steroid drug
- takes other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- takes more or for a longer time than directed

Severe throat warning: Severe or persistent sore throat or sore throat accompanied by high fever, headache, nausea, and vomiting may be serious. Consult doctor promptly. Do not use more than 2 days or administer to children under 3 years of age unless directed by doctor.

Do not use

- if the child has ever had an allergic reaction to ibuprofen or any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- stomach bleeding warning applies to your child
- child has a history of stomach problems, such as heartburn
- child has problems or serious side effects from taking pain relievers or fever reducers
- child has not been drinking fluids child has lost a lot of fluid due to vomiting or diarrhea
- child has high blood pressure, heart disease, liver cirrhosis, or kidney disease
- child has asthma child is taking a diuretic

Ask a doctor or pharmacist before use if the child is

- under a doctor's care for any serious condition taking any other drug

When using this product

- take with food or milk if stomach upset occurs
- the risk of heart attack or stroke may increase if you use more than directed or for longer than directed

Stop use and ask a doctor if

- child experiences any of the following signs of stomach bleeding:
 - feels faint vomits blood has bloody or black stools
 - has stomach pain that does not get better
- the child does not get any relief within first day (24 hours) of treatment
- fever or pain gets worse or lasts more than 3 days
- redness or swelling is present in the painful area any new symptoms appear

Drug Facts (continued)

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions

- this product does not contain directions or complete warnings for adult use
- do not give more than directed
- shake well before using
- first right dose on chart below. If possible, use weight to dose; otherwise use age.
- if needed, repeat dose every 6-8 hours
- do not use more than 4 times a day

Dosing Chart

Weight (lb)	Age (mos)	Dose (mL)
under 6 mos		ask a doctor
12-17 lbs	6-11 mos	1.25 mL
18-23 lbs	12-23 mos	1.875 mL

Other information

- store between 20-25°C (68-77°F)
- do not use if cap seal is broken
- "SAFETY SEAL" or "USE WITH ENCLOSED DOSING DEVICE ONLY" is broken or missing

Inactive ingredients: anhydrous citric acid, carmellose, FD&C red no. 40, flavors, glycerin, polyacrylate 10, pregelatinized starch, purified water, sodium benzoate, sorbitol solution, sucrose, xanthan gum

Questions or comments?
 Call 1-877-965-3665 (ask for) or 215-223-8255 (collect)

New Information. Read Entire Label
 NDC 50580-100-18
For Ages 6 Mos. to 23 Mos.

**Concentrated Motrin®
 Infants' Drops**
 contains Ibuprofen
Oral Suspension (NSAID)
 50 mg per 1.25 mL
Pain Reliever/Fever Reducer

**Lasts up to
 8 hours**

*Original
 Berry Flavor*










1/2 fl oz (15 mL)

Use only with enclosed syringe.



Actual Size

Distributed by: **JOHNSON & JOHNSON CONSUMER INC.**
 McNeil Consumer Healthcare Division
 Fort Washington, PA 19034 USA ©J&JCI 2016
www.motrin.com



(b) (4)

Drug Facts

Active ingredient (in each 1.25 mL)
Ibuprofen (NSAID)
Pain reliever/fever reducer

Purpose
Pain reliever/fever reducer

Uses temporarily: ■ fever ■ pain ■ headache ■ sore throat ■ teething ■ earache ■ toothache

Warnings
■ Do not use if child has ever had an allergic reaction to ibuprofen or any other pain reliever/fever reducer
■ Do not use if child has ever had heart surgery

Directions
■ Shake well before using
■ Follow directions on chart. If possible, use weight to dose, otherwise use age.

Drug Facts (continued)

Warnings
Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin.
Symptoms may include: ■ hives ■ facial swelling ■ asthma (wheezing) ■ shock ■ skin redness ■ rash ■ blisters
If an allergic reaction occurs, stop use and seek medical help right away.
Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if your child:
■ has had stomach ulcers or bleeding problems
■ takes other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
■ takes more or for a longer time than directed
Sore throat warning: Severe or persistent sore throat or sore throat accompanied by high fever, headache, nausea, and vomiting may be serious. Consult doctor promptly. Do not use more than 2 days or administer to children under 3 years of age unless directed by doctor.

Do not use
■ If the child has ever had an allergic reaction to ibuprofen or any other pain reliever/fever reducer
■ Right before or after heart surgery

Ask a doctor before use if
■ Stomach bleeding warning applies to your child
■ Child has a history of stomach problems, such as heartburn
■ Child has problems or serious side effects from taking pain relievers or fever reducers
■ Child has not been drinking fluids ■ Child has lost a lot of fluid due to vomiting or diarrhea
■ Child has high blood pressure, heart disease, liver cirrhosis, or kidney disease
■ Child has asthma ■ Child is taking a diuretic

Ask a doctor or pharmacist before use if the child is
■ Under a doctor's care for any serious condition ■ Taking any other drug

When using this product
■ Take with food or milk if stomach upset occurs
■ The risk of heart attack or stroke may increase if you use more than directed or for longer than directed

Stop use and ask a doctor if
■ Child experiences any of the following signs of stomach bleeding:
■ Feels faint ■ Vomits blood ■ Has bloody or black stools
■ Has stomach pain that does not get better
■ The child does not get any relief within first day (24 hours) of treatment
■ Fever or pain gets worse or lasts more than 3 days
■ Redness or swelling is present in the painful area ■ Any new symptoms appear

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions
■ This product does not contain directions or complete warnings for adult use
■ Do not give more than directed ■ Shake well before using
■ Find right dose on chart. If possible, use weight to dose, otherwise use age.

Drug Facts (continued)

■ mL = milliliter
■ Measure with the dosing device provided. Do not use with any other device.
■ Disperse liquid slowly into the child's mouth, toward the inner cheek.
■ If needed, repeat dose every 6-8 hours
■ Do not use more than 4 times a day

Dosing Chart

Weight (lb)	Age (mos)	Dose (mL)
under 6 mos		ask a doctor
12-17 lbs	6-11 mos	1.25 mL
8-23 lbs	12-23 mos	1.875 mL

Other information
■ Store between 20-25°C (68-77°F)
■ Do not use if carton is opened or carton flap is imprinted "SAFETY SEAL" or bottle was imprinted "SAFETY SEAL" and "USE WITH ENCLOSED DOSING DEVICE ONLY" is broken or missing

Inactive ingredients
anhydrous citric acid, caramel, flavors, glycerin, polyorbital 80, pregelatinized starch, purified water, sodium benzoate, sorbitol solution, sucrose, xanthan gum

Questions or comments?
Call 1-877-895-3665 (toHear) or 215-275-4755 (collect)

Concentrated Motrin® Infants' Drops DYE-FREE



Concentrated Motrin® Infants' Drops
contains Ibuprofen Oral Suspension (NSAID)
50 mg per 1.25 mL
Pain Reliever/Fever Reducer
Non-Staining Original Berry Flavor
1 fl oz (30 mL)

Actual Size

New Information. Read Entire Label
NDC 50580-198-50

For Ages 6 Mos. to 23 Mos.

Concentrated Motrin® Infants' Drops DYE-FREE

contains **Ibuprofen Oral Suspension (NSAID)**
50 mg per 1.25 mL
Pain Reliever/Fever Reducer

Lasts up to 8 hours

Use only with enclosed syringe.

1 fl oz (30 mL)

LOT: _____

EXP: _____

Distributed by: **JOHNSON & JOHNSON CONSUMER INC.**
McNeil Consumer Healthcare Division
Fort Washington, PA 19034 USA ©J&JCI 2016
www.motrin.com

USA
000000
887

Drug Facts (continued)

■ mL = milliliter
■ Measure with the dosing device provided. Do not use with any other device.
■ Disperse liquid slowly into the child's mouth, toward the inner cheek.
■ If needed, repeat dose every 6-8 hours
■ Do not use more than 4 times a day








Dosing Chart

Weight (lb)	Age (mos)	Dose (mL)
under 6 mos		ask a doctor
12-17 lbs	6-11 mos	1.25 mL
8-23 lbs	12-23 mos	1.875 mL

Other information
■ Store between 20-25°C (68-77°F)
■ Do not use if carton is opened or carton flap is imprinted "SAFETY SEAL" or bottle was imprinted "SAFETY SEAL" and "USE WITH ENCLOSED DOSING DEVICE ONLY" is broken or missing

Inactive ingredients
anhydrous citric acid, caramel, flavors, glycerin, polyorbital 80, pregelatinized starch, purified water, sodium benzoate, sorbitol solution, sucrose, xanthan gum

Questions or comments?
Call 1-877-895-3665 (toHear) or 215-275-4755 (collect)



(b) (4)