

**3-DAY
TREATMENT**
COMBINATION PACK

Suppository



Vaginal Suppositories
with 3 Applicators

plus External Cream
for Itch Relief

MONISTAT[®] 3

COMBINATION PACK
Miconazole Nitrate Vaginal Suppositories (200 mg) and
Miconazole Nitrate Vaginal Cream (2%) — Vaginal Antifungal

TAMPER-EVIDENT UNIT
DO NOT USE IF
PRINTED SUPPOSITORY
BLISTERS OR SEAL
OVER TUBE ARE TORN,
OPENED OR
INCOMPLETELY
SEALED

MONISTAT[®] 3

COMBINATION PACK
Miconazole Nitrate Vaginal Suppositories (200 mg) and
Miconazole Nitrate Vaginal Cream (2%) — Vaginal Antifungal

Exp
Lot



- ✓ CURES most vaginal YEAST INFECTIONS
- ✓ Relieves associated external itching and irritation
- ✓ Suppositories contain just 2 Ingredients

3 Suppositories Net Wt. 2.5 g each + 0.32 oz (9 g) Tube



Dist. by Insight Pharmaceuticals LLC
Tarrytown, NY 10591
A Prestige Consumer Healthcare company
©2021 Trade dress is owned by
Insight Pharmaceuticals LLC
All rights reserved. MNUS028105

If you have any
questions or
comments, please call
1-877-MONISTAT
(1-877-666-4782)

MONISTAT[®] 3
COMBINATION PACK
Miconazole Nitrate Vaginal Suppositories (200 mg) and
Miconazole Nitrate Vaginal Cream (2%) — Vaginal Antifungal

Drug Facts

Active ingredients
Miconazole nitrate (200 mg in each suppository) Vaginal antifungal
Miconazole nitrate 2% (external cream) Vaginal antifungal

Purpose
If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions
Before using this product read the enclosed consumer information leaflet for complete directions and information.
adults and children 12 years of age and over:
-suppositories: insert 1 suppository into the vagina at bedtime for three nights in a row. Throw applicator away after use.
-external cream: squeeze a small amount of cream onto your fingertip. Apply the cream onto the itchy, irritated skin outside the vagina. Use 2 times daily for up to 7 days, as needed.
children under 12 years of age: ask a doctor.

Other information
do not use if printed suppository blisters are torn, open or incompletely sealed
do not use if seal over tube opening has been punctured
do not purchase if carton is open
do not use if seal over tube opening has been punctured
store at 20°-25°C (68°-77°F)

Inactive ingredients
suppository: hydrogenated vegetable oil base
external cream: benzoic acid, cetyl alcohol, isopropyl myristate, polyorbate 60, potassium hydroxide, propylene glycol, purified water, stearyl alcohol

Warnings
For vaginal use only
Do not use if you have never had a vaginal yeast infection diagnosed by a doctor.
Ask a doctor before use if you have
vaginal itching and discomfort for the first time
vaginal abdominal, back or shoulder pain, fever, chills, nausea, vomiting, or foul-smelling vaginal discharge. You may have a more serious condition.
vaginal yeast infections often (such as once a month or 3 in 6 months). You could be pregnant or have a serious underlying medical cause for your symptoms, including diabetes, or a weakened immune system.
been exposed to the human immunodeficiency virus (HIV) that causes AIDS
Ask a doctor or pharmacist before use if you are taking the prescription blood thinning medicine warfarin, because bleeding or bruising may occur.

Uses
relieves external itching and irritation due to a vaginal yeast infection
Treats vaginal yeast infections

When using this product
do not use tampons, douches, spermicides or other vaginal products. Condoms and diaphragms may be damaged and fail to prevent pregnancy or sexually transmitted diseases (STDs).

Questions?
If you have any questions or comments, please call 1-877-666-4782.

Drug Facts (continued)
do not have vaginal intercourse
mild increase in vaginal burning, itching or irritation may occur
if you do not get complete relief ask a doctor before using another product
Stop use and ask a doctor if:
symptoms do not get better in 3 days
symptoms last more than 7 days
you get a rash or hives, abdominal pain, fever, chills, nausea, vomiting, or foul-smelling vaginal discharge
If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

This is a representation of an electronic record that was signed electronically. Following this are manifestations of any and all electronic signatures for this electronic record.

/s/

PAMELA J HORN
11/17/2023 09:22:08 AM