

Lot/EXP Area



© 2014, Watson Consumer Health, Inc.
 All rights reserved. www.watson.com

02008783



02008783



TAMPER-EVIDENT BOTTLE
 DO NOT USE IF INNER FOIL SEAL IMPRINTED WITH
 "SEALED FOR YOUR PROTECTION" IS BROKEN OR MISSING

Drug Facts

Active ingredients (in each caplet)	Purposes
Acetaminophen 325 mg	Pain reliever
Aspirin 250 mg (NSAID)*	Pain reliever
Caffeine 65 mg	Pain reliever aid

*nonsteroidal anti-inflammatory drug

Use • treats migraine

Warnings
 Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: • skin reddening • blisters • rash

If a skin reaction occurs, stop use and seek medical help right away.

Allergy alert: Aspirin may cause a severe allergic reaction which may include: • hives • facial swelling • asthma (wheezing) • shock

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 2 caplets in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Caffeine warning: The recommended dose of this product contains about as much caffeine as a cup of coffee. Limit the use of caffeine-containing medications, foods, or beverages while taking this product because too much caffeine may cause nervousness, irritability, sleeplessness, and, occasionally, rapid heart beat.

Do not use

- if you have ever had an allergic reaction to acetaminophen, aspirin or any other pain reliever/fever reducer
- with any other drug containing acetaminophen (prescription or nonprescription).

If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist

Ask a doctor before use if

- you have never had migraines diagnosed by a health professional
- you have a headache that is different from your usual migraines
- you have the worst headache of your life
- you have fever and stiff neck

Drug Facts (continued)

- you have headaches beginning after or caused by head injury, exertion, coughing or bending
- you experienced your first headache after the age of 50
- you have daily headaches
- you have a migraine so severe as to require bed rest
- you have liver disease
- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have vomiting with your migraine headache

Ask a doctor or pharmacist before use if you are

- taking a prescription drug for: • diabetes • gout • arthritis
- under a doctor's care for any serious condition
- taking any other drug
- taking any other product that contains aspirin, acetaminophen, or any other pain reliever/fever reducer

Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
 - feel faint • vomit blood • have bloody or black stools
 - have stomach pain that does not get better
- your migraine is not relieved or worsens after first dose
- new or unexpected symptoms occur
- ringing in the ears or loss of hearing occurs

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.
 In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not use more than directed
- adults: take 2 caplets with a glass of water
- if symptoms persist or worsen, ask your doctor
- do not take more than 2 caplets in 24 hours, unless directed by a doctor
- under 18 years of age: ask a doctor

Other information

- store at controlled room temperature 20°-25°C (68°-77°F)
- close cap tightly after use
- read all product information before using. Keep this box for important information.

Inactive ingredients benzoic acid, carnauba wax, FD&C blue #1, hydroxypropylcellulose, hypromellose, light mineral oil, microcrystalline cellulose, polysorbate 20, povidone, propylene glycol, simethicone emulsion, sorbitan monooleate, stearic acid, titanium dioxide

Questions or comments? 1-800-468-7746





02008783



02008783

Glue





Lot/EXP Area



Distributed by: Hevert's Co. Inc. 12477 24C
 Reserp. No. 051-052 © 2014
 Visit us at www.heverts.com

12477 24C



12477 24C



TAMPER EVIDENT BOTTLE
 DO NOT USE IF INNER FOIL SEAL IMPRINTED WITH
 "SEALED BY YOUR PROTECTION" IS BROKEN OR MISSING

Drug Facts

Active ingredients (in each tablet) Purposes
 Acetaminophen 250 mg..... Pain reliever
 Aspirin 250 mg (NSAID)..... Pain reliever
 Caffeine 65 mg..... Pain reliever aid
 *nonsteroidal anti-inflammatory drug

Use • treats migraine

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.
Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:
 • skin reddening • blisters • rash
 If a skin reaction occurs, stop use and seek medical help right away.
Allergy alert: Aspirin may cause a severe allergic reaction which may include:
 • hives • facial swelling • asthma (wheezing) • shock
Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take • more than 2 tablets in 24 hours, which is the maximum daily amount • with other drugs containing acetaminophen • 3 or more alcoholic drinks every day while using this product.

Drug Facts (continued)

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you • are age 60 or older • have had stomach ulcers or bleeding problems • take a blood thinning (anticoagulant) or steroid drug • take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) • have 3 or more alcoholic drinks every day while using this product • take more or for a longer time than directed.
Caffeine warning: The recommended dose of this product contains about as much caffeine as a cup of coffee. Limit the use of caffeine-containing medications, foods, or beverages while taking this product because too much caffeine may cause nervousness, irritability, sleeplessness, and, occasionally, rapid heart beat.

Do not use • if you have ever had an allergic reaction to acetaminophen, aspirin or any other pain reliever/fever reducer • with any other drug containing acetaminophen (prescription or nonprescription), if you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist

Ask a doctor before use if • you have never had migraines diagnosed by a health professional • you have a headache that is different from your usual migraines • you have the worst headache of your life • you have fever and stiff neck • you have headaches beginning after or caused by head injury, exertion, coughing or bending • you experienced your first headache after the age of 50 • you have daily headaches • you have a migraine so severe as to require bed rest • you have liver disease • stomach bleeding warning applies to you • you have a history of stomach problems, such as heartburn • you have high blood pressure, heart disease,

Drug Facts (continued)

liver cirrhosis, or kidney disease • you are taking a diuretic • you have asthma • you have problems or serious side effects from taking pain relievers or fever reducers • you have vomiting with your migraine headache

Ask a doctor or pharmacist before use if you are • taking a prescription drug for: • diabetes • gout • arthritis • under a doctor's care for any serious condition • taking any other drug • taking any other product that contains aspirin, acetaminophen, or any other pain reliever/fever reducer

Stop use and ask a doctor if • an allergic reaction occurs. Seek medical help right away. • you experience any of the following signs of stomach bleeding: • feel faint • vomit blood • have bloody or black stools • have stomach pain that does not get better • your migraine is not relieved or worsens after first dose • new or unexpected symptoms occur • ringing in the ears or loss of hearing occurs

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions • do not use more than directed • adults: take 2 tablets with a glass of water • if symptoms persist

Drug Facts (continued)


or worse, ask your doctor • do not take more than 2 tablets in 24 hours, unless directed by a doctor • under 18 years of age, ask a doctor
Other information • store at controlled room temperature 20°-25° (68°-77°) • close cap tightly after use • read all product information before using. Keep this box for important information.
Inactive ingredients: benzocaine, croscarmellose, FD&C blue #1, hydroxypropylcellulose, hydroxypropyl methylcellulose, polyethylene glycol, polyethylene glycol, simethicone emulsion, sodium monohydrogen phosphate, stearic acid, titanium dioxide
Questions or comments? 1-800-463-7746

12477 24C



12477 24C





Lot/EXP Area



Distributed by: Novartis Consumer Health, Inc.
 Parsippany, NJ 07654-0222 ©2014, 12477 100C
 Visit us at www.novartis.com

TAMPER EVIDENT BOTTLE
 DO NOT USE IF INNER FOIL SEAL IMPRINTED WITH
 "SEALED BY YOUR PROTECTION" IS BROKEN OR MISSING

Drug Facts
Active ingredients (in each tablet)
 Acetaminophen 250 mg... Pain reliever
 Aspirin 250 mg (NSAID)*... Pain reliever
 Caffeine 65 mg... Pain reliever aid
 *nonsteroidal anti-inflammatory drug

Use • treats migraine

Warnings
Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.
Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include • skin reddening • blisters • rash. If a skin reaction occurs, stop use and seek medical help right away.
Allergy alert: Aspirin may cause a severe allergic reaction which may include: • hives • facial swelling • asthma (wheezing) • shock
Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take
 • more than 2 tablets in 24 hours, which is the maximum daily amount
 • with other drugs containing acetaminophen
 • 3 or more alcoholic drinks every day while using this product
Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you
 • are age 60 or older • have had stomach ulcers or bleeding problems
 • take a blood thinning (anticoagulant) or steroid drug
 • take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
 • have 3 or more alcoholic drinks every day while using this product
 • take more or for a longer time than directed
Caffeine warning: The recommended dose of this product contains about as much caffeine as a cup of coffee. Limit the use of caffeine-containing medications, foods, or beverages while taking this product because too much caffeine may cause nervousness, irritability, sleeplessness, and, occasionally rapid heart beat.
Do not use • if you have ever had an allergic reaction to acetaminophen, aspirin or any other pain reliever/fever reducer
 • with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist

EXCEDRIN
MIGRAINE
 Acetaminophen, Aspirin (NSAID) and Caffeine
 Pain Reliever/Pain Reliever Aid
 100 TABLETS

Drug Facts (continued)
Ask a doctor before use if
 • you have never had migraines diagnosed by a health professional
 • you have a headache that is different from your usual migraines
 • you have the worst headache of your life
 • you have fever and stiff neck
 • you have headaches beginning after or caused by head injury, exertion, coughing or bending
 • you experienced your first headache after the age of 50
 • you have daily headaches
 • you have a migraine so severe as to require bed rest
 • you have liver disease • stomach bleeding warning applies to you
 • you have a history of stomach problems, such as heartburn
 • you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
 • you are taking a diuretic • you have asthma
 • you have problems or serious side effects from taking pain relievers or fever reducers
 • you have vomiting with your migraine headache
Ask a doctor or pharmacist before use if you are
 • taking a prescription drug for: • diabetes • gout • arthritis
 • under a doctor's care for any serious condition
 • taking any other drug
 • taking any other product that contains aspirin, acetaminophen, or any other pain reliever/fever reducer
Stop use and ask a doctor if
 • an allergic reaction occurs. Seek medical help right away.
 • you experience any of the following signs of stomach bleeding:
 • feel faint • vomit blood • have bloody or black stools
 • have stomach pain that does not get better
 • your migraine is not relieved or worsens after first dose
 • new or unexpected symptoms occur
 • ringing in the ears or loss of hearing occurs
If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. Keep out of reach of children.
In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.
Directions • do not use more than directed
 • adults: take 2 tablets with a glass of water

EXCEDRIN
MIGRAINE
 Acetaminophen, Aspirin (NSAID) and Caffeine
 Pain Reliever/Pain Reliever Aid
 100 TABLETS

Drug Facts (continued)
 • If symptoms persist or worsen, ask your doctor
 • do not take more than 2 tablets in 24 hours, unless directed by a doctor
 • under 18 years of age, ask a doctor

Other information
 • store at controlled room temperature 20°-25°C (68°-77°F); excursions permitted to 15°-30°C (59°-86°F) when use is for a short period of time
 • see expiration date on product label
 • keep this and all other medicines away from children
 • keep this and all other medicines away from heat and moisture
 • keep this and all other medicines away from light
 • keep this and all other medicines away from fire
 • keep this and all other medicines away from water
 • keep this and all other medicines away from alcohol
 • keep this and all other medicines away from strong odors
 • keep this and all other medicines away from strong colors
 • keep this and all other medicines away from strong tastes
 • keep this and all other medicines away from strong smells
 • keep this and all other medicines away from strong colors
 • keep this and all other medicines away from strong tastes
 • keep this and all other medicines away from strong smells

Questions or comments? 1-800-465-7146

CODE 128 FPO
 12477 100C

CODE 128 FPO
 12477 100C

Glue

NDC 0067-2035-91

EXCEDRIN®

MIGRAINE


Acetaminophen, Aspirin (NSAID) and Caffeine

Pain Reliever/Pain Reliever Aid

100 GELTABS®
*GELATIN COATED TABLETS


100 GELTABS IN TWO 50 COUNT BOTTLES

135063C100




135063C100

135063C100




135063C100

135063C100




135063C100

135063C100



135063C100

Lot/EXP Area



3 0067 2035 91 2

Do not use if the seal is broken or missing.
Do not use if the seal is broken or missing.
Do not use if the seal is broken or missing.

Distributed by: Rowland Consumer Products, Inc.
Parapet, NJ 07652-0208 ©2014 135063C100
Visit us at www.excedrin.com

Drug Facts (continued)

Ask a doctor before use if you:

- have never had migraines diagnosed by a health professional
- you have a headache that is different from your usual migraines
- you have the worst headache of your life
- you have fever and stiff neck
- you have headaches beginning after or caused by head injury, exertion, coughing or bending
- you experienced your first headache after the age of 50
- you have daily headaches
- you have a migraine so severe as to require bed rest
- you have liver disease
- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have vomiting with your migraine headache

Ask a doctor or pharmacist before use if you are taking a prescription drug for:

- diabetes
- gout
- arthritis
- under a doctor's care for any serious condition
- taking any other drug
- taking any other product that contains aspirin, acetaminophen, or any other pain reliever/fever reducer

Stop use and ask a doctor if:

- you experience any of the following signs of stomach bleeding:
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
- your migraine is not relieved or worsens after first dose
- new or worsened symptoms occur
- ringing in the ears or loss of hearing occurs

Drug Facts

Active ingredients (in each geltab)

Acetaminophen 250 mg Pain reliever
Aspirin 250 mg (NSAID) Pain reliever
Caffeine 65 mg Pain reliever aid
*nonsteroidal anti-inflammatory drug

Use • treats migraine

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin rash
- hives
- facial swelling
- asthma (wheezing)
- shock

Allergy alert: Aspirin may cause a severe allergic reaction which may include:

- hives
- facial swelling
- asthma (wheezing)
- shock

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take more than 2 geltabs in 24 hours, which is the maximum daily amount.

- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chances are higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- take 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Caffeine warning: The recommended dose of this product contains about as much caffeine as a cup of coffee. Limit the use of caffeine-containing medications, foods, or beverages while taking this product because too much caffeine may cause nervousness, irritability, sleeplessness, and, occasionally, rapid heart beat.

Do not use:

- if you're ever had an allergic reaction to acetaminophen, aspirin or any other pain reliever/fever reducer
- with any other drug containing acetaminophen (prescription or nonprescription), if you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist

Drug Facts (continued)

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not use more than directed
- adults: take 2 geltabs with a glass of water
- if symptoms persist or worsen, ask your doctor
- do not take more than 2 geltabs in 24 hours, unless directed by a doctor
- under 18 years of age: ask a doctor

Other information

- store at controlled room temperature 20°-25°C (68°-77°F)
- close cap tightly after use
- read all product information before using. Keep this box for important information.

Inactive ingredients benzoic acid, D&C yellow #10 lake, disodium EDTA, FD&C blue #1 lake, FD&C red #40 lake, ferric oxide, gelatin, glycerin, hydroxypropylcellulose, hypromellose, light mineral oil, maltitol solution, microcrystalline cellulose, pepsin, polysorbate 20, povidone, propylene glycol, propyl gallate, simethicone emulsion, sorbitan monolaurate, stearic acid, titanium dioxide

Questions or comments? 1-800-468-7746

NDC 0067-2035-91

EXCEDRIN®

MIGRAINE

Acetaminophen, Aspirin (NSAID) and Caffeine

Pain Reliever/Pain Reliever Aid

100 GELTABS®
*GELATIN COATED TABLETS

100 GELTABS IN TWO 50 COUNT BOTTLES

This is a representation of an electronic record that was signed electronically and this page is the manifestation of the electronic signature.

/s/

THERESA M MICHELE
07/15/2014