

See new warnings information NDC 0067-2039-02

EXCEDRIN[®]

MIGRAINE

Acetaminophen, Aspirin (NSAID) and Caffeine
Pain Reliever/Pain Reliever Aid 2 CAPLETS

Active ingredients (in each caplet)	Purposes
Acetaminophen 250 mg	Pain reliever
Aspirin 250 mg (NSAID)*	Pain reliever
Caffeine 65 mg	Pain reliever aid

(nonsteroidal anti-inflammatory drug) **Use** • treats migraine **Warnings** **Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: • skin reddening • blisters • rash. If a skin reaction occurs, stop use and seek medical help right away.

Allergy alert: Aspirin may cause a severe allergic reaction which may include: • hives • facial swelling • asthma (wheezing) • shock. **Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take • more than 2 caplets in 24 hours, which is the maximum daily amount • with other drugs containing acetaminophen • 3 or more alcoholic drinks every day while using this product. **Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you • are age 60 or older • have had stomach ulcers or bleeding problems • take a blood thinning (anticoagulant) or steroid drug • take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) • have 3 or more alcoholic drinks every day while using this product • take more or for a longer time than directed. **Medication overuse headache warning:** Headaches may worsen if this product is used for 10 or more days per month. **Caffeine warning:** The recommended dose of this product contains about as much caffeine as a cup of coffee. Limit the use of caffeine-containing medications, foods, or beverages while taking this product because too much caffeine may cause nervousness, irritability, sleeplessness, and, occasionally, rapid heart beat.

Do not use • if you have ever had an allergic reaction to acetaminophen, aspirin or

any other pain reliever/fever reducer • with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. **Ask a doctor before use if** • you have never had migraines diagnosed by a health professional • you have a headache that is different from your usual migraines • you have the worst headache of your life • you have fever and stiff neck • you have headaches beginning after or caused by head injury, exertion, coughing or bending • you experienced your first headache after the age of 50 • you have daily headaches • you have a migraine so severe as to require bed rest • you have liver disease • stomach bleeding warning applies to you • you have a history of stomach problems, such as heartburn • you have high blood pressure, heart disease, liver cirrhosis, or kidney disease • you are taking a diuretic • you have asthma • you have problems or serious side effects from taking pain relievers or fever reducers • you have vomiting with your migraine headache. **Ask a doctor or pharmacist before use if you are** • taking a prescription drug for: • diabetes • gout • arthritis • under a doctor's care for any serious condition • taking any other drug • taking any other product that contains aspirin, acetaminophen, or any other pain reliever/fever reducer. **Stop use and ask a doctor if** • an allergic reaction occurs. Seek medical help right away. • you experience any of the following signs of stomach bleeding: • feel faint • vomit blood • have bloody or black stools • have stomach pain that does not get better • your migraine is not relieved or worsens after first dose • new or unexpected symptoms occur • ringing in the ears or loss of hearing occurs. **If pregnant or breast-feeding,** ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms. **Directions** • do not use more than directed • adults: take 2 caplets with a glass of water • if symptoms persist or worsen, ask your doctor • do not take more than 2 caplets in 24 hours, unless directed by a doctor • under 18 years of age: ask a doctor. **Other information** • store at room temperature 20°-25°C (68°-77°F) • read all product information before using.

Inactive ingredients: benzoic acid, camellia wax, FD&C blue #1, hydroxypropylcellulose, hypromellose, light mineral oil, microcrystalline cellulose, poly sorbate 20, povidone, propylene glycol, simethicone emulsion, sorbitan monolaurate, stearic acid, titanium dioxide. **Questions? 1-800-452-0051**
Dist. by: GSK Consumer Healthcare, Warren, NJ 07059
©2017 GSK or its licensor. 167274-2


LINE AND TEAR AT SLIT
DO NOT USE IF SEAL IS BROKEN

TO OPEN, FOLD ALONG DOTTED LINE AND TEAR AT SLIT
TAMPER EVIDENT PACKET

LOT/EXP AREA

167274-2

DRUG FACT FORMATTING: MODIFIED		
DRUG FACTS TEXT	TYPE SIZE	TYPE STYLE
HEADINGS	4.5 pt.	Helvetica Cond Bold Obl
SUBHEADS	4.5 pt.	Helvetica Cond Bold
BODY TEXT	4.5 pt.	Helvetica Cond
TRACKING	0	
CONDENSATION	83%	
LEADING	4.54 pt.	
BULLETS	4 pt.	



DRUG FACT FORMATTING: MODIFIED		
DRUG FACTS TEXT	TYPE SIZE	TYPE STYLE
DRUG FACTS TITLE	9 pt.	Helvetica Cond Bold Obl
DRUG FACTS CONT.	8 pt.	Helvetica Cond Bold Obl
HEADINGS	8 pt.	Helvetica Cond Bold
SUBHEADS	6 pt.	Helvetica Cond
BODY TEXT	6 pt.	Helvetica Cond
TRACKING	0	
CONDENSATION	83%	
LEADING	6 pt.	
BARLINES, HAIRLINES	1.5, 0.5 pt.	
CHARACTERS/INCH	<39	
BULLETS	5 pt.	



DRUG FACT FORMATTING: MODIFIED

DRUG FACTS TEXT	TYPE SIZE	TYPE STYLE
DRUG FACTS TITLE	9 pt.	Helvetica Cond Bold Obl
DRUG FACTS CONT.	8 pt.	Helvetica Cond Bold Obl
HEADINGS	8 pt.	Helvetica Cond Bold
SUBHEADS	6 pt.	Helvetica Cond
BODY TEXT	6 pt.	Helvetica Cond
TRACKING	0	
CONDENSATION	83%	
LEADING	6.3 pt.	
BARLINES, HAIRLINES	1.5, 0.5 pt.	
CHARACTERS/INCH	<39	
BULLETS	5 pt.	

LOT/EXP AREA

3 0067 2039 91 0

UPC FPO

16/274-100 Visit us at www.excedrin.com
©2017 SSK Group of companies or its licensor. All rights reserved.
Distributed by: GSK Consumer Healthcare, Warren, NJ 07059

TAMPER-EVIDENT BOTTLE
DO NOT USE IF BUBBLE FOLK SEAL IMPRINTED WITH
"SEALED FOR YOUR PROTECTION" IS BROKEN OR MISSING

See new warnings information

EXCEDRIN[®]

MIGRAINE

Acetaminophen, Aspirin (NSAID) and Caffeine

Pain Reliever/Pain Reliever Aid

100 CAPLETS

CAPLETS SHOWN

NDC 0067-2039-91

46170648

CODE 128 FPO

46170648

46170648

CODE 128 FPO

46170648

46170648

CODE 128 FPO

46170648

46170648

CODE 128 FPO

46170648

46170648

CODE 128 FPO

46170648

46170648

CODE 128 FPO

46170648

Glue

Drug Facts

Active ingredients (in each caplet) **Purposes**

Acetaminophen 250 mg Pain reliever

Aspirin 250 mg (NSAID)¹ Pain reliever

Caffeine 65 mg Pain reliever aid

¹nonsteroidal anti-inflammatory drug

Use • treats migraine

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: • skin reddening • blisters • rash. If a skin reaction occurs, stop use and seek medical help right away.

Allergy alert: Aspirin may cause a severe allergic reaction which may include: • hives • facial swelling • asthma (wheezing) • shock

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 2 caplets in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Medication overuse headache warning: Headaches may worsen if this product is used for 10 or more days per month.

Caffeine warning: The recommended dose of this product contains about as much caffeine as a cup of coffee. Limit the use of caffeine-containing medications, foods, or beverages while taking this product because too much caffeine may cause nervousness, irritability, sleeplessness, and, occasionally, rapid heart beat.

Do not use • if you have ever had an allergic reaction to acetaminophen, aspirin or any other pain reliever/fever reducer • with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist

Ask a doctor before use if

- you have never had migraines diagnosed by a health professional
- you have a headache that is different from your usual migraines
- you have the worst headache of your life
- you have fever and stiff neck
- you have headaches beginning after or caused by head injury, exertion, coughing or bending
- you experienced your first headache after the age of 50
- you have daily headaches
- you have a migraine so severe as to require bed rest
- you have liver disease
- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have vomiting with your migraine headache

Ask a doctor or pharmacist before use if you are

- taking a prescription drug for: • diabetes • gout • arthritis
- under a doctor's care for any serious condition
- taking any other drug
- taking any other product that contains aspirin, acetaminophen, or any other pain reliever/fever reducer

Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
- your migraine is not relieved or worsens after first dose
- new or unexpected symptoms occur
- ringing in the ears or loss of hearing occurs

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions • do not use more than directed

- adults: take 2 caplets with a glass of water

Drug Facts (continued)

Other information

- store at controlled room temperature 20°-25°C (68°-77°F)
- dose cap tightly after use
- read all product information before using. Keep this box for important information.

Other ingredients benzoic acid, camphor wax, E130 Blue #1, hydroxypropylcellulose, hypromellose, light mineral oil, microcrystalline cellulose, polyethylene glycol, propylene glycol, stearic acid, sodium monohydrogen phosphate, titanium dioxide

Questions? 1-800-468-7746

DRUG FACT FORMATTING: STANDARD		
DRUG FACTS TEXT	TYPE SIZE	TYPE STYLE
DRUG FACTS TITLE	9 pt.	Helvetica Cond Bold Obl
DRUG FACTS CONT.	8 pt.	Helvetica Cond Bold Obl
HEADINGS	8 pt.	Helvetica Cond Bold
SUBHEADS	6 pt.	Helvetica Cond
BODY TEXT	6 pt.	Helvetica Cond
TRACKING	0	
CONDENSATION	83%	
LEADING	6.5 pt.	
BARLINES, HAIRLINES	1.5, 0.5 pt.	
CHARACTERS/INCH	<39	
BULLETS	5 pt.	

LOT/EXP AREA

3 0067-2039-83 5

FP0

167274-125 Visit us at www.excedrin.com
©2017 GSK Group of Companies or its licensor. All rights reserved.
Distributed by GSK Consumer Healthcare, Warren, NJ 07059

gsk

46170263

46170263

46170263

46170263

46170263

46170263

46170263

46170263

46170263

46170263

46170263

46170263

DO NOT USE IF INNER FOIL SEAL IMPRINTED WITH "SEALED BY YOUR PROTECTOR" IS BROKEN OR MISSING

TAMPER-EVIDENT BOTTLE

See new warnings information

EXCEDRIN

MIGRAINE

Acetaminophen, Aspirin (NSAID)
and Caffeine

Pain Reliever / Pain Reliever Aid

125
100
CAPLETS

25%
MORE
FREE

NDC 0067-2039-83

See new warnings information

EXCEDRIN

MIGRAINE

Acetaminophen, Aspirin (NSAID)
and Caffeine

Pain Reliever / Pain Reliever Aid

125
100
CAPLETS

25%
MORE
FREE

NDC 0067-2039-83

Drug Facts

Active ingredients (in each caplet)

Acetaminophen 250 mg Pain reliever
Aspirin 250 mg (NSAID)* Pain reliever
Caffeine 65 mg Pain reliever aid
*nonsteroidal anti-inflammatory drug

Use • treats migraine

Warnings
Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.
Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: • skin reddening • blisters • rash
If a skin reaction occurs, stop use and seek medical help right away.
Allergy alert: Aspirin may cause a severe allergic reaction which may include: • hives • facial swelling • asthma (wheezing) • shock
Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take
• more than 2 caplets in 24 hours, which is the maximum daily amount
• with other drugs containing acetaminophen
• 3 or more alcoholic drinks every day while using this product
Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you
• are age 60 or older • have had stomach ulcers or bleeding problems
• take a blood thinning (anticoagulant) or steroid drug
• take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
• have 3 or more alcoholic drinks every day while using this product
• take more or for a longer time than directed
Medication overuse headache warning: Headaches may worsen if this product is used for 10 or more days per month.
Caffeine warning: The recommended dose of this product contains about as much caffeine as a cup of coffee. Limit the use of caffeine-containing medications, foods, or beverages while taking this product because too much caffeine may cause nervousness, irritability, sleeplessness, and, occasionally, rapid heart beat.
Do not use • if you have ever had an allergic reaction to acetaminophen, aspirin or any other pain reliever/fever reducer
• with any other drug containing acetaminophen (prescription or

Drug Facts (continued)

nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist

Ask a doctor before use if

- you have never had migraines diagnosed by a health professional
- you have a headache that is different from your usual migraines
- you have the worst headache of your life
- you have fever and stiff neck
- you have headaches beginning after or caused by head injury, exertion, coughing or bending
- you experienced your first headache after the age of 50
- you have daily headaches
- you have a migraine so severe as to require bed rest
- you have liver disease • stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease • you are taking a diuretic • you have asthma
- you have problems or serious side effects from taking pain relievers or fever reducers • you have vomiting with your migraine headache

Ask a doctor or pharmacist before use if you are

- taking a prescription drug for: • diabetes • gout • arthritis
- under a doctor's care for any serious condition
- taking any other drug
- taking any other product that contains aspirin, acetaminophen, or any other pain reliever/fever reducer

Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
 - feel faint • vomit blood • have bloody or black stools
 - have stomach pain that does not get better
 - your migraine is not relieved or worsens after first dose
 - new or unexpected symptoms occur
 - ringing in the ears or loss of hearing occurs

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.
In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions • do not use more than directed
• adults: take 2 caplets with a glass of water

Drug Facts (continued)

- if symptoms persist or worsen, ask your doctor
- do not take more than 2 caplets in 24 hours, unless directed by a doctor
- under 13 years of age, ask a doctor

Other information

- store at room temperature 20°-25°C (68°-77°F)
- close cap tightly after use
- read all product information before using. Keep this box for important information.

Inactive ingredients: benzoic acid, carnauba wax, FD&C Blue #1, hydroxypropylcellulose, hypromellose, light mineral oil, microcrystalline cellulose, polyethylene glycol, polyethylene glycol, stearic acid, stearic acid, titanium dioxide

Questions? 1-800-452-0061

DRUG FACTS FORMATTING: STANDARD		
DRUG FACTS TEXT	TYPE SIZE	TYPE STYLE
DRUG FACTS TITLE	9 pt.	Helvetica Cond Bold Obl
DRUG FACTS CONT.	8 pt.	Helvetica Cond Bold Obl
HEADINGS	8 pt.	Helvetica Cond Bold
SUBHEADS	6 pt.	Helvetica Cond
BODY TEXT	6 pt.	Helvetica Cond
TRACKING	0	
CONDENSATION	83%	
LEADING	6.5 pt.	
BARLINES, HAIRLINES	1.5, 0.5 pt.	
CHARACTERS/INCH	<39	
BULLETS	5 pt.	

LOT/EXP AREA

3 0067 2039 92 7

UFC FPO

16724-200 Visit us at www.excedrin.com
©2017 GSK group of companies or its licensor. All rights reserved.
GSK

4-6170720

CODE 128 FPO

4-6170720

CODE 128 FPO

4-6170720

Glue

4-6170720

4-6170720

4-6170720

See new warnings information

NDC 0067-2039-92

EXCEDRIN

MIGRAINE

Acetaminophen, Aspirin (NSAID) and Caffeine

Pain Reliever/Pain Reliever Aid

200 CAPLETS[®]

*CAPULET-SHAPED TABLETS

See new warnings information

NDC 0067-2039-92

EXCEDRIN

MIGRAINE

Acetaminophen, Aspirin (NSAID) and Caffeine

Pain Reliever/Pain Reliever Aid

200 CAPLETS[®]

*CAPULET-SHAPED TABLETS

TAMPER-EVIDENT BOTTLE
DO NOT USE IF INNER FOIL SEAL IMPRINTED WITH
"SEALED for YOUR PROTECTION" IS BROKEN OR MISSING

Drug Facts

Active ingredients (in each caplet)

Active ingredients (in each caplet)	Purposes
Acetaminophen 250 mg	Pain reliever
Aspirin 250 mg (NSAID) [*]	Pain reliever
Caffeine 65 mg	Pain reliever aid

^{*}nonsteroidal anti-inflammatory drug

Use • treats migraine

Warnings
Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.
Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:
• skin reddening • blisters • rash
If a skin reaction occurs, stop use and seek medical help right away.
Allergy alert: Aspirin may cause a severe allergic reaction which may include:
• hives • facial swelling • asthma (wheezing) • shock
Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take
• more than 2 caplets in 24 hours, which is the maximum daily amount
• with other drugs containing acetaminophen
• 3 or more alcoholic drinks every day while using this product
Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you • are age 60 or older
• have had stomach ulcers or bleeding problems
• take a blood thinning (anticoagulant) or steroid drug
• take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
• have 3 or more alcoholic drinks every day while using this product
• take more or for a longer time than directed
Medication overuse headache warning: Headaches may worsen if this product is used for 10 or more days per month.
Caffeine warning: The recommended dose of this product contains about as much caffeine as a cup of coffee. Limit the use of caffeine-containing medications, foods, or beverages while taking this product because too much caffeine may cause nervousness, irritability, sleeplessness, and, occasionally, rapid heart beat.
Do not use • if you have ever had an allergic reaction to acetaminophen, aspirin or any other pain reliever/fever reducer
• with any other drug containing acetaminophen (prescription or nonprescription).
If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist
Ask a doctor before use if
• you have never had migraines diagnosed by a health professional

Drug Facts (continued)

- you have a headache that is different from your usual migraines
- you have the worst headache of your life
- you have fevers and stiff neck
- you have headaches beginning after or caused by head injury, exertion, coughing or bending
- you experienced your first headache after the age of 50
- you have daily headaches
- you have a migraine so severe as to require bed rest
- you have liver disease
- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have vomiting with your migraine headache

Ask a doctor or pharmacist before use if you are

- taking a prescription drug for: • diabetes • gout • arthritis
- under a doctor's care for any serious condition
- taking any other drug
- taking any other product that contains aspirin, acetaminophen, or any other pain reliever/fever reducer

Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
- your migraine is not relieved or worsens after first dose
- new or unexpected symptoms occur
- ringing in the ears or loss of hearing occurs

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.
Keep out of reach of children.
In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions • do not use more than directed

- adults: take 2 caplets with a glass of water
- if symptoms persist or worsen, ask your doctor
- do not take more than 2 caplets in 24 hours, unless directed by a doctor
- under 18 years of age: ask a doctor

Other information • store at controlled room temperature 20°-25°C (68°-77°F)
• close cap tightly after use
• read all product information before using. Keep this box for important information.

Inactive ingredients benzoic acid, carnauba wax, FD&C blue #1, hydroxypropylcellulose, hypromellose, light mineral oil, microcrystalline cellulose, poly sorbate 20, povidone, propylene glycol, simethicone emulsion, sorbitan monolaurate, stearic acid, titanium dioxide

Questions? 1-800-468-7746

DRUG FACT FORMATTING: STANDARD		
DRUG FACTS TEXT	TYPE SIZE	TYPE STYLE
DRUG FACTS TITLE	9 pt.	Helvetica Cond Bold Obi
DRUG FACTS CONT.	8 pt.	Helvetica Cond Bold Obi
HEADINGS	8 pt.	Helvetica Cond Bold
SUBHEADS	6 pt.	Helvetica Cond
BODY TEXT	6 pt.	Helvetica Cond
TRACKING	0	
CONDENSATION	83%	
LEADING	6.5 pt.	
BARLINES, HAIRLINES	1.5, 0.5 pt.	
CHARACTERS/INCH	<39	
BULLETS	5 pt.	

LOT/EXP AREA



Distributed by: GSK Consumer Healthcare, Warren, NJ 07059
© 2017 GSK group of companies or its licensor. All rights reserved.
167274-250 Visit us at www.excedrin.com

46170647



46170647



See new warnings information NDC 0067-2039-77

EXCEDRIN
MIGRAINE
Acetaminophen,
Aspirin (NSAID) and Caffeine
Pain Reliever/Pain Reliever Aid

25% MORE FREE

250 200 CAPLETS
*CAPSULE-SHAPED TABLETS

See new warnings information
EXCEDRIN
Acetaminophen, Aspirin (NSAID) and Caffeine
Pain Reliever/Pain Reliever Aid
NDC 0067-2039-77
250 200 CAPLETS
*CAPSULE-SHAPED TABLETS

25% MORE FREE

See new warnings information
EXCEDRIN
Acetaminophen, Aspirin (NSAID) and Caffeine
Pain Reliever/Pain Reliever Aid
NDC 0067-2039-77
250 200 CAPLETS
*CAPSULE-SHAPED TABLETS

25% MORE FREE

TAMPER-EVIDENT BOTTLE
DO NOT USE IF INNER FOIL SEAL IMPRINTED WITH
"SEALED FOR YOUR PROTECTION" IS BROKEN OR MISSING

Drug Facts
Active ingredients (in each caplet)
Acetaminophen 250 mg Pain reliever
Aspirin 250 mg (NSAID) Pain reliever
Caffeine 65 mg Pain reliever aid
*nonsteroidal anti-inflammatory drug

Purposes
Pain reliever
Pain reliever
Pain reliever aid

Use • treats migraine

Warnings
Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.
Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:
• skin reddening • blisters • rash
If a skin reaction occurs, stop use and seek medical help right away.
Allergy alert: Aspirin may cause a severe allergic reaction which may include:
• hives • facial swelling • asthma (wheezing) • shock
Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take
• more than 2 caplets in 24 hours, which is the maximum daily amount
• with other drugs containing acetaminophen
• 3 or more alcoholic drinks every day while using this product
Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you • are age 60 or older
• have had stomach ulcers or bleeding problems
• take a blood thinning (anticoagulant) or steroid drug
• take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
• have 3 or more alcoholic drinks every day while using this product
• take more or for a longer time than directed
Medication overuse headache warning: Headaches may worsen if this product is used for 10 or more days per month.
Caffeine warning: The recommended dose of this product contains about as much caffeine as a cup of coffee. Limit the use of caffeine-containing medications, foods, or beverages while taking this product because too much caffeine may cause nervousness, irritability, sleeplessness, and, occasionally, rapid heart beat.

Do not use • if you have ever had an allergic reaction to acetaminophen, aspirin or any other pain reliever/fever reducer
• with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist

Ask a doctor before use if
• you have never had migraines diagnosed by a health professional



46170647

Drug Facts (continued)
• you have a headache that is different from your usual migraines
• you have the worst headache of your life • you have fever and stiff neck
• you have headaches beginning after or caused by head injury, exertion, coughing or bending
• you experienced your first headache after the age of 50
• you have daily headaches • you have a migraine so severe as to require bed rest
• you have liver disease • stomach bleeding warning applies to you
• you have a history of stomach problems, such as heartburn
• you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
• you are taking a diuretic • you have asthma
• you have problems or serious side effects from taking pain relievers or fever reducers
• you have vomiting with your migraine headache

Ask a doctor or pharmacist before use if you are
• taking a prescription drug for: • diabetes • gout • arthritis
• under a doctor's care for any serious condition • taking any other drug
• taking any other product that contains aspirin, acetaminophen, or any other pain reliever/fever reducer

Stop use and ask a doctor if • an allergic reaction occurs. Seek medical help right away.
• you experience any of the following signs of stomach bleeding:
• feel faint • vomit blood • have bloody or black stools
• have stomach pain that does not get better
• your migraine is not relieved or worsens after first dose
• new or unexpected symptoms occur • ringing in the ears or loss of hearing occurs

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.
Keep out of reach of children.
In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions • do not use more than directed
• adults: take 2 caplets with a glass of water
• if symptoms persist or worsen, ask your doctor
• do not take more than 2 caplets in 24 hours, unless directed by a doctor
• under 18 years of age: ask a doctor

Other information • store at room temperature 20°-25°C (68°-77°F)
• close cap tightly after use
• read all product information before using. Keep this box for important information.

Inactive ingredients benzoic acid, carnauba wax, FD&C blue #1, hydroxypropylcellulose, hypromellose, light mineral oil, microcrystalline cellulose, polysorbate 20, povidone, propylene glycol, simethicone emulsion, sorbitan monolaurate, stearic acid, titanium dioxide

Questions? 1-800-452-0051



46170647

DRUG FACT FORMATTING: STANDARD		
DRUG FACTS TEXT	TYPE SIZE	TYPE STYLE
DRUG FACTS TITLE	9 pt.	Helvetica Cond Bold Obl
DRUG FACTS CONT.	8 pt.	Helvetica Cond Bold Obl
HEADINGS	8 pt.	Helvetica Cond Bold
SUBHEADS	6 pt.	Helvetica Cond
BODY TEXT	6 pt.	Helvetica Cond
TRACKING	0	
CONDENSATION	83%	
LEADING	6.5 pt.	
BARLINES, HAIRLINES	1.5, 0.5 pt.	
CHARACTERS/INCH	<39	
BULLETS	5 pt.	

LOT/EXP AREA



Distributed by GSK Consumer Healthcare, Warren, NJ 07059
©2017 GSK group of companies or its licensor. All rights reserved.
167274-300 Visit us at www.excedrin.com



46170261



46170261



TAMPER-EVIDENT BOTTLE
DO NOT USE IF INNER FOIL SEAL IMPRINTED WITH
"SEALED for YOUR PROTECTION" IS BROKEN OR MISSING

Active ingredients (in each caplet)	Purposes
Acetaminophen 250 mg	Pain reliever
Aspirin 250 mg (NSAID)	Pain reliever
Caffeine 65 mg	Pain reliever aid

Use • treats migraine

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: • skin reddening • blisters • rash
If a skin reaction occurs, stop use and seek medical help right away.

Allergy alert: Aspirin may cause a severe allergic reaction which may include: • hives • facial swelling • asthma (wheezing) • shock

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 2 caplets in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen

• 3 or more alcoholic drinks every day while using this product
Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Medication overuse headache warning: Headaches may worsen if this product is used for 10 or more days per month.

Caffeine warning: The recommended dose of this product contains about as much caffeine as a cup of coffee. Limit the use of caffeine-containing medications, foods, or beverages while taking this product because too much caffeine may cause nervousness, irritability, sleeplessness, and, occasionally, rapid heart beat.

Do not use

- if you have ever had an allergic reaction to acetaminophen, aspirin or any other pain reliever/fever reducer
 - with any other drug containing acetaminophen (prescription or nonprescription).
- If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

Ask a doctor before use if

- you have never had migraines diagnosed by a health professional
- you have a headache that is different from your usual migraines
- you have the worst headache of your life
- you have fever and stiff neck

Drug Facts (continued)

- you have headaches beginning after or caused by head injury, exertion, coughing or bending
- you experienced your first headache after the age of 50
- you have daily headaches
- you have a migraine so severe as to require bed rest
- you have liver disease
- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have vomiting with your migraine headache

Ask a doctor or pharmacist before use if you are

- taking a prescription drug for: • diabetes • gout • arthritis
- under a doctor's care for any serious condition
- taking any other drug
- taking any other product that contains aspirin, acetaminophen, or any other pain reliever/fever reducer

Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
 - feel faint • vomit blood • have bloody or black stools
 - have stomach pain that does not get better
- your migraine is not relieved or worsens after first dose
- new or unexpected symptoms occur
- ringing in the ears or loss of hearing occurs

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not use more than directed
- adults: take 2 caplets with a glass of water
- if symptoms persist or worsen, ask your doctor
- do not take more than 2 caplets in 24 hours, unless directed by a doctor
- under 18 years of age: ask a doctor

Other information

- store at controlled room temperature 20°-25°C (68°-77°F)
- close cap tightly after use
- read all product information before using. Keep this box for important information.

Inactive ingredients benzoic acid, carnauba wax, FD&C blue #1, hydroxypropylcellulose, hypromellose, light mineral oil, microcrystalline cellulose, polysorbate 20, povidone, propylene glycol, simethicone emulsion, sorbitan monolaurate, stearic acid, titanium dioxide

Questions? 1-800-468-7746

Glue




46170261




46170261

DRUG FACT FORMATTING- STANDARD		
DRUG FACTS TEXT	TYPE SIZE	TYPE STYLE
DRUG FACTS TITLE	9 pt.	Helvetica Cond Bold Obi
DRUG FACTS CONT.	8 pt.	Helvetica Cond Bold Obi
HEADINGS	8 pt.	Helvetica Cond Bold
SUBHEADS	6 pt.	Helvetica Cond
BODY TEXT	6 pt.	Helvetica Cond
TRACKING	0	
CONDENSATION	83%	
LEADING	6.5 pt.	
BARLINES, HAIRLINES	1.5, 0.5 pt.	
CHARACTERS/INCH	<39	
BULLETS	5 pt.	



DRUG FACT FORMATTING: MODIFIED		
DRUG FACTS TEXT	TYPE SIZE	TYPE STYLE
DRUG FACTS TITLE	7.5 pt.	Helvetica Cond Bold Obl
DRUG FACTS CONT.	7 pt.	Helvetica Cond Bold Obl
HEADINGS	7 pt.	Helvetica Cond Bold
SUBHEADS	6 pt.	Helvetica Cond
BODY TEXT	6 pt.	Helvetica Cond
TRACKING	0	
CONDENSATION	83%	
LEADING	5.8 pt.	
BARLINES, HAIRLINES	1.5, 0.5 pt.	
CHARACTERS/INCH	<39	
BULLETS	5 pt.	



LOT/EXP AREA

3 0067-2035-80 6



UPG FPO

Distributed by: GSK Consumer Healthcare, Warren, NJ 07059
©2017 GSK group of companies or its licensor. All rights reserved.
167287-80 Visit us at www.excedrin.com

46170725



CODE 128 FPO

46170725



CODE 128 FPO

TAMPER-EVIDENT BOTTLE
DO NOT USE IF INNER FOIL SEAL IMPRINTED WITH
"SEALED FOR YOUR PROTECTION" IS BROKEN OR MISSING

Drug Facts

Active ingredients (in each GelTab) **Purposes**
Acetaminophen 250 mg.....Pain reliever
Aspirin 250 mg (NSAID).....Pain reliever
Caffeine 65 mg.....Pain reliever aid
*nonsteroidal anti-inflammatory drug

Use • treats migraine

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.
Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: • skin reddening • blisters • rash
If a skin reaction occurs, stop use and seek medical help right away.
Allergy alert: Aspirin may cause a severe allergic reaction which may include:

- hives • facial swelling • asthma (wheezing) • shock

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 2 gels in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Medication overuse headache warning: Headaches may worsen if this product is used for 10 or more days per month.

Caffeine warning: The recommended dose of this product contains about as much caffeine as a cup of coffee. Limit the use of caffeine-containing medications, foods, or beverages while taking

Drug Facts (continued)

this product because too much caffeine may cause nervousness, irritability, sleeplessness, and, occasionally, rapid heart beat.

Do not use

- if you have ever had an allergic reaction to acetaminophen, aspirin or any other pain reliever/fever reducer
- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist

Ask a doctor before use if

- you have never had migraines diagnosed by a health professional
- you have a headache that is different from your usual migraines
- you have the worst headache of your life
- you have fever and stiff neck
- you have headaches beginning after or caused by head injury, exertion, coughing or bending
- you experienced your first headache after the age of 50
- you have daily headaches
- you have a migraine so severe as to require bed rest
- you have liver disease
- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have vomiting with your migraine headache

Ask a doctor or pharmacist before use if you are

- taking a prescription drug for: • diabetes • gout • arthritis
- under a doctor's care for any serious condition
- taking any other drug
- taking any other product that contains aspirin, acetaminophen, or any other pain reliever/fever reducer

Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
 - feel faint • vomit blood • have bloody or black stools
 - have stomach pain that does not get better
- your migraine is not relieved or worsens after first dose
- new or unexpected symptoms occur
- ringing in the ears or loss of hearing occurs

Drug Facts (continued)

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not use more than directed
- adults: take 2 gels with a glass of water
- if symptoms persist or worsen, ask your doctor
- do not take more than 2 gels in 24 hours, unless directed by a doctor
- under 18 years of age: ask a doctor

Other information

- store at controlled room temperature 20°-25°C (68°-77°F)
- close cap tightly after use
- read all product information before using. Keep this box for important information.

Inactive ingredients

benzoic acid, D&C yellow #10 aluminum lake, edetate disodium, FD&C blue #1, FD&C blue #1 aluminum lake, FD&C red #40 aluminum lake, gelatin, glycerin, hypromellose, iron oxide black, isopropyl alcohol, lactose monohydrate, light mineral oil, low-substituted hydroxypropyl cellulose, maltitol syrup, medium-chain triglycerides, microcrystalline cellulose, pepsin, polysorbate 20, povidone, propyl gallate, propylene glycol, simethicone, sorbitan monolaurate, stearic acid, titanium dioxide

Questions?

1-800-452-0051


EXCEDRIN
MIGRAINE
Acetaminophen, Aspirin (NSAID) and Caffeine
Pain Reliever/Pain Reliever Aid
80 GELTABS
GELTABS

CODE 128 FPO



46170725

CODE 128 FPO



46170725

DRUG FACT FORMATTING- STANDARD		
DRUG FACTS TEXT	TYPE SIZE	TYPE STYLE
DRUG FACTS TITLE	9 pt.	Helvetica Cond Bold Obl
DRUG FACTS CONT.	8 pt.	Helvetica Cond Bold Obl
HEADINGS	8 pt.	Helvetica Cond Bold
SUBHEADS	6 pt.	Helvetica Cond
BODY TEXT	6 pt.	Helvetica Cond
TRACKING	0	
CONDENSATION	83%	
LEADING	6.5 pt.	
BARLINES, HAIRLINES	1.5, 0.5 pt.	
CHARACTERS/INCH	<39	
BULLETS	5 pt.	