

EXCEDRIN SAFETY SEALED • EXCEDRIN SAFETY SEALED
TO OPEN PUSH DOWN THEN TURN → TO OPEN PUSH DOWN THEN TURN →

Drug Facts (continued)

- you have fever and stiff neck
- you have headaches beginning after or caused by head injury, exertion, coughing or bending
- you experienced your first headache after the age of 50
- you have daily headaches
- you have a migraine so severe as to require bed rest
- you have liver disease
- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have vomiting with your migraine headache

Ask a doctor or pharmacist before use if you are

- taking a prescription drug for:
 - diabetes
 - gout
 - arthritis
- under a doctor's care for any serious condition
- taking any other drug
- taking any other product that contains aspirin, acetaminophen, or any other pain reliever/fever reducer

Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
- your migraine is not relieved or worsens after first dose
- new or unexpected symptoms occur
- ringing in the ears or loss of hearing occurs

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions • do not use more than directed

- adults: take 2 caplets with a glass of water
- if symptoms persist or worsen, ask your doctor
- do not take more than 2 caplets in 24 hours, unless directed by a doctor
- under 18 years of age: ask a doctor

Other information

- store at controlled room temperature 20°–25°C (68°–77°F)
- close cap tightly after use
- read all product information before using

Inactive ingredients benzoic acid, carnauba wax, FD&C blue no. 1, hydroxypropyl cellulose, low-substituted hydroxypropyl cellulose, microcrystalline cellulose, polyborate 20, povidone, propylene glycol, simethicone, sorbitan monolaurate, stearic acid, titanium dioxide

Questions? 1-800-468-7746

Distributed by: GSK Consumer Healthcare, Warren, NJ 07059
©2021 GSK group of companies or its licensee.
Ingredients not verified or licensed to the GSK group of companies.

DO NOT USE IF EXCEDRIN SAFETY SEAL AROUND CAP IS BROKEN OR MISSING.

FPO

NDC 0000-0000-00

EXCEDRIN
MIGRAINE
Acetaminophen, Aspirin (ASA) and Caffeine
Pain Reliever/Pain Reliever AS

Drug Facts

Active ingredients (in each caplet)	Purposes
Acetaminophen 250 mg	Pain reliever
Aspirin 250 mg (NSAID)	Pain reliever
Caffeine 65 mg	Pain reliever aid
monoclonal anti-inflammatory drug	

PEEL RIGHT CORNER FOR FULL DRUG FACTS ▲ FEEL HERE

Drug Facts (continued)

Use • treats migraine

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Allergy alert: Aspirin may cause a severe allergic reaction which may include:

- hives
- facial swelling
- asthma (wheezing)
- shock

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 2 caplets in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Medication overdose headache warning: Headaches may worsen if this product is used for 10 or more days per month.

Caffeine warning: The recommended dose of this product contains about as much caffeine as a cup of coffee. Limit the use of caffeine-containing medications, foods, or beverages while taking this product because too much caffeine may cause nervousness, irritability, sleeplessness, and, occasionally, rapid heart beat.

Do not use

- if you have ever had an allergic reaction to acetaminophen, aspirin or any other pain reliever/fever reducer
- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

Ask a doctor before use if

- you have never had migraines diagnosed by a health professional
- you have a headache that is different from your usual migraines
- you have the worst headache of your life

GSK Regulatory Spec Box		Verified Date: 05.07.21
Drug Facts Info		
Drug Facts	Helvetica Neue LT Std Bold Cond Oblique	9 pt.
Drug Facts (continued)	Helvetica Neue LT Std Bold Cond Oblique / Cond	8 pt.
Headings	Helvetica Neue LT Std Bold Cond Oblique	8 pt.
Subheadings	Helvetica Neue LT Std Bold Cond	6 pt.
Body text	Helvetica Neue LT Std Bold Cond / Cond	6 pt.
Bullets	Helvetica Neue LT Std Roman	5 pt.
Bullets on same lines: end of statement separated from bulleted statement by two ems		Yes
Spacing of the hair lines from edge of box -i.e. Minimum of 2 spaces either side of Drug Fact Box		Yes
Tracking:	0 pt. Horizontal Scale:	93%
Leading:	6 pt. Maximum Characters/Inch:	< 39
Barlines:	1.5 pt. Hairlines:	0.5 pt.
Primary Display Panel Info		
Size of Net Wt/Contents (Smallest character height in inches)*		>0.0625 in.
PDP dimensions (in square inches)		< 5 sq. in.
Ratio of Statement of Identity to Logo/Largest Copy on PDP		
Statement of Identity (pt.) (Divided by) Largest Logo Copy (pt.)		
	Vertical PDP: N/A	Horizontal PDP: 42%


NDC 0000-0000-00

EXCEDRIN

MIGRAINE

Acetaminophen, Aspirin (NSAID) and Caffeine
Pain Reliever/Pain Reliever Aid

10
CAPLETS[®]
*CAPSULE-SHAPED TABLETS




Drug Facts	Purposes	Drug Facts (continued)
Active ingredients (in each caplet)		
Acetaminophen 250 mg.....	Pain reliever	
Aspirin 250 mg (NSAID)*.....	Pain reliever	
Caffeine 65 mg.....	Pain reliever aid	
<small>*nonsteroidal anti-inflammatory drug</small>		
Use		
• treats migraine		
Warnings		
Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.		
Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: • skin reddening • blisters • rash If a skin reaction occurs, stop use and seek medical help right away.		
Allergy alert: Aspirin may cause a severe allergic reaction which may include: • hives • facial swelling • asthma (wheezing) • shock		
Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take • more than 2 caplets in 24 hours, which is the maximum daily amount • with other drugs containing acetaminophen • 3 or more alcoholic drinks every day while using this product		
Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you • are age 60 or older • have had stomach ulcers or bleeding problems • take a blood thinning (anticoagulant) or steroid drug • take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) • have 3 or more alcoholic drinks every day while using this product • take more or for a longer time than directed		

NDC 0000-0000-00

FPO

0 00000 00000 0

DO NOT USE IF EXCEDRIN SAFETY SEAL AROUND CAP IS BROKEN OR MISSING.

Drug Facts (continued)	Drug Facts (continued)
Medication overuse headache warning: Headaches may worsen if this product is used for 10 or more days per month.	• your migraine is not relieved or worsens after first dose
Caffeine warning: The recommended dose of this product contains about as much caffeine as a cup of coffee. Limit the use of caffeine-containing medications, foods, or beverages while taking this product because too much caffeine may cause nervousness, irritability, sleeplessness, and, occasionally, rapid heart beat.	• new or unexpected symptoms occur
Do not use	• ringing in the ears or loss of hearing occurs
• if you have ever had an allergic reaction to acetaminophen, aspirin or any other pain reliever/fever reducer	If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.
• with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.	Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.
Ask a doctor before use if	Directions
• you have never had migraines diagnosed by a health professional	• do not use more than directed
• you have a headache that is different from your usual migraines	• adults: take 2 caplets with a glass of water
• you have the worst headache of your life	• if symptoms persist or worsen, ask your doctor
• you have fever and stiff neck	• do not take more than 2 caplets in 24 hours, unless directed by a doctor
• you have headaches beginning after or caused by head injury, exertion, coughing or bending	• under 18 years of age: ask a doctor
• you experienced your first headache after the age of 50	Other information
• you have daily headaches	• store at controlled room temperature 20°-25°C (68°-77°F)
• you have a migraine so severe as to require bed rest	• close cap tightly after use
• you have liver disease	• read all product information before using. Keep this card for important information.
• stomach bleeding warning applies to you	Inactive ingredients benzoic acid, carnauba wax, FD&C blue no. 1, hypromellose, light mineral oil,
• you have a history of stomach problems, such as heartburn	low-substituted hydroxypropyl cellulose, microcrystalline cellulose, polysorbate 20, povidone, propylene glycol, simethicone, sorbitan monolaurate, stearic acid, titanium dioxide
• you have high blood pressure, heart disease, liver cirrhosis, or kidney disease	
• you are taking a diuretic	Questions? 1-800-468-7746
• you have asthma	
• you have problems or serious side effects from taking pain relievers or fever reducers	Distributed by: GSK Consumer Healthcare Warren, NJ 07059 ©2021 GSK or licensor. Trademarks owned or licensed by GSK. NPA004384
• you have vomiting with your migraine headache	
Ask a doctor or pharmacist before use if you are	
• taking a prescription drug for: • diabetes • gout • arthritis	
• under a doctor's care for any serious condition	
• taking any other drug	
• taking any other product that contains aspirin, acetaminophen, or any other pain reliever/fever reducer	
Stop use and ask a doctor if	
• an allergic reaction occurs. Seek medical help right away.	
• you experience any of the following signs of stomach bleeding: • feel faint • vomit blood • have bloody or black stools • have stomach pain that does not get better	

GSK Regulatory Spec Box		Verified Date: 05.10.21
Drug Facts Info		
Drug Facts	Helvetica Neue LT Std Bold Cond Oblique	9 pt.
Drug Facts (continued)	Helvetica Neue LT Std Bold Cond Oblique / Cond	8 pt.
Headings	Helvetica Neue LT Std Bold Cond Oblique	8 pt.
Subheadings	Helvetica Neue LT Std Bold Cond	6 pt.
Body text	Helvetica Neue LT Std Bold Cond / Cond	6 pt.
Bullets	Helvetica Neue LT Std Roman	5 pt.
Bullets on same lines: end of statement separated from bulleted statement by two ems		Yes
Spacing of the hair lines from edge of box -i.e. Minimum of 2 spaces either side of Drug Fact Box		Yes
Tracking:	0	Horizontal Scale: 85%
Leading:	6.25 pt.	Maximum Characters/Inch: < 39
Barlines:	1.5 pt.	Hairlines: 0.5 pt.
Primary Display Panel Info		
Size of Net Wt/Contents (Smallest character height in inches)*		> 0.125 in.
PDP dimensions (in square inches)		5-25 sq. in.
Ratio of Statement of Identity to Logo/Largest Copy on PDP Statement of Identity (pt.) (Divided by) Largest Logo Copy (pt.)		
	Vertical PDP: N/A	Horizontal PDP: 35%

This is a representation of an electronic record that was signed electronically. Following this are manifestations of any and all electronic signatures for this electronic record.

/s/

NUSHIN F TODD
12/07/2021 12:28:03 PM