


# EXCEDRIN

## MIGRAINE


Acetaminophen, Aspirin (NSAID) and Caffeine  
Pain Reliever/Pain Reliever Aid

**8**  
CAPLETS<sup>®</sup>


\*CAPSULE-SHAPED TABLETS




<p><b>Drug Facts</b></p> <p><b>Active ingredients (in each caplet)</b></p> <p>Acetaminophen 250 mg..... Pain reliever Aspirin 250 mg (NSAID)*..... Pain reliever Caffeine 65 mg..... Pain reliever aid <small>*nonsteroidal anti-inflammatory drug</small></p> <p><b>Use</b></p> <ul style="list-style-type: none"> <li>treats migraine</li> </ul> <p><b>Warnings</b></p> <p><b>Reye's syndrome:</b> Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.</p> <p><b>Allergy alert:</b> Acetaminophen may cause severe skin reactions. Symptoms may include:</p> <ul style="list-style-type: none"> <li>skin reddening</li> <li>blisters</li> <li>rash</li> </ul> <p>If a skin reaction occurs, stop use and seek medical help right away.</p>	<p><b>Drug Facts (continued)</b></p> <p><b>Allergy alert:</b> Aspirin may cause a severe allergic reaction which may include:</p> <ul style="list-style-type: none"> <li>hives</li> <li>facial swelling</li> <li>asthma (wheezing)</li> <li>shock</li> </ul> <p><b>Liver warning:</b> This product contains acetaminophen. Severe liver damage may occur if you take</p> <ul style="list-style-type: none"> <li>more than 2 caplets in 24 hours, which is the maximum daily amount</li> <li>with other drugs containing acetaminophen</li> <li>3 or more alcoholic drinks every day while using this product</li> </ul> <p><b>Stomach bleeding warning:</b> This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you</p> <ul style="list-style-type: none"> <li>are age 60 or older</li> <li>have had stomach ulcers or bleeding problems</li> <li>take a blood thinning (anticoagulant) or steroid drug</li> <li>take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)</li> <li>have 3 or more alcoholic drinks every day while using this product</li> <li>take more or for a longer time than directed</li> </ul>
--	---



**DO NOT USE IF EXCEDRIN SAFETY SEAL AROUND CAP IS BROKEN OR MISSING.**



3 0067266750 2

<p><b>Drug Facts (continued)</b></p> <p><b>Medication overuse headache warning:</b> Headaches may worsen if this product is used for 10 or more days per month.</p> <p><b>Caffeine warning:</b> The recommended dose of this product contains about as much caffeine as a cup of coffee. Limit the use of caffeine-containing medications, foods, or beverages while taking this product because too much caffeine may cause nervousness, irritability, sleeplessness, and, occasionally, rapid heart beat.</p> <p><b>Do not use</b></p> <ul style="list-style-type: none"> <li>if you have ever had an allergic reaction to acetaminophen, aspirin or any other pain reliever/fever reducer</li> <li>with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.</li> </ul> <p><b>Ask a doctor before use if</b></p> <ul style="list-style-type: none"> <li>you have never had migraines diagnosed by a health professional</li> <li>you have a headache that is different from your usual migraines</li> <li>you have the worst headache of your life</li> <li>you have fever and stiff neck</li> <li>you have headaches beginning after or caused by head injury, exertion, coughing or bending</li> <li>you experienced your first headache after the age of 50</li> <li>you have daily headaches</li> <li>you have a migraine so severe as to require bed rest</li> <li>you have liver disease</li> <li>stomach bleeding warning applies to you</li> <li>you have a history of stomach problems, such as heartburn</li> <li>you have high blood pressure, heart disease, liver cirrhosis, or kidney disease</li> <li>you are taking a diuretic</li> <li>you have asthma</li> <li>you have problems or serious side effects from taking pain relievers or fever reducers</li> <li>you have vomiting with your migraine headache</li> </ul> <p><b>Ask a doctor or pharmacist before use if you are</b></p> <ul style="list-style-type: none"> <li>taking a prescription drug for: <ul style="list-style-type: none"> <li>diabetes</li> <li>gout</li> <li>arthritis</li> </ul> </li> <li>under a doctor's care for any serious condition</li> <li>taking any other drug</li> <li>taking any other product that contains aspirin, acetaminophen, or any other pain reliever/fever reducer</li> </ul> <p><b>Stop use and ask a doctor if</b></p> <ul style="list-style-type: none"> <li>an allergic reaction occurs. Seek medical help right away.</li> <li>you experience any of the following signs of stomach bleeding: <ul style="list-style-type: none"> <li>feel faint</li> <li>vomit blood</li> <li>have bloody or black stools</li> </ul> </li> <li>have stomach pain that does not get better</li> </ul>	<p><b>Drug Facts (continued)</b></p> <ul style="list-style-type: none"> <li>your migraine is not relieved or worsens after first dose</li> <li>new or unexpected symptoms occur</li> <li>ringing in the ears or loss of hearing occurs</li> </ul> <p><b>If pregnant or breast-feeding,</b> ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.</p> <p><b>Keep out of reach of children.</b></p> <p>In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.</p> <p><b>Directions</b></p> <ul style="list-style-type: none"> <li>do not use more than directed</li> <li>adults: take 2 caplets with a glass of water</li> <li>if symptoms persist or worsen, ask your doctor</li> <li>do not take more than 2 caplets in 24 hours, unless directed by a doctor</li> <li>under 18 years of age: ask a doctor</li> </ul> <p><b>Other information</b></p> <ul style="list-style-type: none"> <li>store at or below 25°C (77°F)</li> <li>close cap tightly after use</li> <li>read all product information before using</li> </ul> <p><b>Inactive ingredients</b> benzoic acid, carnauba wax, FD&amp;C blue no. 1, hypromellose, light mineral oil, low-substituted hydroxypropyl cellulose, microcrystalline cellulose, polyorbate 20, povidone, propylene glycol, simethicone, sorbitan monolaurate, stearic acid, titanium dioxide</p> <p><b>Questions or comments?</b> 1-800-468-7746</p> <p>Distributed by: <b>GSK Consumer Healthcare</b> Warren, NJ 07059 ©2022 GSK or licensor. Trademarks owned or licensed by GSK. 1001792</p> 
--	---

<b>GSK Regulatory Spec Box</b>		<b>Verified Date: 08.03.22</b>
<b>Drug Facts Info</b>		
<b>Drug Facts</b>	Helvetica Neue LT Std Bold Cond Oblique	9 pt.
<b>Drug Facts (continued)</b>	Helvetica Neue LT Std Bold Cond Oblique / Cond	8 pt.
<b>Headings</b>	Helvetica Neue LT Std Bold Cond Oblique	8 pt.
<b>Subheadings</b>	Helvetica Neue LT Std Bold Cond	6 pt.
<b>Body text</b>	Helvetica Neue LT Std Bold Cond / Cond	6 pt.
<b>Bullets</b>	Helvetica Neue LT Std Roman	5 pt.
<b>Bullets on same lines: end of statement separated from bulleted statement by two ems</b>		Yes
<b>Spacing of the hair lines from edge of box-i.e. Minimum of 2 spaces either side of Drug Fact Box</b>		Yes
<b>Tracking:</b>	0	<b>Horizontal Scale:</b> 85%
<b>Leading:</b>	6,25 pt.	<b>Maximum Characters/Inch:</b> < 39
<b>Barlines:</b>	1,5 pt.	<b>Hairlines:</b> 0,5 pt.
<b>Primary Display Panel Info</b>		
<b>Size of Net Wt/Contents (Smallest character height in inches)*</b>		> 0.125 in.
<b>PDP dimensions (in square inches)</b>		5-25 sq. in.
<b>Ratio of Statement of Identity to Logo/Largest Copy on PDP</b>		
<b>Statement of Identity (pt.) (Divided by) Largest Logo Copy (pt.)</b>		
<b>Vertical PDP:</b> N/A		<b>Horizontal PDP:</b> 35%

EXCEDRIN SAFETY SEALED • EXCEDRIN SAFETY SEALED  
TO OPEN PUSH DOWN THEN TURN → TO OPEN PUSH DOWN THEN TURN →

**Drug Facts (continued)**

- you have fever and stiff neck
- you have headaches beginning after or caused by head injury, exertion, coughing or bending
- you experienced your first headache after the age of 50
- you have daily headaches
- you have a migraine so severe as to require bed rest
- you have liver disease
- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have vomiting with your migraine headache

**Ask a doctor or pharmacist before use if you are**

- taking a prescription drug for:
  - diabetes
  - gout
  - arthritis
- under a doctor's care for any serious condition
- taking any other drug
- taking any other product that contains aspirin, acetaminophen, or any other pain reliever/fever reducer

**Stop use and ask a doctor if**

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
  - feel faint
  - vomit blood
  - have bloody or black stools
  - have stomach pain that does not get better
- your migraine is not relieved or worsens after first dose
- new or unexpected symptoms occur
- ringing in the ears or loss of hearing occurs

**If pregnant or breast-feeding, ask a health professional before use.** It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions** • do not use more than directed

- adults: take 2 caplets with a glass of water
- if symptoms persist or worsen, ask your doctor
- do not take more than 2 caplets in 24 hours, unless directed by a doctor
- under 18 years of age: ask a doctor

**Other information**


- store at or below 25°C (77°F)
- close cap tightly after use
- read all product information before using

**Inactive ingredients** benzoic acid, carnauba wax, FD&C blue no. 1, hypromellose, light mineral oil, low-substituted hydroxypropyl cellulose, microcrystalline cellulose, poly sorbate 20, povidone, propylene glycol, simethicone, sorbitan mono laurate, stearic acid, titanium dioxide

**Questions or comments?** 1-800-468-7746

Distributed by GSK Consumer Healthcare, Warren, NJ 07059  
©2022 GSK. All rights reserved. © 2022 GSK. All rights reserved. All trademarks are owned by or licensed to the GSK group of companies.

**DO NOT USE IF EXCEDRIN SAFETY SEAL AROUND CAP IS BROKEN OR MISSING.**



3 0067266760 1

**EXCEDRIN**  
**MIGRAINE**  
Acetaminophen, Aspirin (NSAID) and Caffeine  
Pain Reliever/Pain Reliever Aid

**Drug Facts**

**Active ingredients (in each caplet)**

Acetaminophen 250 mg ..... Pain reliever

Aspirin 250 mg (NSAID) ..... Pain reliever

Caffeine 65 mg ..... Pain reliever aid

transdermal anti-inflammatory drug

PEEL RIGHT CORNER FOR FULL DRUG FACTS

**Drug Facts (continued)**

**Use** • treats migraine

**Warnings**

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

**Allergy alert:** Aspirin may cause a severe allergic reaction which may include:

- hives
- facial swelling
- asthma (wheezing)
- shock

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take

- more than 2 caplets in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

**Medication overdose headache warning:** Headaches may worsen if this product is used for 10 or more days per month.

**Caffeine warning:** The recommended dose of this product contains about as much caffeine as a cup of coffee. Limit the use of caffeine-containing medications, foods, or beverages while taking this product because too much caffeine may cause nervousness, irritability, sleeplessness, and, occasionally, rapid heart beat.

**Do not use**

- if you have ever had an allergic reaction to acetaminophen, aspirin or any other pain reliever/fever reducer
- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

**Ask a doctor before use if**

- you have never had migraines diagnosed by a health professional
- you have a headache that is different from your usual migraines
- you have the worst headache of your life

GSK Regulatory Spec Box		Verified Date: 08.02.22
<b>Drug Facts Info</b>		
<b>Drug Facts</b>	Helvetica Neue LT Std Bold Cond Oblique	9 pt.
<b>Drug Facts (continued)</b>	Helvetica Neue LT Std Bold Cond Oblique / Cond	8 pt.
<b>Headings</b>	Helvetica Neue LT Std Bold Cond Oblique	8 pt.
<b>Subheadings</b>	Helvetica Neue LT Std Bold Cond	6 pt.
<b>Body text</b>	Helvetica Neue LT Std Bold Cond / Cond	6 pt.
<b>Bullets</b>	Helvetica Neue LT Std Roman	5 pt.
<b>Bullets on same lines: end of statement separated from bulleted statement by two ems</b>		Yes
<b>Spacing of the hair lines from edge of box--i.e., Minimum of 2 spaces either side of Drug Fact Box</b>		Yes
<b>Tracking:</b>	0 pt.	Horizontal Scale: 93%
<b>Leading:</b>	6 pt.	Maximum Characters/Inch: < 39
<b>Barlines:</b>	1.5 pt.	Hairlines: 0.5 pt.
<b>Primary Display Panel Info</b>		
<b>Size of Net Wt/Contents (Smallest character height in inches)*</b>		>0.0625 in.
<b>PDP dimensions (in square inches)</b>		< 5 sq. in.
<b>Ratio of Statement of Identity to Logo/Largest Copy on PDP</b>		
<b>Statement of Identity (pt.) (Divided by) Largest Logo Copy (pt.)</b>		
<b>Vertical PDP:</b>	N/A	<b>Horizontal PDP:</b> 42%