

Artwork No. 6-775480, Chloraprep One-Step 1mL Drug Facts Insert

(b) (4)

4.75" width

Front

(b) (4)



Cat. No. 260480 / 260480NSB NDC 054365-400-06

## Chloraprep® One-Step

**2% w/v chlorhexidine gluconate (CHG) and 70% v/v isopropyl alcohol (IPA)  
Patient Preoperative Skin Preparation**

**Non-sterile Solution  
Applicator is sterile if package is intact**

**1 ml APPLICATOR**

 Not made with natural rubber latex Do Not ReuseProfessional Use Only  
External Use Only

<b>Drug Facts</b>	
<b>Active ingredients</b> Chlorhexidine gluconate 2% w/v..... Isopropyl alcohol 70% v/v.....	<b>Purposes</b> .....Antiseptic .....Antiseptic
<b>Use</b> for the preparation of the patient's skin prior to surgery. Helps to reduce bacteria that potentially can cause skin infection.	
<b>Warnings</b> For external use only. Flammable, keep away from fire or flame. ■ do not use with electrocautery procedures	
<b>Allergy Alert</b> This product may cause a severe allergic reaction. Symptoms may include: ■ wheezing/difficulty breathing ■ shock ■ facial swelling ■ hives ■ rash If an allergic reaction occurs, stop use and seek medical help right away.	
<b>Do not use</b> ■ on patients allergic to chlorhexidine gluconate or any other ingredient in this product ■ for lumbar puncture or in contact with the meninges ■ on open skin wounds or as a general skin cleanser	

3.8" length

## Chloraprep® One-Step

**2% w/v chlorhexidine gluconate (CHG) and 70% v/v isopropyl alcohol (IPA)**

<b>Drug Facts (continued)</b> <b>When using this product</b> keep out of eyes, ears, and mouth. May cause serious or permanent injury if permitted to enter and remain. If contact occurs, rinse with cold water right away and contact a doctor. <b>Stop use and ask a doctor</b> if irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition. <b>Keep out of reach of children.</b> If swallowed, get medical help or contact a Poison Control Center right away.
<b>Directions</b> ■ use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns. ■ maximal treatment area for one applicator is approximately 2.5 in. x 2.5 in. (42 cm <sup>2</sup> ) ■ remove applicator from package; do not touch sponge ■ hold the applicator with the sponge down. Pinch wings <u>only once</u> to activate the ampule and release the antiseptic. ■ wet the sponge by pressing and releasing the sponge against the treatment area until liquid is visible on the skin ■ completely wet the treatment area with antiseptic ■ <b>dry surgical sites</b> (e.g., abdomen or arm): use gentle repeated back-and-forth strokes for approximately 30 seconds. Allow the area to air dry for approximately 30 seconds. ■ <b>moist surgical sites</b> (e.g., inguinal fold): use gentle repeated back-and-forth strokes for approximately 2 minutes. Allow the area to air dry for approximately 1 minute. ■ do not blot or wipe away ■ discard the applicator after a single use along with any portion of the solution not required to cover the prep area. It is not necessary to use the entire amount available.
<b>Other information</b> ■ store between 15–30 °C (59–86 °F) ■ avoid freezing and excessive heat above 40 °C (104 °F)
<b>Inactive ingredient</b> USP purified water
<b>Questions?</b> ■ <a href="http://www.chloraprep.com">www.chloraprep.com</a> ■ call 1-800-523-0502 (M-F 8 a.m.–5 p.m. CST)

CareFusion, El Paso, TX 79912 • Artwork 6-775480 • EXXXXX • Made in the USA of US and imported parts


Back

This label may not be the latest approved by FDA.  
For current labeling information, please visit <https://www.fda.gov/drugsatfda>

Artwork No. 6-001480, ChloroPrep One-Step 1 mL (b) (4)


(b) (4)

(b) (4)



Exp. Date and Lot No.: placement - for  
manufacturing purpose only.

Artwork No. 6-824480, Chloraprep One-Step 1mL Intermediate Carton



Artwork No. 6-775403F, ChloroPrep 2% Frepp Applicator, Non-Sterile - Folded Drug Facts Insert

(b) (4)

(b) (4)

Dimensions: 7.5" x 2.5"

← 2.5" →

**ChloroPrep® One-Step**  
 2% w/v chlorhexidine gluconate (CHG)  
 and 70% v/v isopropyl alcohol (IPA)  
**Patient Preoperative Skin Preparation**  
 Non-sterile Solution  
 Applicator is sterile if package is intact  
 1.5 ml FREPP® APPLICATOR

Active ingredients	Purposes
Chlorhexidine gluconate 2% w/v	Antiseptic
Isopropyl alcohol 70% v/v	Antiseptic

**Use** For the preparation of the patient's skin prior to surgery. Helps to reduce bacteria that potentially can cause skin infection.

**Warnings**  
 For external use only  
**Flammable, keep away from fire or flame.**  
 ■ do not use with electrocautery procedures

**Allergy Alert**  
 This product may cause a severe allergic reaction. Symptoms may include:  
 ■ wheezing/difficulty breathing ■ shock  
 ■ facial swelling ■ hives ■ rash  
 If an allergic reaction occurs, stop use and seek medical help right away.

**Do not use**  
 ■ on patients allergic to chlorhexidine gluconate or any other ingredient in this product  
 ■ for lumbar puncture or in contact with the meninges  
 ■ on open skin wounds or as a general cleanser

**When using this product** keep out of eyes, ears, and mouth. May cause serious or permanent injury if permitted to enter and remain. If contact occurs, rinse with cold water right away and contact a doctor.

**Stop use and ask a doctor if** irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition.

**Keep out of reach of children.** If swallowed, get medical help or contact a Poison Control Center right away.

**Directions**  
 ■ use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns.  
 ■ maximal treatment area for one applicator is approximately 2.5 in. x 2.5 in. (42 cm<sup>2</sup>)  
 ■ remove applicator from package; do not touch sponge  
 ■ hold the applicator horizontally with the sponge down. Pinch wings **only once** to activate the ampule and release the antiseptic.  
 ■ wet the sponge by pressing and releasing the sponge against the treatment area until liquid is visible on the skin  
 ■ completely wet the treatment area with antiseptic  
 ■ **dry surgical sites** (e.g., abdomen or arm): use gentle repeated back-and-forth strokes for approximately 30 seconds. Allow the area to air dry for approximately 30 seconds.  
 ■ **moist surgical sites** (e.g., inguinal fold): use gentle repeated back-and-forth strokes for approximately 2 minutes. Allow the area to air dry for approximately 1 minute.  
 ■ do not blot or wipe away  
 ■ discard the applicator after a single use along with any portion of the solution not required to cover the prep area. It is not necessary to use the entire amount available.

**Other information** ■ store between 15–30 °C (59–86 °F)  
 ■ avoid freezing and excessive heat above 40 °C (104 °F)

**Inactive ingredient** USP purified water

**Questions?** ■ [www.chloroprep.com](http://www.chloroprep.com)  
 ■ call 1-800-523-0502 (M-F 8 a.m.–5 p.m. CST)

See other side for additional information


7.5"


Front

NDC 054365-400-02  
 Cat. No. 260299NSB / 260599NSB

**ChloroPrep® One-Step**  
 2% w/v chlorhexidine gluconate (CHG)  
 and 70% v/v isopropyl alcohol (IPA)  
**Patient Preoperative Skin Preparation**  
 Non-sterile Solution  
 Applicator is sterile if package is intact  
 1.5 ml FREPP® APPLICATOR

See Drug Facts for Full Labeling Information

 Do Not Reuse  
  
External Use Only

 Not made with natural rubber latex  
  
Professional Use Only

CareFusion, El Paso, TX 79912

Artwork 6-775403F · PXXXXXX  
 Made in the USA of US and imported parts

Back

(b) (4)

This label may not be the latest approved by FDA.  
For current labeling information, please visit <https://www.fda.gov/drugsatfda>

Artwork No. 6-000299, ChloroPrep One-Step Frepp

(b) (4)

(b) (4)

(b) (4)



Trim Area

This label may not be the latest approved by FDA.  
For current labeling information, please visit <https://www.fda.gov/drugsatfda>

Artwork No. 6-001299, ChloroPrep One-Step Frepp (b) (4) (No Lot/Exp. Date)

(b) (4)

(b) (4)

(b) (4)

Cat. No. 260299 NDC 054365-400-02  
**ChloroPrep® One-Step**  
2% w/v chlorhexidine gluconate (CHG)  
and 70% v/v isopropyl alcohol (IPA)  
Patient Preoperative Skin Preparation  
**Non-sterile Solution**  
Applicator is sterile if package is intact  
1.5 ml FREPP® APPLICATOR  
**WARNING. FLAMMABLE.**  
**KEEP AWAY FROM FIRE OR FLAME.**  
CareFusion, El Paso, TX 79912  
1-800-523-0502 (M-F 8 a.m. - 5 p.m. CST)  
Artwork No. 6-001299 · F090314  
Made in the USA of US and imported parts

Placeholder for lot and expiry data

Do Not Reuse

Bend and Peel


66 mm

40 mm

Exp. Date and Lot No.: placement - for  
manufacturing purpose only.

Reference ID: 4151410

Artwork No. 6-824299, ChloroPrep One-Step Frepp Intermediate Carton



Artwork No. 6-775400, Chloraprep One-Step and Chloraprep With Tint Drug Facts Insert for 3-mL  
Applicator Sterile/Non-Sterile/ (b) (4)




(b) (4)

(b) (4)

4.75" width

Front

(b) (4)

<b>WARNING</b> <b>FLAMMABLE</b>	Cat. No. 260415 / 260415NS / 260415NSB NDC 054365-400-11 Cat. No. 260400 / 260400NS / 260500NS NDC 054365-400-01
 <b>Keep away from fire or flame. To reduce risk of fire, PREP CAREFULLY:</b> <ul style="list-style-type: none"><li>solution contains alcohol and gives off <b>flammable vapors</b></li><li>avoid getting solution into hairy areas. Hair may take up to 1 hour to dry. <b>Wet hair is flammable.</b></li><li>do not drape or use ignition source (e.g., cautery, laser) until solution is completely dry (minimum of 3 minutes on hairless skin; up to 1 hour in hair)</li><li>do not allow solution to pool</li><li>remove wet materials from prep area</li></ul>	<b>Chloraprep® One-Step Chloraprep® With Tint</b> 2% w/v chlorhexidine gluconate (CHG) and 70% v/v isopropyl alcohol (IPA) <b>Patient Preoperative Skin Preparation</b> <b>Non-sterile Solution</b> <b>Applicator is sterile if package is intact</b> <b>3 ml APPLICATOR</b> Professional Use Only External Use Only  Do Not Reuse  Not made with natural rubber latex
<b>Drug Facts</b>	
<b>Active ingredients</b> Chlorhexidine gluconate 2% w/v..... Isopropyl alcohol 70% v/v.....	<b>Purposes</b> .....Antiseptic .....Antiseptic
<b>Use</b> for the preparation of the patient's skin prior to surgery. Helps to reduce bacteria that potentially can cause skin infection.	
<b>Warnings</b> <b>For external use only. Flammable, keep away from fire or flame. To reduce risk of fire, PREP CAREFULLY:</b> <ul style="list-style-type: none"><li>solution contains alcohol and gives off <b>flammable vapors</b></li><li>avoid getting solution into hairy areas. Hair may take up to 1 hour to dry. <b>Wet hair is flammable.</b></li><li>do not drape or use ignition source (e.g., cautery, laser) until solution is completely dry (minimum of 3 minutes on hairless skin; up to 1 hour in hair)</li><li>do not allow solution to pool</li><li>remove wet materials from prep area</li></ul>	
<b>Allergy Alert</b> This product may cause a severe allergic reaction. Symptoms may include: ■ wheezing/difficulty breathing ■ shock ■ facial swelling ■ hives ■ rash If an allergic reaction occurs, stop use and seek medical help right away.	

3.8" length

<b>Chloraprep® One-Step / Chloraprep® With Tint</b> 2% w/v chlorhexidine gluconate (CHG) and 70% v/v isopropyl alcohol (IPA)
<b>Drug Facts (continued)</b>
<b>Do not use</b> <ul style="list-style-type: none"><li>on patients allergic to chlorhexidine gluconate or any other ingredient in this product</li><li>for lumbar puncture or in contact with the meninges</li><li>on open skin wounds or as a general skin cleanser</li></ul>
<b>When using this product</b> keep out of eyes, ears, and mouth. May cause serious or permanent injury if permitted to enter and remain. If contact occurs, rinse with cold water right away and contact a doctor.
<b>Stop use and ask a doctor</b> if irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition.
<b>Keep out of reach of children.</b> If swallowed, get medical help or contact a Poison Control Center right away.
<b>Directions</b> <ul style="list-style-type: none"><li>use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns.</li><li>use in a well ventilated area</li><li>maximal treatment area for one applicator is approximately 4 in. x 5 in. (130 cm<sup>2</sup>)</li><li>remove applicator from package; do not touch sponge</li><li>hold the applicator with the sponge down. Pinch wings <b>only once</b> to activate the ampule and release the antiseptic.</li><li>wet the sponge by pressing and releasing the sponge against the treatment area until liquid is visible on the skin</li><li>completely wet the treatment area with antiseptic</li><li><b>dry surgical sites</b> (e.g., abdomen or arm): use gentle repeated back-and-forth strokes of the sponge for approximately 30 seconds. Allow solution to air dry for approximately 30 seconds. Do not blot or wipe away.</li><li><b>moist surgical sites</b> (e.g., inguinal fold): use gentle repeated back-and-forth strokes of the sponge for approximately 2 minutes. Allow solution to air dry for approximately 1 minute. Do not blot or wipe away.</li><li><b>if using an ignition source, allow the solution to completely dry</b> (minimum of 3 minutes on hairless skin; up to 1 hour in hair). Do not blot or wipe away.</li><li>discard the applicator after a single use along with any portion of the solution not required to cover the prep area. It is not necessary to use the entire amount available.</li></ul>
<b>Other information</b> ■ store between 15–30 °C (59–86 °F) ■ avoid freezing and excessive heat above 40 °C (104 °F) ■ the tint will slowly fade from the skin. Soap and water, or alcohol may be used to remove the tint if desired.
<b>Inactive ingredients</b> FD&C yellow #6 dye (Hi-Lite Orange applicators only), USP purified water
<b>Questions?</b> ■ <a href="http://www.chloraprep.com">www.chloraprep.com</a> ■ call 1-800-523-0502 (M-F 8 a.m.–5 p.m. CST)
CareFusion, El Paso, TX 79912 · Artwork 6-775400 · IXXXXX · Made in the USA of US and imported parts


Back

# Artwork No. 6-786415, ChloroPrep w/Tint (Orange) 3ml Applicator Lidding

(b) (4)




Lot/Exp Placeholder -  
for reference only.



# Artwork No.6-786315, ChloroPrep One-Step 3ml Applicator Lidding


(b) (4)



Lot/Exp Placeholder - for reference only.

Artwork No. 6-824400, ChloroPrep One-Step 3mL Intermediate Carton

1 15/32



ChloroPrep® One-Step  
2% w/v chlorhexidine gluconate (CHG)  
and 70% v/v isopropyl alcohol (IPA)  
Patient Preoperative Skin Preparation  
Non-sterile Solution  
Applicator is sterile if package is intact

**WARNING: FLAMMABLE. KEEP AWAY FROM FIRE OR FLAME.**

**Drug Facts**

**Active Ingredients**  
Chlorhexidine gluconate 2% w/v  
Isopropyl alcohol 70% v/v

**Purposes**  
Antiseptic  
Antiseptic

**Use** For the preparation of the patient's skin prior to surgery. Helps to reduce bacteria that potentially can cause skin infection.

**Warnings**  
For external use only. Flammable, keep away from fire or flame. To reduce the risk of fire, PREP CAREFULLY:  
• avoid getting solution into hairy areas. Hair may take up to 1 hour to dry. Wet hair is flammable.  
• do not drap or use ignition source (e.g., cautery, laser) until solution is completely dry (minimum of 3 minutes on hairless skin; up to 1 hour in hair)  
• do not allow solution to pool  
• remove wet materials from prep area

**Caution:** This product may cause a severe allergic reaction. Symptoms may include:  
• anaphylactic shock  
• facial swelling  
• hives  
• rash  
If an allergic reaction occurs, stop use and seek medical help right away.

**Do not use**  
• on patients allergic to chlorhexidine gluconate or any other ingredient in this product  
• for laceration or in contact with the mucous membranes  
• on open skin wounds or as a general skin cleanser

**When using this product** keep out of eyes, ears, and mouth. May cause serious or permanent injury if permitted to enter and remain. If contact occurs, rinse with cold water right away and contact a doctor.

**Stop use and ask a doctor** if irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition.  
Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

**Directions**  
• Use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns.  
• Use in a well-ventilated area.  
• maximal treatment area for one applicator is approximately 4 in. x 5 in. (100 cm)  
• remove applicator from package; do not touch sponge  
• hold the applicator with the sponge down. Pinch wings only to activate the ampule and release the antiseptic.  
• wet the sponge by pressing and releasing the sponge against the treatment area until liquid is visible on the skin  
• completely wet the treatment area with antiseptic  
• dry surgical sites (e.g., abdomen or arm) use gentle repeated back-and-forth strokes of the sponge for approximately 30 seconds. Allow solution to air dry for approximately 30 seconds. Do not blot or wipe area.  
• moist surgical sites (e.g., inguinal fold) use gentle repeated back-and-forth strokes of the sponge for approximately 2 minutes. Allow solution to air dry for approximately 1 minute. Do not blot or wipe away.  
• if using on ignition sources, allow the solution to completely dry (minimum of 3 minutes on hairless skin; up to 1 hour in hair). Do not blot or wipe away.  
• discard the applicator after a single use along with any portion of the solution not required to cover the prep area. It is not necessary to use the entire amount available.

**Other information**  
• avoid freezing and excessive heat above 40 °C (104 °F)  
**Tactile ingredient** USP purified water

**Questions?**  
• visit [www.chloroprep.com](http://www.chloroprep.com)  
• call 1-800-523-0502 (M-F 9 a.m. - 4 p.m. CST)

25 applicators

3 ml  
APPLICATORS  
Clear

ChloroPrep® One-Step  
2% w/v chlorhexidine gluconate (CHG)  
and 70% v/v isopropyl alcohol (IPA)  
Patient Preoperative Skin Preparation  
Non-sterile Solution  
Applicator is sterile if package is intact

3 ml  
APPLICATORS  
Clear

ChloroPrep® One-Step  
2% w/v chlorhexidine gluconate (CHG)  
and 70% v/v isopropyl alcohol (IPA)  
Patient Preoperative Skin Preparation  
Non-sterile Solution  
Applicator is sterile if package is intact

3 ml  
APPLICATORS  
Clear

25 applicators

3 ml  
APPLICATORS  
Clear

Do Not Reuse  
Not made with natural rubber latex  
STERILE

Professional Use Only  
Store between 15-30 °C (59-86 °F)  
Avoid freezing and excessive heat above 40 °C (104 °F)

25 applicators  
0.10 fl. oz. (3 mL) each

ChloroPrep® One-Step  
2% w/v chlorhexidine gluconate (CHG)  
and 70% v/v isopropyl alcohol (IPA)  
Patient Preoperative Skin Preparation  
Non-sterile Solution  
Applicator is sterile if package is intact

**WARNING  
FLAMMABLE**  
Keep away from fire or flame.  
To reduce risk of fire, PREP CAREFULLY:  
• solution contains alcohol and gives off flammable vapors  
• avoid getting solution into hairy areas.  
• hair may take up to 1 hour to dry. Wet hair is flammable.  
• do not drap or use ignition source (e.g., cautery, laser) until solution is completely dry (minimum of 3 minutes on hairless skin; up to 1 hour in hair)  
• do not allow solution to pool  
• remove wet materials from prep area

ChloroPrep® One-Step  
2% w/v chlorhexidine gluconate (CHG)  
and 70% v/v isopropyl alcohol (IPA)  
Patient Preoperative Skin Preparation  
Non-sterile Solution  
Applicator is sterile if package is intact

3 ml  
APPLICATORS  
Clear

LOT XXXXX    EXPIRY DATE XXXX  
No Variab

BARCODE PLACEHOLDER

©2009 Cardinal. All rights reserved. ChloroPrep is a registered trademark of Cardinal Corporation in one of its subsidiaries.

3 1/4

6 1/2

2 5 1/6 4

1 7 1/4

3 1 1/16

1 1/16

1 2 7/8

5 5/8

1 2 7/8


5 1 9/32

3 7 1 9/32

Artwork No. 6-824415, ChloroPrep with Tint 3ml Intermediate Carton

1 15/32

**ChloroPrep® With Tint**  
2% w/v chlorhexidine gluconate (CHG) and 70% v/v isopropyl alcohol (IPA)  
Patient Preoperative Skin Preparation  
Non-sterile Solution  
Applicator is sterile if package is intact



**WARNING, FLAMMABLE. KEEP AWAY FROM FIRE OR FLAME.**

Drug Facts	
<b>Active ingredients</b>	<b>Purposes</b>
Chlorhexidine gluconate 2% w/v	Antiseptic
Isopropyl alcohol 70% v/v	Antiseptic
<b>Uses</b> For the preparation of the patient's skin prior to surgery, helps to reduce bacteria that potentially can cause skin infection.	
<b>Warnings</b>	
For external use only. Flammable, keep away from fire or flame. To reduce the risk of fire, PREP CAREFULLY:	
<ul style="list-style-type: none"> <li>• solution contains alcohol and gives off flammable vapors.</li> <li>• avoid getting solution into hairy areas. Hair may take up to 1 hour to dry. Wet hair is flammable.</li> <li>• do not drag or use sponge around (e.g., cautery, heat) until solution is completely dry (minimum of 3 minutes on hairless skin; up to 1 hour in hair)</li> <li>• do not allow solution to pool</li> <li>• remove wet materials from prep area</li> </ul>	
<b>Allergy Alert:</b> This product may cause a severe allergic reaction. Symptoms may include:	
<ul style="list-style-type: none"> <li>• anaphylactic shock</li> <li>• hives</li> <li>• facial swelling</li> <li>• rash</li> </ul>	
If an allergic reaction occurs, stop use and seek medical help right away.	
<b>Do not use</b>	
<ul style="list-style-type: none"> <li>• if allergic to chlorhexidine gluconate or any other ingredient in this product</li> <li>• for lumbar puncture or in contact with the meninges</li> <li>• on open skin wounds or as a general skin cleanser</li> </ul>	
<b>When using this product</b> keep out of eyes, ears, and mouth. May cause serious or permanent injury if permitted to enter and remain. If contact occurs, rinse with cold water right away and contact a doctor.	
<b>Stop use and ask a doctor if</b> irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition.	
<b>Keep out of reach of children.</b> If swallowed, get medical help or contact a Poison Control Center right away.	
<b>Directions</b>	
<ul style="list-style-type: none"> <li>• use in a well-ventilated area</li> <li>• maximal treatment area for one applicator is approximately 4 in. x 5 in. (130 cm<sup>2</sup>)</li> <li>• remove applicator from package, do not touch sponge</li> <li>• hold the applicator with the sponge down. Pinch wings only once to activate the ampule and release the antiseptic.</li> <li>• wet the sponge by pinching and releasing the sponge against the treatment area until liquid is visible on the skin.</li> <li>• completely wet the treatment area with antiseptic.</li> <li>• dry surgical sites (e.g., abdomen or arm): use gentle repeated back-and-forth strokes of the sponge for approximately 30 seconds. Allow solution to air dry for approximately 30 seconds. Do not blot or wipe area.</li> <li>• moist surgical sites (e.g., inguinal fold): use gentle repeated back-and-forth strokes of the sponge for approximately 2 minutes. Allow solution to air dry for approximately 1 minute. Do not blot or wipe away.</li> <li>• if using on lighter-colored skin, allow the solution to completely dry (minimum of 3 minutes on hairless skin; up to 1 hour in hair). Do not blot or wipe away.</li> <li>• discard the applicator after a single use along with any portion of the solution not required to cover the prep area. It is not necessary to use the entire amount available.</li> </ul>	
<b>Other information</b>	
<ul style="list-style-type: none"> <li>• store between 15-30 °C (59-86 °F)</li> <li>• avoid freezing and excessive heat above 40 °C (104 °F)</li> </ul>	
<b>Inactive ingredients</b> FD&C yellow #6 dye, USP purified water	
<b>Questions?</b>	
<ul style="list-style-type: none"> <li>• <a href="http://www.chloroprep.com">www.chloroprep.com</a></li> <li>• call 1-800-823-9882 (M-F 9 a.m.-5 p.m. CST)</li> </ul>	

25 applicators


**3 ml**  
APPLICATORS  
Hi-Lite Orange®

**ChloroPrep® With Tint**  
2% w/v chlorhexidine gluconate (CHG) and 70% v/v isopropyl alcohol (IPA)

**3 ml**  
APPLICATORS  
Hi-Lite Orange®

**ChloroPrep® With Tint**  
2% w/v chlorhexidine gluconate (CHG) and 70% v/v isopropyl alcohol (IPA)  
Patient Preoperative Skin Preparation  
Non-sterile Solution  
Applicator is sterile if package is intact

**3 ml**  
APPLICATORS  
Hi-Lite Orange®



**3 ml**  
APPLICATORS  
Hi-Lite Orange®


Do Not Reuse  
Not made with natural rubber latex  
STERILE

Professional Use Only  
Store between 15-30 °C (59-86 °F)  
Avoid freezing and excessive heat above 40 °C (104 °F)

25 applicators  
0.10 fl. oz. (3 ml) each

25 applicators

CareFusion  
El Paso, TX 79912  
1.800.523.0502  
[www.chloroprep.com](http://www.chloroprep.com)  
Made in the USA of US and imported parts




**ChloroPrep® With Tint**  
2% w/v chlorhexidine gluconate (CHG) and 70% v/v isopropyl alcohol (IPA)  
Patient Preoperative Skin Preparation  
Non-sterile Solution  
Applicator is sterile if package is intact

NDC 054365-400-11  
Cat. No. 260415

**WARNING FLAMMABLE**  
Keep away from fire or flame. To reduce the risk of fire, PREP CAREFULLY:  
solution contains alcohol and gives off flammable vapors  
avoid getting solution into hairy areas.  
Hair may take up to 1 hour to dry. Wet hair is flammable.  
• do not drag or use sponge around (e.g., cautery, heat) until solution is completely dry (minimum of 3 minutes on hairless skin; up to 1 hour in hair)  
• do not allow solution to pool  
• remove wet materials from prep area

**ChloroPrep® With Tint**  
2% w/v chlorhexidine gluconate (CHG) and 70% v/v isopropyl alcohol (IPA)  
Patient Preoperative Skin Preparation  
Non-sterile Solution  
Applicator is sterile if package is intact

**3 ml**  
APPLICATORS  
Hi-Lite Orange®



LOT XXXXX    EXPIRY DATE XXXX  
No Variants

BARCODE PLACEHOLDER

©2008 CareFusion. All rights reserved. ChloroPrep® is a registered trademark of CareFusion Corporation or one of its subsidiaries.

No Variants / No printing area

3 1/4

6 1/2

2 5 1/6 4

1 7 1/4

3 1 1/16

1 1/16

1 2 7/8

5 5/8

1 2 7/8

5 1 9/32

3 7 1 9/32

Artwork No. 6-640248, 10.5-mL Applicator ChloraPrep® One-Step




(b) (4)

(b) (4)

4.75" width

Front

(b) (4)

<b>WARNING FLAMMABLE</b>		Cat. No. 260700 / 260700NS / 260700NSB Cat. No. 260715 / 260715NS / 260715NSB Cat. No. 260725 / 260725NS / 260725NSB	NDC 054365-400-04 NDC 054365-400-12 NDC 054365-400-09
 <b>Keep away from fire or flame. To reduce risk of fire, PREP CAREFULLY:</b>	<ul style="list-style-type: none"><li>• solution contains alcohol and gives off <b>flammable vapors</b></li><li>• avoid getting solution into hairy areas. Hair may take up to 1 hour to dry. <b>Wet hair is flammable.</b></li><li>• do not drape or use ignition source (e.g., cautery, laser) until solution is completely dry (minimum of 3 minutes on hairless skin; up to 1 hour in hair)</li><li>• do not allow solution to pool</li><li>• remove wet materials from prep area</li></ul>	<b>ChloraPrep® One-Step ChloraPrep® With Tint</b> 2% w/v chlorhexidine gluconate (CHG) and 70% v/v isopropyl alcohol (IPA) <b>Patient Preoperative Skin Preparation</b> <b>Non-sterile Solution</b> <b>Applicator is sterile if package is intact</b> <b>10.5 ml APPLICATOR</b>  Professional Use Only  Do Not Reuse Not made with natural rubber latex	
<b>Drug Facts</b>			
<b>Active ingredients</b> Chlorhexidine gluconate 2% w/v..... Isopropyl alcohol 70% v/v.....		<b>Purposes</b> .....Antiseptic .....Antiseptic	
<b>Use</b> for the preparation of the patient's skin prior to surgery. Helps to reduce bacteria that potentially can cause skin infection.			
<b>Warnings</b> <b>For external use only. Flammable, keep away from fire or flame. To reduce risk of fire, PREP CAREFULLY</b> ■ solution contains alcohol and gives off <b>flammable vapors</b> ■ avoid getting solution into hairy areas. Hair may take up to 1 hour to dry. <b>Wet hair is flammable.</b> ■ do not drape or use ignition source (e.g., cautery, laser) until solution is completely dry (minimum of 3 minutes on hairless skin; up to 1 hour in hair) ■ do not allow solution to pool ■ remove wet materials from prep area			
<b>Allergy Alert</b> This product may cause a severe allergic reaction. Symptoms may include: ■ wheezing/difficulty breathing ■ shock ■ facial swelling ■ hives ■ rash If an allergic reaction occurs, stop use and seek medical help right away.			

3.8" length

<b>ChloraPrep® One-Step / ChloraPrep® With Tint</b> 2% w/v chlorhexidine gluconate (CHG) and 70% v/v isopropyl alcohol (IPA)	
<b>Drug Facts (continued)</b>	
<b>Do not use</b> ■ on patients allergic to chlorhexidine gluconate or any other ingredient in this product ■ for lumbar puncture or in contact with the meninges ■ on open skin wounds or as a general skin cleanser	
<b>When using this product</b> keep out of eyes, ears, and mouth. May cause serious or permanent injury if permitted to enter and remain. If contact occurs, rinse with cold water right away and contact a doctor.	
<b>Stop use and ask a doctor if</b> irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition.	
<b>Keep out of reach of children.</b> If swallowed, get medical help or contact a Poison Control Center right away.	
<b>Directions</b> ■ use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns. ■ use in a well ventilated area ■ maximal treatment area for one applicator is approximately 8.4 in. x 8.4 in. (457 cm <sup>2</sup> ) ■ remove applicator from package; do not touch sponge ■ hold the applicator with the sponge down. Pinch wings <b>only once</b> to activate the ampule and release the antiseptic. ■ wet the sponge by pressing and releasing the sponge against the treatment area until liquid is visible on the skin ■ completely wet the treatment area with antiseptic ■ <b>dry surgical sites</b> (e.g., abdomen or arm): use gentle repeated back-and-forth strokes for 30 seconds ■ <b>moist surgical sites</b> (e.g., inguinal fold): use gentle repeated back-and-forth strokes for 2 minutes ■ do not allow solution to pool; tuck prep towels to absorb solution, and then remove ■ <b>allow the solution to completely dry</b> (minimum of 3 minutes on hairless skin; up to 1 hour in hair). Do not blot or wipe away. ■ discard the applicator after a single use along with any portion of the solution not required to cover the prep area. It is not necessary to use the entire amount available.	
<b>Other information</b> ■ store between 15–30 °C (59–86 °F) ■ avoid freezing and excessive heat above 40 °C (104 °F) ■ the tint will slowly fade from the skin. Soap and water, or alcohol may be used to remove the tint if desired.	
<b>Inactive ingredients</b> FD&C green #3 dye (Scrub-Teal applicators only), FD&C yellow #6 dye (Hi-Lite Orange applicators only), USP purified water	
<b>Questions?</b> ■ <a href="http://www.chloraprep.com">www.chloraprep.com</a> ■ call 1-800-523-0502 (M-F 8 a.m.–5 p.m. CST)	
CareFusion, El Paso, TX 79912 · Artwork No. 6-640248 · MXXXXXX · Made in the USA of US and imported parts	

Back

Artwork No. 6-786408 10.5-ml ChloroPrep Clear Applicator (b) (4) Labeling for K-8




(b) (4)



(b) (4)

Lot/Exp Placeholder - for reference only




(b) (4)

Artwork No. 6-786715, 10.5-mL ChloroPrep With Tint (Orange) Applicator


(b) (4) **Labeling**

(b) (4)



(b) (4)

Lot/Exp Placeholder - for reference only.




Artwork No. 6-786725 10.5-mL ChloroPrep With Tint Applicator (b) (4) Labeling

(b) (4)




(b) (4)

Lot/Exp Placeholder - for reference only.



(b) (4)


Part No. 6-773700  
10.5-ml ChloroPrep One-Step  
Line Mold Labels



(b) (4)

(b) (4)


Part No. 6-773715  
0.5-ml ChloroPrep With Tint  
In-Line Mold Labels



(b) (4)

(b) (4)

Part No. 6-773725  
10.5-ml ChloroPrep With Tint  
In-Line Mold Labels




(b) (4)

(b) (4)

Artwork No. # 6-824248, Intermediate Carton 10.5-mL Applicator ChloroPrep 2% One-Step

1<sup>15</sup>/<sub>32</sub>

ChloroPrep® One-Step  
2% w/v chlorhexidine gluconate (CHG)  
and 70% v/v isopropyl alcohol (IPA)  
Patient Preoperative Skin Preparation  
Non-sterile Solution  
Applicator is sterile if package is intact



25 applicators

10.5 ml  
APPLICATORS  
Clear

WARNING, FLAMMABLE, KEEP AWAY FROM FIRE OR FLAME.


Drug Facts	Purpose
<b>Active Ingredients:</b> Chlorhexidine gluconate 2% w/v Isopropyl alcohol 70% v/v	Antiseptic Antiseptic
<b>Use:</b> For the preparation of the patient's skin prior to surgery. Helps to reduce bacteria that potentially can cause skin infection.	
<b>Warnings:</b> For external use only. Flammable, keep away from fire or flame. To reduce the risk of fire, PREP CAREFULLY: <ul style="list-style-type: none"> <li>avoid getting solution into hairy areas. Hair may take up to 1 hour to dry. Wet hair is flammable.</li> <li>do not close or use goggles, or use eye shields, lenses, or contact lenses completely dry (minimum of 3 minutes on hairless skin; up to 1 hour in hair)</li> <li>do not use solution to wash</li> <li>do not use solution to peel</li> <li>remove wet materials from prep area.</li> </ul> <b>Caution:</b> This product may cause a severe allergic reaction. Symptoms may include: <ul style="list-style-type: none"> <li>anxiety or difficulty breathing</li> <li>swelling</li> <li>itching</li> <li>skin rash</li> <li>if an allergic reaction occurs, stop use and seek medical help right away.</li> </ul> <b>Do not use:</b> <ul style="list-style-type: none"> <li>if you are allergic to chlorhexidine gluconate or any other ingredients in this product</li> <li>for lumbar puncture or in contact with the meninges</li> <li>on open skin wounds or for a general skin cleanser</li> </ul> <b>When using this product:</b> keep out of eyes, ears, and mouth. May cause serious or permanent injury if permitted to enter and remain. If contact occurs, rinse with cold water right away and contact a doctor. <b>Stop use and ask a doctor if:</b> irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition. <b>Keep out of reach of children.</b> If swallowed, get medical help or contact a Poison Control Center right away.	
<b>Directions:</b> <ul style="list-style-type: none"> <li>use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns.</li> <li>use in a well-ventilated area.</li> <li>remove applicator from package; do not touch sponge.</li> <li>hold the applicator with the sponge down. Press sponge onto skin to activate the applicator and release the antiseptic.</li> <li>use the sponge for cleansing and rubbing. Do not scrub the treatment area with liquid or residue on the skin.</li> <li>avoid rubbing the treatment area with clothing.</li> <li>dry surgical sites (e.g., abdomen or arms) use gentle repeated back-and-forth strokes for 30 seconds.</li> <li>avoid surgical sites (e.g., inguinal folds) use gentle repeated back-and-forth strokes for 2 minutes.</li> <li>do not allow solution to pool; back prep towels to absorb solution, and then remove.</li> <li>allow the solution to completely dry (minimum of 3 minutes on hairless skin; up to 1 hour in hair). Do not blot or wipe away.</li> <li>discard the applicator after a single use along with any portion of the solution not required to cover the prep area. It is not necessary to use the entire amount available.</li> </ul>	
<b>Other information:</b> <ul style="list-style-type: none"> <li>store between 15-30 °C (59-86 °F)</li> <li>avoid freezing and excessive heat above 40 °C (104 °F)</li> </ul>	
<b>Inactive ingredient:</b> Sorbic acid	
<b>Questions?</b> <ul style="list-style-type: none"> <li>visit <a href="http://www.chloroprep.com">www.chloroprep.com</a></li> <li>call 1-800-523-0502 (M-F 8 a.m.-5 p.m. CST)</li> </ul>	

ChloroPrep® One-Step  
2% w/v chlorhexidine gluconate (CHG)  
and 70% v/v isopropyl alcohol (IPA)  
Patient Preoperative Skin Preparation  
Non-sterile Solution  
Applicator is sterile if package is intact

10.5 ml  
APPLICATORS  
Clear

ChloroPrep® One-Step  
2% w/v chlorhexidine gluconate (CHG)  
and 70% v/v isopropyl alcohol (IPA)  
Patient Preoperative Skin Preparation  
Non-sterile Solution  
Applicator is sterile if package is intact

10.5 ml  
APPLICATORS  
Clear



25 applicators

CareFusion  
P.O. Box 128, 79912  
1-800-523-0502  
www.chloroprep.com  
Made in the USA of US and imported parts


10.5 ml  
APPLICATORS  
Clear



Professional Use Only

Store between 15-30 °C (59-86 °F)  
Avoid freezing and excessive heat  
above 40 °C (104 °F)

25 applicators  
0.36 fl. oz. (10.5 ml) each



ChloroPrep® One-Step  
2% w/v chlorhexidine gluconate (CHG)  
and 70% v/v isopropyl alcohol (IPA)  
Patient Preoperative Skin Preparation  
Non-sterile Solution  
Applicator is sterile if package is intact

NDC 054365-400-04  
Cat. No. 260700



ChloroPrep® One-Step  
2% w/v chlorhexidine gluconate (CHG)  
and 70% v/v isopropyl alcohol (IPA)  
Patient Preoperative Skin Preparation  
Non-sterile Solution  
Applicator is sterile if package is intact

10.5 ml  
APPLICATORS  
Clear



©2020 CareFusion. All rights reserved. ChloroPrep® is a registered trademark of CareFusion Corporation or one of its subsidiaries.

No Variable / No printing area

3<sup>1</sup>/<sub>4</sub>

1 7<sup>1</sup>/<sub>4</sub>

6<sup>1</sup>/<sub>2</sub>

2<sup>5</sup>/<sub>16</sub>

3<sup>1</sup>/<sub>16</sub>

1<sup>1</sup>/<sub>16</sub>

1 2<sup>7</sup>/<sub>8</sub>

5<sup>5</sup>/<sub>8</sub>

1 2<sup>7</sup>/<sub>8</sub>

5<sup>1</sup>/<sub>32</sub>

3 7<sup>9</sup>/<sub>32</sub>

Artwork No. 6-824715, 10.5-mL ChloraPrep with Tint Applicator Intermediate Carton

1 15/32



ChloraPrep® With Tint  
2% w/v chlorhexidine gluconate (CHG)  
and 70% v/v isopropyl alcohol (IPA)  
Patient Preoperative Skin Preparation  
Non-sterile Solution  
Applicator is sterile if package is intact

**WARNING: FLAMMABLE. KEEP AWAY FROM FIRE OR FLAME.**

Drug Facts	Purposes
<p><b>Active ingredients</b> Chlorhexidine gluconate (CHG) 2% w/v Isopropyl alcohol (IPA) 70% v/v</p> <p><b>Use</b> Use for the preparation of the patient's skin prior to surgery. Helps to reduce bacteria that potentially can cause skin infection.</p> <p><b>Warnings</b> For external use only. Flammable, keep away from fire or flame. To reduce the risk of fire, PREP CAREFULLY.  <ul style="list-style-type: none"> <li>• solution contains alcohol and gives off flammable vapors</li> <li>• avoid getting solution into hairy areas. Hair may take up to 1 hour to dry. Wet hair is flammable.</li> <li>• do not drink or use solution around face, especially, infant's face; solution is completely dry (minimum of 3 minutes on hairless skin up to 1 hour in hair)</li> <li>• do not allow solution to pool</li> <li>• remove wet materials from prep area</li> </ul> </p> <p><b>Caution</b> Allergic Alert: This product may cause a severe allergic reaction. Symptoms may include:  <ul style="list-style-type: none"> <li>• wheezing/rapidly breathing</li> <li>• dizziness</li> <li>• facial swelling</li> <li>• hives</li> <li>• rash</li> </ul>                     If an allergic reaction occurs, stop use and seek medical help right away.</p> <p><b>Do not use</b>  <ul style="list-style-type: none"> <li>• if patient allergic to chlorhexidine gluconate or any other ingredient in this product</li> <li>• to lubricate puncture or to cleanse with the manometer</li> <li>• on open skin wounds or on a genital skin condition</li> </ul>                     When using this product, keep out of eyes, ears, and mouth. May cause serious or permanent injury if permitted to enter and remain. If contact occurs, rinse with cold water right away and contact a doctor.</p> <p><b>Stop use and ask a doctor if</b> irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition.</p> <p><b>Keep out of reach of children.</b> If swallowed, get medical help or contact a Poison Control Center right away.</p> <p><b>Directions</b>  <ul style="list-style-type: none"> <li>• use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns.</li> <li>• use in small treatment area</li> <li>• minimal treatment area for one applicator is approximately 8.4 in. x 8.4 in. (457 cm<sup>2</sup>)</li> <li>• remove applicator from package, do not touch applicator</li> <li>• hold the applicator with the sponge down. Pinch wings only, close to activate the ampule and release the antiseptic.</li> <li>• wet the sponge by squeezing and releasing the sponge against the treatment area until liquid is visible on the skin</li> <li>• completely wet the treatment area with antiseptic</li> <li>• use applicator after 10-15 seconds or until use gentle repeated back-and-forth strokes for 30 seconds</li> <li>• moisten and allow to dry (e.g., inguinal fold) use gentle repeated back-and-forth strokes for 2 minutes</li> <li>• do not allow solution to pool, and dry hands to absorb solution, and then remove</li> <li>• allow the solution to completely dry (minimum of 2 minutes on hairless skin, up to 1 hour in hair). Do not touch or wipe prep area.</li> <li>• discard the applicator after a single use along with any portion of the solution not required to cover the prep area. It is not necessary to use the entire amount available.</li> </ul> </p> <p><b>Other information</b>  <ul style="list-style-type: none"> <li>• store between 15-30 °C (59-86 °F)</li> <li>• avoid freezing and excessive heat above 40 °C (104 °F)</li> </ul> </p> <p><b>Inactive ingredients</b>                      Citric acid, 100% USP purified water</p> <p><b>Questions?</b>  <ul style="list-style-type: none"> <li>• visit <a href="http://www.chloraprep.com">www.chloraprep.com</a></li> <li>• call 1-800-523-6982 (M-F 8 a.m.-5 p.m., CST)</li> </ul> </p>	<p>ChloraPrep® With Tint 2% w/v chlorhexidine gluconate (CHG) and 70% v/v isopropyl alcohol (IPA) Patient Preoperative Skin Preparation Non-sterile Solution Applicator is sterile if package is intact</p> <p>10.5 ml APPLICATORS Hi-Lite Orange®</p> <p>10.5 ml APPLICATORS Hi-Lite Orange®</p> <p>10.5 ml APPLICATORS Hi-Lite Orange®</p> <p>10.5 ml APPLICATORS Hi-Lite Orange®</p> <p>25 applicators</p> <p>25 applicators 0.36 fl. oz. (10.5 ml) each</p> <p>Not made with natural rubber latex</p> <p>STERILE [EO]</p> <p>Professional Use Only</p> <p>Store between 15-30 °C (59-86 °F) Avoid freezing and excessive heat above 40 °C (104 °F)</p> <p>25 applicators 0.36 fl. oz. (10.5 ml) each</p> <p>NDC 054365-400-12 Cat. No. 260715</p> <p>ChloraPrep® With Tint 2% w/v chlorhexidine gluconate (CHG) and 70% v/v isopropyl alcohol (IPA) Patient Preoperative Skin Preparation Non-sterile Solution Applicator is sterile if package is intact</p> <p>10.5 ml APPLICATORS Hi-Lite Orange®</p> <p>ChloraPrep® With Tint 2% w/v chlorhexidine gluconate (CHG) and 70% v/v isopropyl alcohol (IPA) Patient Preoperative Skin Preparation Non-sterile Solution Applicator is sterile if package is intact</p> <p>10.5 ml APPLICATORS Hi-Lite Orange®</p> <p>LOT XXXXXX (printed in blue) No Variish</p> <p>EXP DATE XXXX</p> <p>BARCODE PLACEHOLDER</p> <p>©2009 CareFusion. All rights reserved. ChloraPrep® is a registered trademark of CareFusion Corporation or one of its subsidiaries.</p> <p>No Variish / No printing area</p>

3 1 1/16

1 1/16

1 2 7/8

5 5/8

1 2 7/8

5 1 9/32

3 7 9/32

3 1/4

6 1/2


2 5 1/6 4

1 7 1/4

Artwork No. 6-824725, 10.5-mL ChloroPrep with Tint Intermediate Carton

1<sup>15</sup>/<sub>32</sub>

3<sup>1</sup>/<sub>16</sub>



ChloroPrep® With Tint  
2% w/v chlorhexidine gluconate (CHG)  
and 70% v/v isopropyl alcohol (IPA)  
Patient Preoperative Skin Preparation  
Non-sterile Solution  
Applicator is sterile if package is intact

**WARNING: FLAMMABLE. KEEP AWAY FROM FIRE OR FLAME.**

**Drug Facts**

**Active ingredients**  
Chlorhexidine gluconate 2% w/v  
Isopropyl alcohol 70% v/v

**Purposes**  
Antiseptic  
Anesthetic

**Use**  
For the preparation of the patient's skin prior to surgery. Helps to reduce bacteria that potentially can cause skin infection.

**Warnings**  
For external use only. Flammable, keep away from fire or flame. To reduce the risk of fire, PREP CAREFULLY:  
• solution contains alcohol and gives off flammable vapors.  
• avoid getting solution into hairy areas. Hair may take up to 1 hour to dry. Wet hair is flammable.  
• do not drop or use ignition source (e.g., candle, laser) until solution is completely dry (minimum of 3 minutes on hairless skin; up to 1 hour in hair).  
• do not allow solution to dry on skin.  
• remove wet materials from prep area.

**Allergy Alert:** This product may cause a severe allergic reaction. Symptoms may include:  
• hives  
• difficulty breathing  
• chest swelling  
• rash  
• dizziness  
• an allergic reaction occurs, stop use and seek medical help right away.

**Do not use**  
• on patients allergic to chlorhexidine gluconate or any other ingredient in this product.  
• on facial punctures or in contact with the meninges.  
• on open skin wounds or as a general skin cleanser.

**When using this product:** keep out of eyes, ears, and mouth. May cause serious or permanent injury if permitted to enter and remain. If contact occurs, rinse with cool water right away and contact a doctor.

**Keep out of reach of children.** If swallowed, sensitization, or allergic reaction occurs. These may be signs of a serious condition. Stop use and seek a doctor if irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition. Stop use and seek a doctor if irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition. Stop use and seek a doctor if irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition.

**Directions**  
• use only on non-bleeding, intact or intact under 2 months of age. These products may cause irritation or chemical burns.  
• use in a well-ventilated area.  
• maximal treatment area for one applicator is approximately 8.4 in. x 8.4 in. (457 cm²).  
• remove applicator from package. Do not touch applicator.  
• hold the applicator with the orange foam. Place applicator against the treatment area until liquid is visible on the skin.  
• use the applicator to prepare the treatment area. Rubbing the applicator against the treatment area until liquid is visible on the skin.  
• consistently rub the treatment area with applicator until the treatment area is completely dry (minimum of 3 minutes on hairless skin; up to 1 hour in hair).  
• dry the applicator (e.g., blotting or air-dry) and gently separate back-and-forth strokes to dry applicator.  
• moist surgical sites (e.g., surgical field): use gentle repeated back-and-forth strokes for 2 minutes.  
• do not allow solution to dry back prep towels in direct solution, and then remove.  
• allow the solution to completely dry (minimum of 3 minutes on hairless skin; up to 1 hour in hair). Do not blot or wipe away.  
• discard the applicator after a single use along with any portion of the solution not required to cover the prep area. It is not necessary to use the entire amount available.

**Other information**  
• store between 15-30 °C (59-86 °F).  
• avoid freezing and excessive heat above 40 °C (104 °F).

**Inactive ingredients**  
Ethanol 10% v/v, USP purified water.

**Questions?**  
• [www.chloroprep.com](http://www.chloroprep.com)  
• call 1-800-523-0502 (M-F 8 a.m.-5 p.m. CST).

25 applicators  
Scrub Teal®  
10.5 ml  
APPLICATORS

ChloroPrep® With Tint  
2% w/v chlorhexidine gluconate (CHG)  
and 70% v/v isopropyl alcohol (IPA)  
Patient Preoperative Skin Preparation  
Non-sterile Solution  
Applicator is sterile if package is intact

10.5 ml  
APPLICATORS  
Scrub Teal®

ChloroPrep® With Tint  
2% w/v chlorhexidine gluconate (CHG)  
and 70% v/v isopropyl alcohol (IPA)  
Patient Preoperative Skin Preparation  
Non-sterile Solution  
Applicator is sterile if package is intact

10.5 ml  
APPLICATORS  
Scrub Teal®

CareFusion  
El Paso, TX 79912  
1-800-523-0502  
[www.chloroprep.com](http://www.chloroprep.com)  
Made in the USA of US and imported parts

10.5 ml  
APPLICATORS  
Scrub Teal®

Not made with natural rubber latex  
STERILE EO

Professional Use Only

Store between 15-30 °C (59-86 °F)  
Avoid freezing and excessive heat above 40 °C (104 °F)

25 applicators  
0.36 fl. oz. (10.5 ml) each

NDC 054365-400-09  
Cat. No. 260725

ChloroPrep® With Tint  
2% w/v chlorhexidine gluconate (CHG)  
and 70% v/v isopropyl alcohol (IPA)  
Patient Preoperative Skin Preparation  
Non-sterile Solution  
Applicator is sterile if package is intact

**WARNING FLAMMABLE**  
Keep away from fire or flame.  
To reduce risk of fire, PREP CAREFULLY:  
• solution contains alcohol and gives off flammable vapors  
• avoid getting solution into hairy areas.  
• hair may take up to 1 hour to dry. Wet hair is flammable.  
• do not drop or use ignition source (e.g., candle, laser) until solution is completely dry (minimum of 3 minutes on hairless skin; up to 1 hour in hair).  
• do not allow solution to dry.  
• remove wet materials from prep area.

ChloroPrep® With Tint  
2% w/v chlorhexidine gluconate (CHG)  
and 70% v/v isopropyl alcohol (IPA)  
Patient Preoperative Skin Preparation  
Non-sterile Solution  
Applicator is sterile if package is intact

10.5 ml  
APPLICATORS  
Scrub Teal®

LOT XXXXX (printed here) EXP DATE XXXX  
No Varnish

BARCODE PLACEHOLDER

©2009 CareFusion. All rights reserved. ChloroPrep® is a registered trademark of CareFusion Corporation or one of its subsidiaries.

11/16

1 2<sup>7</sup>/<sub>8</sub>

5<sup>5</sup>/<sub>8</sub>

1 2<sup>7</sup>/<sub>8</sub>

5<sup>1</sup>/<sub>32</sub>

3 7<sup>1</sup>/<sub>32</sub>

1<sup>15</sup>/<sub>32</sub>

3<sup>1</sup>/<sub>4</sub>

6<sup>1</sup>/<sub>2</sub>

2<sup>5</sup>/<sub>64</sub>

1 7<sup>1</sup>/<sub>4</sub>

Artwork No. 6-775815, 26-mL ChloroPrep® One-Step & With Tint Applicator Sterile, Non-Sterile / (b) (4) Insert Sheet - Chlorhexidine Gluconate 2% w/v and Isopropyl Alcohol 70% v/v Patient Preoperative Skin Preparation (b) (4)



5" width Front

**⚠ WARNING**

**FLAMMABLE**

**Keep away from fire or flame.**  
**To reduce risk of fire, PREP CAREFULLY:**

- do not use 26-ml applicator for head and neck surgery or on an area smaller than 8.4 in. x 8.4 in. Use a smaller applicator instead.
- solution contains alcohol and gives off **flammable vapors**
- avoid getting solution into hairy areas. Hair may take up to 1 hour to dry. **Wet hair is flammable.**
- do not drape or use ignition source (e.g., cautery, laser) until solution is completely dry (minimum of 3 minutes on hairless skin; up to 1 hour in hair)
- do not allow solution to pool
- remove wet materials from prep area

Cat. No. 260800 / 260800NS / 260800NSB NDC 054365-400-14  
 Cat. No. 260815 / 260815NS / 260815NSB NDC 054365-400-13  
 Cat. No. 260825 / 260825NS / 260825NSB NDC 054365-400-05

ChloroPrep® One-Step  
ChloroPrep® With Tint

2% w/v chlorhexidine gluconate (CHG)  
and 70% v/v isopropyl alcohol (IPA)

Patient Preoperative Skin Preparation

Non-sterile Solution  
Applicator is sterile if package is intact

26 ml APPLICATOR

Professional Use Only Do Not Reuse Not made with natural rubber latex

Drug Facts

<b>Active ingredients</b>	<b>Purposes</b>
Chlorhexidine gluconate 2% w/v.....	Antiseptic
Isopropyl alcohol 70% v/v.....	Antiseptic

**Use** for the preparation of the patient's skin prior to surgery. Helps to reduce bacteria that potentially can cause skin infection.

**Warnings**

**For external use only. Flammable, keep away from fire or flame. To reduce risk of fire, PREP CAREFULLY:**

- do not use 26-ml applicator for head and neck surgery
- do not use on an area smaller than 8.4 in. x 8.4 in. Use a smaller applicator instead
- solution contains alcohol and gives off **flammable vapors**
- avoid getting solution into hairy areas. Hair may take up to 1 hour to dry. **Wet hair is flammable.**
- do not drape or use ignition source (e.g., cautery, laser) until solution is completely dry (minimum of 3 minutes on hairless skin; up to 1 hour in hair)
- do not allow solution to pool
- remove wet materials from prep area

4" length

ChloroPrep® One-Step / ChloroPrep® With Tint

2% w/v chlorhexidine gluconate (CHG) and 70% v/v isopropyl alcohol (IPA)

**Drug Facts** (continued)

**Allergy Alert**  
This product may cause a severe allergic reaction. Symptoms may include:  
 ■ wheezing/difficulty breathing ■ shock ■ facial swelling ■ hives ■ rash  
 If an allergic reaction occurs, stop use and seek medical help right away.

**Do not use**

- on patients allergic to chlorhexidine gluconate or any other ingredient in this product
- for lumbar puncture or in contact with the meninges
- on open skin wounds or as a general skin cleanser

**When using this product** keep out of eyes, ears, and mouth. May cause serious or permanent injury if permitted to enter and remain. If contact occurs, rinse with cold water right away and contact a doctor.

**Stop use and ask a doctor** if irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition.

**Keep out of reach of children.** If swallowed, get medical help or contact a Poison Control Center right away.

**Directions**

- use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns.
- use in a well ventilated area
- maximal treatment area for one applicator is approximately 13.2 in. x 13.2 in. (1126 cm<sup>2</sup>)
- remove applicator from package; do not touch sponge
- hold the applicator with the sponge down. Pinch wings **only once** to activate the ampules and release the antiseptic.
- wet the sponge by pressing and releasing the sponge against the treatment area until liquid is visible on the skin
- completely wet the treatment area with antiseptic
- **dry surgical sites** (e.g., abdomen or arm): use gentle repeated back-and-forth strokes for 30 seconds
- **moist surgical sites** (e.g., inguinal fold): use gentle repeated back-and-forth strokes for 2 minutes
- do not allow solution to pool; tuck prep towels to absorb solution, and then remove
- **allow the solution to completely dry** (minimum of 3 minutes on hairless skin; up to 1 hour in hair). Do not blot or wipe away.
- discard the applicator after a single use along with any portion of the solution not required to cover the prep area. It is not necessary to use the entire amount available.

**Other information** ■ store between 15–30 °C (59–86 °F) ■ avoid freezing and excessive heat above 40 °C (104 °F) ■ the tint will slowly fade from the skin. Soap and water, or alcohol may be used to remove the tint if desired.

**Inactive ingredients** FD&C green #3 dye (Scrub-Teal applicators only), FD&C yellow #6 dye (Hi-Lite Orange applicators only), USP purified water

**Questions?** ■ [www.chloroprep.com](http://www.chloroprep.com) ■ call 1-800-523-0502 (M-F 8 a.m.–5 p.m. CST)

CareFusion, El Paso, TX 79912 · Artwork No. 6-775815 · LXXXXXX · Made in the USA of US and imported parts

Back

Artwork No. 6-787800, 26-mL ChloroPrep One-Step Applicator Lidding

(b) (4)

(b) (4)

Lot # and Exp. Date are displayed at this end of the package.

Barcode Placeholder

NDC 054365-400-14  
Cat. No. 260800

Single Use  
**ChloroPrep® One-Step**  
2% w/v chlorhexidine gluconate (CHG)  
and 70% v/v isopropyl alcohol (IPA)  
Patient Preoperative Skin Preparation  
Non-sterile Solution  
Applicator is sterile if package is intact  
**26 ml APPLICATOR**

Clear

WARNING. FLAMMABLE. KEEP AWAY FROM FIRE OR FLAME.

**Drug Facts**

Active ingredients	Purposes
Chlorhexidine gluconate 2% w/v	Antiseptic
Isopropyl alcohol 70% v/v	Antiseptic

**Use** for the preparation of the patient's skin prior to surgery. Helps to reduce bacteria that potentially can cause skin infection.

**Warnings**

**For external use only. Flammable, keep away from fire or flame. To reduce risk of fire, PREP CAREFULLY:**

- do not use 26-ml applicator for head and neck surgery
- do not use on an area smaller than 8.4 in. x 8.4 in. Use a smaller applicator instead.
- solution contains alcohol and gives off flammable vapors
- avoid getting solution into hairy areas. Hair may take up to 1 hour to dry. **Wet hair is flammable.**
- do not drape or use ignition source (e.g., cautery, laser) until solution is completely dry (minimum of 3 minutes on hairless skin; up to 1 hour in hair)
- do not allow solution to pool
- remove wet materials from prep area

**Allergy Alert:** This product may cause a severe allergic reaction. Symptoms may include:

- wheezing/difficulty breathing
- shock
- facial swelling
- hives
- rash

If an allergic reaction occurs, stop use and seek medical help right away.

**Do not use**

- on patients allergic to chlorhexidine gluconate or any other ingredient in this product
- for lumbar puncture or in contact with the meninges
- on open skin wounds or as a general skin cleanser

**When using this product** keep out of eyes, ears, and mouth. May cause serious or permanent injury if permitted to enter and remain. If contact occurs, rinse with cold water right away and contact a doctor.

**Stop use and ask a doctor if** irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition.

**Keep out of reach of children.** If swallowed, get medical help or contact a Poison Control Center right away.

**Directions**

- use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns.
- use in a well ventilated area
- maximal treatment area for one applicator is approximately 13.2 in. x 13.2 in. (1126 cm<sup>2</sup>). Do not use 26-ml applicator for area smaller than 8.4 in. x 8.4 in. Use a smaller applicator instead.
- do not use 26-ml applicator for head and neck surgery
- remove applicator from package; do not touch sponge
- hold the applicator with the sponge down. Pinch wing only once to activate the ampules and release the antiseptic.
- wet the sponge by pressing and releasing the sponge against the treatment area until liquid is visible on the skin
- completely wet the treatment area with antiseptic
- dry surgical sites** (e.g., abdomen or arm): use gentle repeated back-and-forth strokes for 30 seconds
- moist surgical sites** (e.g., inguinal fold): use gentle repeated back-and-forth strokes for 2 minutes
- do not allow solution to pool; tuck prep towels to absorb solution, and then remove
- allow the solution to completely dry** (minimum of 3 minutes on hairless skin; up to 1 hour in hair). Do not blot or wipe away.
- discard the applicator after a single use along with any portion of the solution not required to cover the prep area. It is not necessary to use the entire amount available.

**Other information**

- store between 15–30 °C (59–86 °F)
- avoid freezing and excessive heat above 40 °C (104 °F)

**Inactive ingredient**

USP purified water

**Questions?**

- www.chloroprep.com
- call 1-800-523-0502 (M-F 8 a.m.–5 p.m. CST)

CareFusion, El Paso, TX 79912 • Artwork No. 6-787800 • MXXXXXX  
Made in the USA of US and imported parts

243

305

8.25

1


60

80.5

Lot/Exp Placeholder -  
for reference only

Artwork No. 6-786815, 26-mL ChloroPrep with Tint (Orange) Applicator Lidding

(b) (4)



(b) (4)

Lot # and Exp. Date are displayed at this end of the package.

Barcode Placeholder

NDC 054365-400-13  
 Cat. No. 260815

Single Use  
**ChloroPrep® With Tint**  
 2% w/v chlorhexidine gluconate (CHG)  
 and 70% v/v isopropyl alcohol (IPA)  
 Patient Preoperative Skin Preparation  
 Non-sterile Solution  
 Applicator is sterile if package is intact  
**26 ml APPLICATOR**

**Hi-Lite Orange®**

WARNING. FLAMMABLE. KEEP AWAY FROM FIRE OR FLAME.

**Drug Facts**

Active ingredients	Purposes
Chlorhexidine gluconate 2% w/v	Antiseptic
Isopropyl alcohol 70% v/v	Antiseptic

**Use** for the preparation of the patient's skin prior to surgery. Helps to reduce bacteria that potentially can cause skin infection.

**Warnings**  
 For external use only. Flammable, keep away from fire or flame. To reduce risk of fire, PREP CAREFULLY:

- do not use 26-ml applicator for head and neck surgery
- do not use on an area smaller than 8.4 in. x 8.4 in. Use a smaller applicator instead.
- solution contains alcohol and gives off flammable vapors
- avoid getting solution into hairy areas. Hair may take up to 1 hour to dry. Wet hair is flammable.
- do not drape or use ignition source (e.g., cautery, laser) until solution is completely dry (minimum of 3 minutes on hairless skin; up to 1 hour in hair)
- do not allow solution to pool
- remove wet materials from prep area

**Allergy Alert:** This product may cause a severe allergic reaction. Symptoms may include:

- wheezing/difficulty breathing
- shock
- facial swelling
- hives
- rash

If an allergic reaction occurs, stop use and seek medical help right away.

**Do not use**

- on patients allergic to chlorhexidine gluconate or any other ingredient in this product
- for lumbar puncture or in contact with the meninges
- on open skin wounds or as a general skin cleanser

**When using this product** keep out of eyes, ears, and mouth. May cause serious or permanent injury if permitted to enter and remain. If contact occurs, rinse with cold water right away and contact a doctor.

**Stop use and ask a doctor if** irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition.

**Keep out of reach of children.** If swallowed, get medical help or contact a Poison Control Center right away.

**Directions**

- use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns.
- use in a well ventilated area
- maximal treatment area for one applicator is approximately 13.2 in. x 13.2 in. (1126 cm<sup>2</sup>). Do not use 26-ml applicator for area smaller than 8.4 in. x 8.4 in. Use a smaller applicator instead.
- do not use 26-ml applicator for head and neck surgery
- remove applicator from package; do not touch sponge
- hold the applicator with the sponge down. Pinch wing **only once** to activate the ampules and release the antiseptic.
- wet the sponge by pressing and releasing the sponge against the treatment area until liquid is visible on the skin
- completely wet the treatment area with antiseptic
- dry surgical sites** (e.g., abdomen or arm): use gentle repeated back-and-forth strokes for 30 seconds
- moist surgical sites** (e.g., inguinal fold): use gentle repeated back-and-forth strokes for 2 minutes
- do not allow solution to pool; tuck prep towels to absorb solution, and then remove
- allow the solution to completely dry** (minimum of 3 minutes on hairless skin; up to 1 hour in hair). Do not blot or wipe away.
- discard the applicator after a single use along with any portion of the solution not required to cover the prep area. It is not necessary to use the entire amount available.

**Other information**

- store between 15–30 °C (59–86 °F)
- avoid freezing and excessive heat above 40 °C (104 °F)

**Inactive ingredients**  
 FD&C yellow #6 dye, USP purified water

**Questions?**

- www.chloroprep.com
- call 1-800-523-0502 (M-F 8 a.m.–5 p.m. CST)

CareFusion, El Paso, TX 79912 - Artwork No. 6-786815 - OXXXXXX  
 Made in the USA of US and imported parts

243

305

Lot/Exp Placeholder -  
 for reference only

8.25


1

60

80.5


Artwork No. 6-786825, 26-mL ChloroPrep with Tint (Teal) Applicator Lidding

(b) (4)



(b) (4)


Lot/Exp Placeholder -  
for reference only



(b) (4)

Part No. 6-773800-L  
26-ml ChloroPrep One-Step  
In-Line Mold Labels


(b) (4)



(b) (4)

(b) (4)

Part No. 6-773800-R  
26-ml ChloroPrep One-Step  
In-Line Mold Labels




(b) (4)

(b) (4)

Part No. 6-773815-L  
26-ml ChloroPrep with tint  
In-Line Mold Labels


(b) (4)

(b) (4)



(b) (4)

Part No. 6-773815-R  
26-ml ChloroPrep with tint  
In-Line Mold Labels



(b) (4)


(b) (4)

(b) (4)

(b) (4)

Part No. 6-773825-L  
26-ml ChloroPrep with tint  
In-Line Mold Labels

(b) (4)




(b) (4)

(b) (4)

Part No. 6-773825-R  
26-ml ChloroPrep with tint  
In-Line Mold Labels

(b) (4)



(b) (4)

---

**This is a representation of an electronic record that was signed electronically and this page is the manifestation of the electronic signature.**

---

/s/  
-----

VALERIE S PRATT  
09/12/2017