

Allegra Hives 24 Hour Tablets

5 Count Blister Print Mat

03.18.21

ALG Hives 24hr 5ct Blister FDA 803911-01 DSN4.ai
(PDF of art being shown is submitted at 100% actual size)





Allegra Hives 24 Hour 30 Count Tablets

30 Count Label

04.01.21

ALG Hives 24hr 30ct LBL FDA 803914-01 DSN4.ai
 (PDF of art being shown is submitted at 100% actual size)





Allegra Hives 24 Hour 45 Count Tablets

45 Count Label

04.01.21

ALG Hives 24hr 45ct LBL FDA 803916-01 DSN5.ai
 (PDF of art being shown is submitted at 100% actual size)




Allegra Hives 24 Hour 60 Count Tablets

60 Count Label

04.01.21

ALG Hives 24hr 60ct LBL FDA 803930-01 DSN5.ai
 (PDF of art being shown is submitted at 100% actual size)



Allegra Hives 24 Hour 30 Count Tablets
 30 Tablet Carton
 06.09.21
 ALG Hives 24hr 30ct CTN FDA 803913-01 DSN6.ai
 (PDF of art being shown is submitted at 100% actual size)



Drug Facts
 Fexofenadine HCl 180 mg
Allegra Hives 24 HR
 30 TABLETS
 NON-DROWSY
 This product will not prevent hives or an allergic skin reaction from occurring.
 Reduces hives and relieves itching due to hives (urticaria).
Uses
 Active ingredient (in each tablet) Purpose

Drug Facts (continued)
Other information
 ■ safety sealed; do not use if carton is opened or if printed foil inner seal on bottle is torn or missing
 ■ store between 20° and 25°C (68° and 77°F)
 ■ protect from excessive moisture
Inactive ingredients
 colloidal silicon dioxide, croscarmellose sodium, hypromellose, iron oxide blends, magnesium stearate, microcrystalline cellulose, polyethylene glycol, povidone, pregelatinized starch, titanium dioxide
Questions or comments?
 call toll-free 1-800-633-1610

The makers of Allegra® do not make store brand products. The trade dress of this Allegra® package is subject to trademark protection.
 Diet. By: Chatham, Inc. a Sanofi Company, Chattanooga, TN 37409-0219
 ©2021 Origin Germany

New!
Allegra Hives
 fexofenadine HCl tablet 180 mg/antihistamine **24 HR**
REDUCES HIVES and RELIEVES ITCHING due to hives
30 TABLETS
 NON-DROWSY
 NDC 41167-4126-4
 — NO ART HERE — WINDOW
 Actual Size

803913-01
 LOT
 EXP

 0 41167 41264 0

Drug Facts (continued)
Warnings
Severe Allergy Warning: Get emergency help immediately if you have hives along with any of the following symptoms:
 ■ trouble swallowing ■ dizziness or loss of consciousness
 ■ swelling of tongue ■ swelling in or around mouth
 ■ trouble speaking ■ wheezing or problems breathing
 ■ drooling
 These symptoms may be signs of anaphylactic shock. This condition can be life threatening if not treated by a health professional immediately. Symptoms of anaphylactic shock may occur when hives first appear or up to a few hours later.
Not a Substitute for Epinephrine. If your doctor has prescribed an epinephrine injector for "anaphylaxis" or severe allergy symptoms that could occur with your hives, never use this product as a substitute for the epinephrine injector. If you have been prescribed an epinephrine injector, you should carry it with you at all times.
Do not use
 ■ to prevent hives from any known cause such as:
 ■ foods ■ insect stings ■ medicines ■ latex or rubber gloves because this product will not stop hives from occurring. Avoiding the cause of your hives is the only way to prevent them. Hives can sometimes be serious. If you do not know the cause of your hives, see your doctor for a medical exam. Your doctor may be able to help you find a cause.
 ■ if you have ever had an allergic reaction to this product or any of its ingredients.

Drug Facts (continued)
Ask a doctor before use if you have
 ■ kidney disease. Your doctor should determine if you need a different dose.
 ■ hives that are an unusual color, look bruised or blistered
 ■ hives that do not itch
When using this product
 ■ do not take more than directed
 ■ do not take at the same time as aluminum or magnesium antacids
 ■ do not take with fruit juices (see Directions)
Stop use and ask a doctor if
 ■ an allergic reaction to this product occurs. Seek medical help right away.
 ■ symptoms do not improve after 3 days of treatment
 ■ the hives have lasted more than 6 weeks
If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.
Directions

adults and children 12 years of age and over	take one 180 mg tablet with water once a day; do not take more than 1 tablet in 24 hours
children under 12 years of age	do not use
adults 65 years of age and older	ask a doctor
consumers with kidney disease	ask a doctor

Allegra Hives 24 Hour 45 Count Tablets
 45 Tablet Carton
 06.09.21
 ALG Hives 24hr 45ct CTN FDA 803915-01 DSN5.ai
 (PDF of art being shown is submitted at 100% actual size)



Drug Facts
 Active ingredient (in each tablet) Purpose
 Fexofenadine HCl 180 mg...
 Uses
 Reduces hives and relieves itching due to hives (urticaria).
 This product will not prevent hives or an allergic skin reaction from occurring.

Allegra Hives 24 HR
 NON-DROWSY
 45 TABLETS

Drug Facts (continued)
Other information
 ■ safety sealed; do not use if carton is opened or if printed foil inner seal on bottle is torn or missing
 ■ store between 20° and 25°C (68° and 77°F)
 ■ protect from excessive moisture
Inactive ingredients
 colloidal silicon dioxide, croscarmellose sodium, hypromellose, iron oxide blends, magnesium stearate, microcrystalline cellulose, polyethylene glycol, povidone, pregelatinized starch, titanium dioxide
Questions or comments?
 call toll-free 1-800-633-1610

New!
Allegra Hives
 fexofenadine HCl tablet 180 mg/antihistamine **24 HR**
REDUCES HIVES and RELIEVES ITCHING due to hives
 NON-DROWSY
 NDC 41167-4126-7
 45 TABLETS
 — NO ART HERE — WINDOW
 Actual Size

Allegra Hives 24HR

Drug Facts (continued)
Warnings
Severe Allergy Warning: Get emergency help immediately if you have hives along with any of the following symptoms:
 ■ trouble swallowing ■ dizziness or loss of consciousness
 ■ swelling of tongue ■ swelling in or around mouth
 ■ trouble speaking ■ wheezing or problems breathing
 ■ drooling
 These symptoms may be signs of anaphylactic shock. This condition can be life threatening if not treated by a health professional immediately. Symptoms of anaphylactic shock may occur when hives first appear or up to a few hours later.
Not a Substitute for Epinephrine. If your doctor has prescribed an epinephrine injector for "anaphylaxis" or severe allergy symptoms that could occur with your hives, never use this product as a substitute for the epinephrine injector. If you have been prescribed an epinephrine injector, you should carry it with you at all times.
Do not use
 ■ to prevent hives from any known cause such as:
 ■ foods ■ insect stings ■ medicines ■ latex or rubber gloves because this product will not stop hives from occurring. Avoiding the cause of your hives is the only way to prevent them. Hives can sometimes be serious. If you do not know the cause of your hives, see your doctor for a medical exam. Your doctor may be able to help you find a cause.
 ■ if you have ever had an allergic reaction to this product or any of its ingredients.

Drug Facts (continued)
Ask a doctor before use if you have
 ■ kidney disease. Your doctor should determine if you need a different dose.
 ■ hives that are an unusual color, look bruised or blistered
 ■ hives that do not itch
When using this product
 ■ do not take more than directed
 ■ do not take at the same time as aluminum or magnesium antacids
 ■ do not take with fruit juices (see Directions)
Stop use and ask a doctor if
 ■ an allergic reaction to this product occurs. Seek medical help right away.
 ■ symptoms do not improve after 3 days of treatment
 ■ the hives have lasted more than 6 weeks
If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.
Directions

adults and children 12 years of age and over	take one 180 mg tablet with water once a day; do not take more than 1 tablet in 24 hours
children under 12 years of age	do not use
adults 65 years of age and older	ask a doctor
consumers with kidney disease	ask a doctor

803915-01
 LOT
 EXP



Allegra Hives 24 Hour 60 Count Tablets
 60 Tablet Carton
 06.09.21
 ALG Hives 24hr 60ct CTN FDA 803929-01 DSN5.ai
 (PDF of art being shown is submitted at 100% actual size)



Drug Facts
 Fexofenadine HCl 180 mg
Active ingredient (in each tablet) Purpose
Uses
 Reduces hives and relieves itching due to hives (urticaria).
 This product will not prevent hives or an allergic skin reaction from occurring.

Allegra Hives 24HR
 NON-DROWSY
 60 TABLETS

Drug Facts (continued)
Other information
 ■ safety sealed; do not use if carton is opened or if printed foil inner seal on bottle is torn or missing
 ■ store between 20° and 25°C (68° and 77°F)
 ■ protect from excessive moisture
Inactive ingredients
 colloidal silicon dioxide, croscarmellose sodium, hypromellose, iron oxide blends, magnesium stearate, microcrystalline cellulose, polyethylene glycol, povidone, pregelatinized starch, titanium dioxide
Questions or comments?
 call toll-free 1-800-633-1610

New!
Allegra Hives
 fexofenadine HCl tablet 180 mg/antihistamine **24HR**
NON-DROWSY
 NDC 41167-4126-1
60 TABLETS

Allegra Hives 24HR

Drug Facts (continued)
Warnings
Severe Allergy Warning: Get emergency help immediately if you have hives along with any of the following symptoms:
 ■ trouble swallowing ■ dizziness or loss of consciousness
 ■ swelling of tongue ■ swelling in or around mouth
 ■ trouble speaking ■ wheezing or problems breathing
 ■ drooling
 These symptoms may be signs of anaphylactic shock. This condition can be life threatening if not treated by a health professional immediately. Symptoms of anaphylactic shock may occur when hives first appear or up to a few hours later.
Not a Substitute for Epinephrine: If your doctor has prescribed an epinephrine injector for "anaphylaxis" or severe allergy symptoms that could occur with your hives, never use this product as a substitute for the epinephrine injector. If you have been prescribed an epinephrine injector, you should carry it with you at all times.
Do not use
 ■ to prevent hives from any known cause such as:
 ■ foods ■ insect stings ■ medicines ■ latex or rubber gloves because this product will not stop hives from occurring. Avoiding the cause of your hives is the only way to prevent them. Hives can sometimes be serious. If you do not know the cause of your hives, see your doctor for a medical exam. Your doctor may be able to help you find a cause.
 ■ if you have ever had an allergic reaction to this product or any of its ingredients.

Drug Facts (continued)
Ask a doctor before use if you have
 ■ kidney disease. Your doctor should determine if you need a different dose.
 ■ hives that are an unusual color, look bruised or blistered
 ■ hives that do not itch
When using this product
 ■ do not take more than directed
 ■ do not take at the same time as aluminum or magnesium antacids
 ■ do not take with fruit juices (see Directions)
Stop use and ask a doctor if
 ■ an allergic reaction to this product occurs. Seek medical help right away.
 ■ symptoms do not improve after 3 days of treatment
 ■ the hives have lasted more than 6 weeks
If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.
Directions

adults and children 12 years of age and over	take one 180 mg tablet with water once a day; do not take more than 1 tablet in 24 hours
children under 12 years of age	do not use
adults 65 years of age and older	ask a doctor
consumers with kidney disease	ask a doctor

The makers of Allegra® do not make store brand products. The trade dress of this Allegra® package is subject to trademark protection.
 Div. By: Chatham, Inc., a Sanofi Company, Chattanooga, TN 37409-0219
 ©2021 Origin Germany

803929-01
 LOT
 EXP

 0 41167 41261 9

