This label may not be the latest approved by FDA.
For current labeling information, please visit https://www.fda.gov/drugsatfda

Allegra® ALLERGY 24 HR

NON-DROWSY

1 TABLET

NDC 41167-4124-2

Allegra® ALLERGY

fexofenadine HCl tablet 180 mg/antihistamine 24 HR

INDOOR / OUTDOOR ALLERGY RELIEF

- ✓ Sneezing
- ✓ Runny Nose
- ✓ Itchy, Watery Eyes
- ✓ Itchy Nose or Throat

1 TABLET
Actual Size

Allegra® ALLERGY

The makers of Allegra® do not make store brand products.
The trade dress of this Allegra® package is subject to trademark protection.

SAMPLE:
NOT FOR
RETAIL SALE

Dist By: Chattem, Inc., a Sanofi Company, Chattanooga, TN 37409-0219 ©2022 Origin India

863472

LOT
EXP

Drug Facts

Active ingredient (in each tablet) — **Purpose**
Fexofenadine HCl 180 mg ..... Antihistamine

Uses
temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:
- runny nose
- sneezing
- itchy, watery eyes
- itching of the nose or throat

Warnings
Do not use if you have ever had an allergic reaction to this product or any of its ingredients.
Ask a doctor before use if you have kidney disease. Your doctor should determine if you need a different dose.
When using this product
- do not take more than directed
- do not take at the same time as aluminum or magnesium antacids
- do not take with fruit juices (see Directions)

Stop use and ask a doctor if an allergic reaction to this product occurs. Seek medical help right away.
If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Drug Facts (continued)

Directions

adults and children 12 years of age and over	take one 180 mg tablet with water once a day; do not take more than 1 tablet in 24 hours
children under 12 years of age	do not use
adults 65 years of age and older	ask a doctor
consumers with kidney disease	ask a doctor

Other information
- safety sealed: do not use if carton is opened or if individual blister unit is torn or opened
- store between 20° and 25°C (68° and 77°F)
- protect from excessive moisture

Inactive ingredients
colloidal silicon dioxide, croscarmellose sodium, hypromellose, iron oxide blends, magnesium stearate, microcrystalline cellulose, polyethylene glycol, povidone, pregelatinized starch, titanium dioxide

Questions or comments?
call toll-free **1-866-844-2798** or www.allegra.com

(4) (b)

This label may not be the latest approved by FDA.
For current labeling information, please visit https://www.fda.gov/drugsatfda



(b) (4)

This label may not be the latest approved by FDA.
For current labeling information, please visit https://www.fda.gov/drugsatfda

Reference ID: 5218613

(b) (4)



This label may not be the latest approved by FDA.
For current labeling information, please visit https://www.fda.gov/drugsatfda

Reference ID: 5218613

(b) (4)

Drug Facts

Active ingredient (in each tablet) **Purpose**
Fexofenadine HCl 180 mg ..... Antihistamine

Uses
temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:
- runny nose
- sneezing
- itchy, watery eyes
- itching of the nose or throat

Warnings
Do not use if you have ever had an allergic reaction to this product or any of its ingredients.

Ask a doctor before use if you have kidney disease. Your doctor should determine if you need a different dose.

When using this product
- do not take more than directed
- do not take at the same time as aluminum or magnesium antacids
- do not take with fruit juices (see Directions)

Stop use and ask a doctor if an allergic reaction to this product occurs. Seek medical help right away.

If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

▶

Drug Facts (continued)

Directions

adults and children 12 years of age and over	take one 180 mg tablet with water once a day; do not take more than 1 tablet in 24 hours
children under 12 years of age	do not use
adults 65 years of age and older	ask a doctor
consumers with kidney disease	ask a doctor

Other information
- safety sealed: do not use if carton is opened or if individual blister units are torn or opened
- store between 20° and 25°C (68° and 77°F)
- protect from excessive moisture

Inactive ingredients
colloidal silicon dioxide, croscarmellose sodium, hypromellose, iron oxide blends, magnesium stearate, microcrystalline cellulose, polyethylene glycol, povidone, pregelatinized starch, titanium dioxide

Questions or comments?
call toll-free **1-866-844-2798** or www.allegra.com

Allegra® ALLERGY 24HR

Dist. By: Chattem, Inc., a Sanofi Company, Chattanooga, TN 37409-0219 ©2022 Origin India 858012

LOT EXP

The makers of Allegra® do not make store brand products.
The trade dress of this Allegra® package is subject to trademark protection.

SAMPLE:
NOT FOR RETAIL SALE

Allegra® ALLERGY

Actual Size
5 TABLETS

✓ Sneezing
✓ Runny Nose
✓ Itchy, Watery Eyes
✓ Itchy Nose or Throat

INDOOR / OUTDOOR ALLERGY RELIEF

Allegra® ALLERGY 24 HR
fexofenadine HCl tablet
180 mg/antihistamine

NON-DROWSY

NDC 41167-4120-0

Allegra® ALLERGY 24 HR
5 TABLETS
NON-DROWSY

858012

This label may not be the latest approved by FDA.
For current labeling information, please visit https://www.fda.gov/drugsatfda

Reference ID: 5218613

(b) (4)



This label may not be the latest approved by FDA.
For current labeling information, please visit https://www.fda.gov/drugsatfda

849966

NON-DROWSY

Allegra® ALLERGY 24 HR

30 TABLETS

NDC 41167-4120-3

NON-DROWSY

Allegra® ALLERGY fexofenadine HCl tablet 180 mg/antihistamine 24 HR

INDOOR / OUTDOOR ALLERGY RELIEF

- ✓ Sneezing
- ✓ Runny Nose
- ✓ Itchy, Watery Eyes
- ✓ Itchy Nose or Throat

30 TABLETS
Actual Size

LOT

EXP

0 41167 41251 0

Allegra® ALLERGY 24 HR

Drug Facts

Active ingredient (in each tablet) **Purpose**
Fexofenadine HCl 180 mg ..... Antihistamine

Uses
temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:
- runny nose
- sneezing
- itchy, watery eyes
- itching of the nose or throat

Warnings
Do not use if you have ever had an allergic reaction to this product or any of its ingredients.

Ask a doctor before use if you have kidney disease. Your doctor should determine if you need a different dose.

When using this product
- do not take more than directed
- do not take at the same time as aluminum or magnesium antacids
- do not take with fruit juices (see Directions)

Stop use and ask a doctor if an allergic reaction to this product occurs. Seek medical help right away.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. ▶

Drug Facts (continued)

Directions

adults and children 12 years of age and over	take one 180 mg tablet with water once a day; do not take more than 1 tablet in 24 hours
children under 12 years of age	do not use
adults 65 years of age and older	ask a doctor
consumers with kidney disease	ask a doctor

Other information
- safety sealed: do not use if carton is opened or if printed foil inner seal on bottle is torn or missing
- store between 20° and 25°C (68° and 77°F)
- protect from excessive moisture

Inactive ingredients
colloidal silicon dioxide, croscarmellose sodium, hypromellose, iron oxide blends, magnesium stearate, microcrystalline cellulose, polyethylene glycol, povidone, pregelatinized starch, titanium dioxide

Questions or comments?
call toll-free **1-866-844-2798** or www.allegra.com

The makers of Allegra® do not make store brand products. The trade dress of this Allegra® package is subject to trademark protection.
Dist. By: Chattem, Inc., a Sanofi Company, Chattanooga, TN 37409-0219 ©2022 Origin India 849966

849966

(b) (4)

This label may not be the latest approved by FDA.
For current labeling information, please visit https://www.fda.gov/drugsatfda

(b) (4)



Active ingredient (in each tablet)	***Purpose***
Fexofenadine HCl 180 mg	Antihistamine

Uses temporarily relieves these symptoms due to hay fever or other upper respiratory allergies: ■ runny nose ■ sneezing ■ itchy, watery eyes ■ itching of the nose or throat ***Warnings*** **Do not use** if you have ever had an allergic reaction to this product or any of its ingredients. **Ask a doctor before use if you have** kidney disease. Your doctor should determine if you need a different dose. **When using this product** ■ do not take more than directed ■ do not take at the same time as aluminum or magnesium antacids ■ do not take with fruit juices (see Directions) **Stop use and ask a doctor if** an allergic reaction to this product occurs. Seek medical help right away. **If pregnant or breast-feeding,** ask a health professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. ***Directions*** ■ **adults and children 12 years of age and over** take one 180 mg tablet with water once a day; do not take more than 1 tablet in 24 hours ■ **children under 12 years of age** do not use ■ **adults 65 years of age and older** ask a doctor ■ **consumers with kidney disease** ask a doctor ***Other information*** ■ safety sealed: do not use if carton was opened or if printed foil inner seal on bottle is torn or missing ■ store between 20° and 25°C (68° and 77°F) ▶

■ protect from excessive moisture
Inactive ingredients colloidal silicon dioxide, croscarmellose sodium, hypromellose, iron oxide blends, magnesium stearate, microcrystalline cellulose, polyethylene glycol, povidone, pregelatinized starch, titanium dioxide
Questions or comments? call toll-free **1-866-844-2798** or www.allegra.com

Dist. By: Chattem, Inc., a Sanofi Company
Chattanooga, TN 37409-0219
©2022 Origin India



This label may not be the latest approved by FDA.
For current labeling information, please visit https://www.fda.gov/drugsatfda

849968

Allegra® ALLERGY 24 HR

NON-DROWSY

~~30~~ 40 TABLETS

33% MORE | 10 FREE TABLETS

NDC 41167-4121-4

Allegra® ALLERGY

fexofenadine HCl tablet
180 mg/antihistamine 24 HR

INDOOR / OUTDOOR ALLERGY RELIEF

- ✓ Sneezing
- ✓ Runny Nose
- ✓ Itchy, Watery Eyes
- ✓ Itchy Nose or Throat

40
~~30~~ TABLETS
Actual Size

LOT

EXP

0 41167 41214 5

Drug Facts

Active ingredient (in each tablet) **Purpose**
Fexofenadine HCl 180 mg ..... Antihistamine

Uses
temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:
- runny nose
- sneezing
- itchy, watery eyes
- itching of the nose or throat

Warnings
Do not use if you have ever had an allergic reaction to this product or any of its ingredients.

Ask a doctor before use if you have kidney disease. Your doctor should determine if you need a different dose.

When using this product
- do not take more than directed
- do not take at the same time as aluminum or magnesium antacids
- do not take with fruit juices (see Directions)

Stop use and ask a doctor if an allergic reaction to this product occurs. Seek medical help right away.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. ▶

Drug Facts (continued)

Directions

adults and children 12 years of age and over	take one 180 mg tablet with water once a day; do not take more than 1 tablet in 24 hours
children under 12 years of age	do not use
adults 65 years of age and older	ask a doctor
consumers with kidney disease	ask a doctor

Other information
- safety sealed: do not use if carton is opened or if printed foil inner seal on bottle is torn or missing
- store between 20° and 25°C (68° and 77°F)
- protect from excessive moisture

Inactive ingredients
colloidal silicon dioxide, croscarmellose sodium, hypromellose, iron oxide blends, magnesium stearate, microcrystalline cellulose, polyethylene glycol, povidone, pregelatinized starch, titanium dioxide

Questions or comments?
call toll-free **1-866-844-2798** or www.allegra.com

The makers of Allegra® do not make store brand products. The trade dress of this Allegra® package is subject to trademark protection.
Dist. By: Chattem, Inc., a Sanofi Company, Chattanooga, TN 37409-0219 ©2022 Origin India 849968

849968

(b) (4)

This label may not be the latest approved by FDA.
For current labeling information, please visit https://www.fda.gov/drugsatfda

(b) (4)



Active ingredient (in each tablet) ***Purpose***
Fexofenadine HCl 180 mg.....Antihistamine
Uses temporarily relieves these symptoms due to hay fever or other upper respiratory allergies: ■ runny nose ■ sneezing ■ itchy, watery eyes ■ itching of the nose or throat ***Warnings*** **Do not use** if you have ever had an allergic reaction to this product or any of its ingredients. **Ask a doctor before use if you have** kidney disease. Your doctor should determine if you need a different dose.
When using this product ■ do not take more than directed ■ do not take at the same time as aluminum or magnesium antacids ■ do not take with fruit juices (see Directions) **Stop use and ask a doctor if** an allergic reaction to this product occurs. Seek medical ▶

help right away. **If pregnant or breast-feeding,** ask a health professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. ***Directions*** ■ **adults and children 12 years of age and over** take one 180 mg tablet with water once a day; do not take more than 1 tablet in 24 hours ■ **children under 12 years of age** do not use ■ **adults 65 years of age and older** ask a doctor ■ **consumers with kidney disease** ask a doctor
Other information ■ safety sealed: do not use if carton was opened or if printed foil inner seal on bottle is torn or missing ■ store between 20° and 25°C (68° and 77°F) ▶

■ protect from excessive moisture
Inactive ingredients colloidal silicon dioxide, croscarmellose sodium, hypromellose, iron oxide blends, magnesium stearate, microcrystalline cellulose, polyethylene glycol, povidone, pregelatinized starch, titanium dioxide
Questions or comments? call toll-free **1-866-844-2798** or www.allegra.com

Dist. By: Chattem, Inc., a Sanofi Company
Chattanooga, TN 37409-0219
©2022 Origin India

NDC 41167-4121-4

Allegra®
ALLERGY 24 HR
fexofenadine HCl tablet
180 mg/antihistamine

INDOOR / OUTDOOR ALLERGY RELIEF

40 TABLETS

This label may not be the latest approved by FDA.
For current labeling information, please visit https://www.fda.gov/drugsatfda

849969

Allegra® ALLERGY 24 HR

NON-DROWSY

45 TABLETS

NDC 41167-4120-4

Allegra® ALLERGY

fexofenadine HCl tablet
180 mg/antihistamine 24 HR

INDOOR / OUTDOOR ALLERGY RELIEF

- ✓ Sneezing
- ✓ Runny Nose
- ✓ Itchy, Watery Eyes
- ✓ Itchy Nose or Throat

45 TABLETS
Actual Size

LOT

EXP

0 41167 41274 9

Allegra® ALLERGY 24 HR

Drug Facts

Active ingredient (in each tablet) **Purpose**
Fexofenadine HCl 180 mg..... Antihistamine

Uses
temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:
- runny nose
- sneezing
- itchy, watery eyes
- itching of the nose or throat

Warnings
Do not use if you have ever had an allergic reaction to this product or any of its ingredients.

Ask a doctor before use if you have kidney disease. Your doctor should determine if you need a different dose.

When using this product
- do not take more than directed
- do not take at the same time as aluminum or magnesium antacids
- do not take with fruit juices (see Directions)

Stop use and ask a doctor if an allergic reaction to this product occurs. Seek medical help right away.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. ▶

Drug Facts (continued)

Directions

adults and children 12 years of age and over	take one 180 mg tablet with water once a day; do not take more than 1 tablet in 24 hours
children under 12 years of age	do not use
adults 65 years of age and older	ask a doctor
consumers with kidney disease	ask a doctor

Other information
- safety sealed: do not use if carton is opened or if printed foil inner seal on bottle is torn or missing
- store between 20° and 25°C (68° and 77°F)
- protect from excessive moisture

Inactive ingredients
colloidal silicon dioxide, croscarmellose sodium, hypromellose, iron oxide blends, magnesium stearate, microcrystalline cellulose, polyethylene glycol, povidone, pregelatinized starch, titanium dioxide

Questions or comments?
call toll-free **1-866-844-2798** or www.allegra.com

The makers of Allegra® do not make store brand products. The trade dress of this Allegra® package is subject to trademark protection.
Dist. By: Chattem, Inc., a Sanofi Company, Chattanooga, TN 37409-0219 ©2022 Origin India 849969

849969

(b) (4)

This label may not be the latest approved by FDA.
For current labeling information, please visit https://www.fda.gov/drugsatfda

(b) (4)

NDC 41167-4120-4

NON-DROWSY
Allegra®
ALLERGY 24 HR
fexofenadine HCl tablet
180 mg/antihistamine

45 TABLETS

INDOOR / OUTDOOR ALLERGY RELIEF
- ✓ Sneezing
- ✓ Runny Nose
- ✓ Itchy, Watery Eyes
- ✓ Itchy Nose or Throat

Dist. By: Chattem, Inc., a Sanofi Company
Chattanooga, TN 37409
©2022
Origin India

849984

Peel Here

Active ingredient (in each tablet) ***Purpose***
Fexofenadine HCl 180 mg.....Antihistamine
Uses temporarily relieves these symptoms due to hay fever or other upper respiratory allergies: ■ runny nose ■ sneezing ■ itchy, watery eyes ■ itching of the nose or throat ***Warnings*** **Do not use** if you have ever had an allergic reaction to this product or any of its ingredients. **Ask a doctor before use if you have** kidney disease. Your doctor should determine if you need a different dose.
When using this product ■ do not take more than directed ■ do not take at the same time as aluminum or magnesium antacids ■ do not take with fruit juices (see Directions) **Stop use and ask a doctor if** an allergic reaction to this product occurs. Seek medical help right away. **If pregnant or breast-feeding,** ask a health professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. ***Directions*** ■ **adults and children 12 years of age and over** take one 180 mg tablet with water once a day; do not take more than 1 tablet in 24 hours ■ **children under 12 years of age** do not use ■ **adults 65 years of age and older** ask a doctor ■ **consumers with kidney disease** ask a doctor
Other information ■ safety sealed: do not use if carton was opened or if printed foil inner seal on bottle is torn or missing ■ store between 20° and 25°C (68° and 77°F) ■ protect from excessive moisture
Inactive ingredients colloidal silicon dioxide, croscarmellose sodium, hypromellose, iron oxide blends, magnesium stearate, microcrystalline cellulose, polyethylene glycol, povidone, pregelatinized starch, titanium dioxide
Questions or comments? call toll-free **1-866-844-2798** or www.allegra.com

NDC 41167-4120-4

Allegra®
ALLERGY 24 HR
fexofenadine HCl tablet
180 mg/antihistamine

INDOOR / OUTDOOR ALLERGY RELIEF

45 TABLETS

Dist. By: Chattem, Inc., a Sanofi Company
Chattanooga, TN 37409-0219
©2022 Origin India

This label may not be the latest approved by FDA.
For current labeling information, please visit https://www.fda.gov/drugsatfda

(b) (4)



Active ingredient (in each tablet)	***Purpose***
Fexofenadine HCl 180 mg	Antihistamine

Uses temporarily relieves these symptoms due to hay fever or other upper respiratory allergies: ■ runny nose ■ sneezing ■ itchy, watery eyes ■ itching of the nose or throat ***Warnings*** **Do not use** if you have ever had an allergic reaction to this product or any of its ingredients. **Ask a doctor before use if you have** kidney disease. Your doctor should determine if you need a different dose. **When using this product** ■ do not take more than directed ■ do not take at the same time as aluminum or magnesium antacids ■ do not take with fruit juices (see Directions) **Stop use and ask a doctor if** an allergic reaction to this product occurs. Seek medical help right away. **If pregnant or breast-feeding,** ask a health professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. ***Directions*** ■ **adults and children 12 years of age and over** take one 180 mg tablet with water once a day; do not take more than 1 tablet in 24 hours ■ **children under 12 years of age** do not use ■ **adults 65 years of age and older** ask a doctor ■ **consumers with kidney disease** ask a doctor ***Other information*** ■ safety sealed: do not use if printed foil inner seal on bottle is torn or missing ■ store between 20° and 25°C (68° and 77°F) ▶

■ protect from excessive moisture
Inactive ingredients colloidal silicon dioxide, croscarmellose sodium, hypromellose, iron oxide blends, magnesium stearate, microcrystalline cellulose, polyethylene glycol, povidone, pregelatinized starch, titanium dioxide
Questions or comments? call toll-free **1-866-844-2798** or www.allegra.com

Dist. By: Chattem, Inc., a Sanofi Company
Chattanooga, TN 37409-0219
©2022 Origin India

NDC 41167-4124-5

Allegra®
ALLERGY 24HR
fexofenadine HCl tablet
180 mg/antihistamine

INDOOR / OUTDOOR ALLERGY RELIEF

55 TABLETS

This label may not be the latest approved by FDA.
For current labeling information, please visit https://www.fda.gov/drugsatfda



(b) (4)

This label may not be the latest approved by FDA.
For current labeling information, please visit https://www.fda.gov/drugsatfda

(b) (4)

NDC 41167-4121-1

Dist. By: Chattem, Inc.,
a Sanofi Company
Chattanooga, TN 37409
©2022
Origin India

Peel Here

INDOOR / OUTDOOR ALLERGY RELIEF
- ✓ Sneezing
- ✓ Runny Nose
- ✓ Itchy, Watery Eyes
- ✓ Itchy Nose or Throat

NON-DROWSY
Allegra®
ALLERGY 24 HR
fexofenadine HCl tablet
180 mg/antihistamine

60 TABLETS

849977

Active ingredient (in each tablet) ***Purpose***
Fexofenadine HCl 180 mg.....Antihistamine

Uses temporarily relieves these symptoms due to hay fever or other upper respiratory allergies: ■ runny nose ■ sneezing ■ itchy, watery eyes ■ itching of the nose or throat

Warnings **Do not use** if you have ever had an allergic reaction to this product or any of its ingredients. **Ask a doctor before use if you have** kidney disease. Your doctor should determine if you need a different dose.

When using this product ■ do not take more than directed ■ do not take at the same time as aluminum or magnesium antacids ■ do not take with fruit juices (see Directions) **Stop use and ask a doctor if** an allergic reaction to this product occurs. Seek medical help right away. **If pregnant or breast-feeding,** ask a health professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

Directions ■ **adults and children 12 years of age and over** take one 180 mg tablet with water once a day; do not take more than 1 tablet in 24 hours ■ **children under 12 years of age** do not use ■ **adults 65 years of age and older** ask a doctor ■ **consumers with kidney disease** ask a doctor

Other information ■ safety sealed: do not use if carton was opened or if printed foil inner seal on bottle is torn or missing ■ store between 20° and 25°C (68° and 77°F) ■ protect from excessive moisture

Inactive ingredients colloidal silicon dioxide, croscarmellose sodium, hypromellose, iron oxide blends, magnesium stearate, microcrystalline cellulose, polyethylene glycol, povidone, pregelatinized starch, titanium dioxide

Questions or comments? call toll-free **1-866-844-2798** or www.allegra.com

Dist. By: Chattem, Inc., a Sanofi Company
Chattanooga, TN 37409-0219
©2022 Origin India

NDC 41167-4121-1

Allegra®
ALLERGY 24 HR
fexofenadine HCl tablet
180 mg/antihistamine

INDOOR / OUTDOOR ALLERGY RELIEF

60 TABLETS

This label may not be the latest approved by FDA.
For current labeling information, please visit https://www.fda.gov/drugsatfda

849972

NON-DROWSY

Allegra® ALLERGY 24 HR

70 TABLETS

VALUE SIZE!
Compare to 30 count

NON-DROWSY

NDC 41167-4120-6

Allegra® ALLERGY
fexofenadine HCl tablet
180 mg/antihistamine 24 HR

INDOOR / OUTDOOR ALLERGY RELIEF
- ✓ Sneezing
- ✓ Runny Nose
- ✓ Itchy, Watery Eyes
- ✓ Itchy Nose or Throat

70 TABLETS
Actual Size

LOT

EXP

0 41167 41206 0

Allegra® ALLERGY 24 HR

Drug Facts

Active ingredient (in each tablet) **Purpose**
Fexofenadine HCl 180 mg..... Antihistamine

Uses
temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:
- runny nose
- sneezing
- itchy, watery eyes
- itching of the nose or throat

Warnings
Do not use if you have ever had an allergic reaction to this product or any of its ingredients.

Ask a doctor before use if you have kidney disease. Your doctor should determine if you need a different dose.

When using this product
- do not take more than directed
- do not take at the same time as aluminum or magnesium antacids
- do not take with fruit juices (see Directions)

Stop use and ask a doctor if an allergic reaction to this product occurs. Seek medical help right away.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. ▶

Drug Facts (continued)

Directions

adults and children 12 years of age and over	take one 180 mg tablet with water once a day; do not take more than 1 tablet in 24 hours
children under 12 years of age	do not use
adults 65 years of age and older	ask a doctor
consumers with kidney disease	ask a doctor

Other information
- safety sealed: do not use if carton is opened or if printed foil inner seal on bottle is torn or missing
- store between 20° and 25°C (68° and 77°F)
- protect from excessive moisture

Inactive ingredients
colloidal silicon dioxide, croscarmellose sodium, hypromellose, iron oxide blends, magnesium stearate, microcrystalline cellulose, polyethylene glycol, povidone, pregelatinized starch, titanium dioxide

Questions or comments?
call toll-free **1-866-844-2798** or www.allegra.com

The makers of Allegra® do not make store brand products. The trade dress of this Allegra® package is subject to trademark protection.
Dist. By: Chattem, Inc., a Sanofi Company, Chattanooga, TN 37409-0219 ©2022 Origin India 849972

849972

(b) (4)

This label may not be the latest approved by FDA.
For current labeling information, please visit https://www.fda.gov/drugsatfda

(b) (4)



Active ingredient (in each tablet) ***Purpose***
Fexofenadine HCl 180 mg.....Antihistamine
Uses temporarily relieves these symptoms due to hay fever or other upper respiratory allergies: ■ runny nose ■ sneezing ■ itchy, watery eyes ■ itching of the nose or throat
Warnings **Do not use** if you have ever had an allergic reaction to this product or any of its ingredients. **Ask a doctor before use if you have** kidney disease. Your doctor should determine if you need a different dose. **When using this product** ■ do not take more than directed ■ do not take at the same time as aluminum or magnesium antacids ■ do not take with fruit juices (see Directions) **Stop use and ask a doctor if** an allergic reaction to this product occurs. Seek medical help right away. **If pregnant or breast-feeding,** ask a health professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.
Directions ■ **adults and children 12 years of age and over** take one 180 mg tablet with water once a day; do not take more than 1 tablet in 24 hours ■ **children under 12 years of age** do not use ■ **adults 65 years of age and older** ask a doctor ■ **consumers with kidney disease** ask a doctor
Other information ■ safety sealed: do not use if carton was opened or if printed foil inner seal on bottle is torn or missing ■ store between 20° and 25°C (68° and 77°F) ▶

■ protect from excessive moisture
Inactive ingredients colloidal silicon dioxide, croscarmellose sodium, hypromellose, iron oxide blends, magnesium stearate, microcrystalline cellulose, polyethylene glycol, povidone, pregelatinized starch, titanium dioxide
Questions or comments? call toll-free **1-866-844-2798** or www.allegra.com

Dist. By: Chattem, Inc., a Sanofi Company, Chattanooga, TN 37409-0219 ©2022 Origin India

NDC 41167-4120-6

Allegra®
ALLERGY 24 HR
fexofenadine HCl tablet
180 mg/antihistamine

INDOOR / OUTDOOR ALLERGY RELIEF

70 TABLETS

This label may not be the latest approved by FDA.
For current labeling information, please visit https://www.fda.gov/drugsatfda

849973

VALUE PACK! Compare to 30 count

NON-DROWSY

NDC 41167-4124-0

Allegra® ALLERGY

fexofenadine HCl tablet
180 mg/antihistamine 24 HR

INDOOR / OUTDOOR ALLERGY RELIEF
- ✓ Sneezing
- ✓ Runny Nose
- ✓ Itchy, Watery Eyes
- ✓ Itchy Nose or Throat

90 TABLETS
Actual Size

Allegra® ALLERGY 24 HR

0 41167 41240 4

849973

— Actual Size —

Allegra® ALLERGY 24 HR

90 Count Tablet Bottle

The makers of Allegra® do not make store brand products. The trade dress of this Allegra® package is subject to trademark protection.

Dist. By: Chattem, Inc., a Sanofi Company
Chattanooga, TN 37409-0219 ©2022
Origin India 849973

Questions or comments?
call toll-free 1-866-844-2798 or
www.allegra.com

Drug Facts

Active ingredient (in each tablet)	Purpose
Fexofenadine HCl 180 mg	Antihistamine

Uses
temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:
- runny nose
- itchy, watery eyes
- sneezing
- itching of the nose or throat

Warnings
Do not use if you have ever had an allergic reaction to this product or any of its ingredients.

Ask a doctor before use if you have kidney disease. Your doctor should determine if you need a different dose.

When using this product
- do not take more than directed
- do not take at the same time as aluminum or magnesium antacids
- do not take with fruit juices (see Directions)

Stop use and ask a doctor if an allergic reaction to this product occurs. Seek medical help right away.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Drug Facts (continued)

Directions

adults and children 12 years of age and over	take one 180 mg tablet with water once a day; do not take more than 1 tablet in 24 hours
children under 12 years of age	do not use
adults 65 years of age and older	ask a doctor
consumers with kidney disease	ask a doctor

Other information
- safety sealed: do not use if carton is opened or if printed foil inner seal on bottle is torn or missing
- store between 20° and 25°C (68° and 77°F)
- protect from excessive moisture

Inactive ingredients
colloidal silicon dioxide, croscarmellose sodium, hypromellose, iron oxide blends, magnesium stearate, microcrystalline cellulose, polyethylene glycol, povidone, pregelatinized starch, titanium dioxide

LOT

EXP

NON-DROWSY

Allegra® ALLERGY 24 HR

90 TABLETS

Reference ID: 5218613

This label may not be the latest approved by FDA.
For current labeling information, please visit https://www.fda.gov/drugsatfda

(b) (4)

NDC 41167-4124-0 **NON-DROWSY**

Allegra®
ALLERGY 24 HR
fexofenadine HCl tablet
180 mg/antihistamine

INDOOR / OUTDOOR ALLERGY RELIEF
- ✓ Sneezing ✓ Runny Nose
- ✓ Itchy, Watery Eyes
- ✓ Itchy Nose or Throat

90 TABLETS

Active ingredient (in each tablet) ***Purpose***
Fexofenadine HCl 180 mg.....Antihistamine

Uses temporarily relieves these symptoms due to hay fever or other upper respiratory allergies: ■ runny nose ■ sneezing ■ itchy, watery eyes ■ itching of the nose or throat

Warnings **Do not use** if you have ever had an allergic reaction to this product or any of its ingredients. **Ask a doctor before use if you have** kidney disease. Your doctor should determine if you need a different dose. **When using this product** ■ do not take more than directed ■ do not take at the same time as aluminum or magnesium antacids ■ do not take with fruit juices (see Directions) **Stop use and ask a doctor if** an allergic reaction to this product occurs. Seek medical help right away. **If pregnant or breast-feeding,** ask a health professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. ▶

Directions ■ **adults and children 12 years of age and over** take one 180 mg tablet with water once a day; do not take more than 1 tablet in 24 hours ■ **children under 12 years of age** do not use ■ **adults 65 years of age and older** ask a doctor ■ **consumers with kidney disease** ask a doctor ***Other information*** ■ safety sealed: do not use if carton was opened or if printed foil inner seal on bottle is torn or missing ■ store between 20° and 25°C (68° and 77°F) ■ protect from excessive moisture ***Inactive ingredients*** colloidal silicon dioxide, croscarmellose sodium, hypromellose, iron oxide blends, magnesium stearate, microcrystalline cellulose, polyethylene glycol, povidone, pregelatinized starch, titanium dioxide ***Questions or comments?*** call toll-free **1-866-844-2798** or www.allegra.com

Dist. By: Chattem, Inc., a Sanofi Company
Chattanooga, TN 37409-0219 ©2022
Origin India 858016

LOT
EXP

This label may not be the latest approved by FDA.
For current labeling information, please visit https://www.fda.gov/drugsatfda



This label may not be the latest approved by FDA.
For current labeling information, please visit https://www.fda.gov/drugsatfda

(b) (4)

(b) (4)

NDC 41167-4124-8 NON-DROWSY

Allegra® ALLERGY 24 HR

fexofenadine HCl tablet
180 mg/antihistamine

INDOOR / OUTDOOR ALLERGY RELIEF

✓ Sneezing ✓ Runny Nose
✓ Itchy, Watery Eyes
✓ Itchy Nose or Throat

100 TABLETS

Active ingredient (in each tablet) ***Purpose***
Fexofenadine HCl 180 mg.....Antihistamine

Uses temporarily relieves these symptoms due to hay fever or other upper respiratory allergies: ■ runny nose ■ sneezing ■ itchy, watery eyes ■ itching of the nose or throat

Warnings **Do not use** if you have ever had an allergic reaction to this product or any of its ingredients. **Ask a doctor before use if you have** kidney disease. Your doctor should determine if you need a different dose. **When using this product** ■ do not take more than directed ■ do not take at the same time as aluminum or magnesium antacids ■ do not take with fruit juices (see Directions) **Stop use and ask a doctor if** an allergic reaction to this product occurs. Seek medical help right away. **If pregnant or breast-feeding,** ask a health professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. ▶

Directions ■ **adults and children 12 years of age and over** take one 180 mg tablet with water once a day; do not take more than 1 tablet in 24 hours ■ **children under 12 years of age** do not use ■ **adults 65 years of age and older** ask a doctor ■ **consumers with kidney disease** ask a doctor ***Other information*** ■ safety sealed: do not use if carton was opened or if printed foil inner seal on bottle is torn or missing ■ store between 20° and 25°C (68° and 77°F) ■ protect from excessive moisture ***Inactive ingredients*** colloidal silicon dioxide, croscarmellose sodium, hypromellose, iron oxide blends, magnesium stearate, microcrystalline cellulose, polyethylene glycol, povidone, pregelatinized starch, titanium dioxide ***Questions or comments?*** call toll-free **1-866-844-2798** or www.allegra.com

Dist. By: Chattem, Inc., a Sanofi Company
Chattanooga, TN 37409-0219 ©2022
Origin India 858017

LOT
EXP

Reference ID: 5218613

Drug Facts

Active ingredient (in each tablet) **Purpose**
Fexofenadine HCl 180 mg ..... Antihistamine

Uses
temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:
- runny nose
- sneezing
- itchy, watery eyes
- itching of the nose or throat

Warnings
Do not use if you have ever had an allergic reaction to this product or any of its ingredients.

Ask a doctor before use if you have kidney disease.
Your doctor should determine if you need a different dose.

When using this product
- do not take more than directed
- do not take at the same time as aluminum or magnesium antacids
- do not take with fruit juices (see Directions)

Stop use and ask a doctor if an allergic reaction to this product occurs. Seek medical help right away.

If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. ▶

Drug Facts (continued)

Directions

adults and children 12 years of age and over	take one 180 mg tablet with water once a day; do not take more than 1 tablet in 24 hours
children under 12 years of age	do not use
adults 65 years of age and older	ask a doctor
consumers with kidney disease	ask a doctor

Other information
- safety sealed: do not use if printed foil inner seal on bottle is torn or missing
- store between 20° and 25°C (68° and 77°F)
- protect from excessive moisture

Inactive ingredients
colloidal silicon dioxide, croscarmellose sodium, hypromellose, iron oxide blends, magnesium stearate, microcrystalline cellulose, polyethylene glycol, povidone, pregelatinized starch, titanium dioxide

Questions or comments?
call toll-free **1-866-844-2798** or www.allegra.com

The makers of Allegra® do not make store brand products.
The trade dress of this Allegra® package is subject to trademark protection.
Dist. By: Chattem, Inc., a Sanofi Company
Chattanooga, TN 37409-0219 ©2022
Origin India

849975

LOT

EXP

0 41167 41243 5

Drug Facts

Active ingredient (in each tablet) **Purpose**
Fexofenadine HCl 180 mg ..... Antihistamine

Uses
temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:
- runny nose
- sneezing
- itchy, watery eyes
- itching of the nose or throat

Warnings
Do not use if you have ever had an allergic reaction to this product or any of its ingredients.

Ask a doctor before use if you have kidney disease.
Your doctor should determine if you need a different dose.

When using this product
- do not take more than directed
- do not take at the same time as aluminum or magnesium antacids
- do not take with fruit juices (see Directions)

Stop use and ask a doctor if an allergic reaction to this product occurs. Seek medical help right away.

If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. ▶

Drug Facts (continued)

Directions

adults and children 12 years of age and over	take one 180 mg tablet with water once a day; do not take more than 1 tablet in 24 hours
children under 12 years of age	do not use
adults 65 years of age and older	ask a doctor
consumers with kidney disease	ask a doctor

Other information
- safety sealed: do not use if printed foil inner seal on bottle is torn or missing
- store between 20° and 25°C (68° and 77°F)
- protect from excessive moisture

Inactive ingredients
colloidal silicon dioxide, croscarmellose sodium, hypromellose, iron oxide blends, magnesium stearate, microcrystalline cellulose, polyethylene glycol, povidone, pregelatinized starch, titanium dioxide

Questions or comments?
call toll-free **1-866-844-2798** or www.allegra.com

The makers of Allegra® do not make store brand products.
The trade dress of this Allegra® package is subject to trademark protection.
Dist. By: Chattem, Inc., a Sanofi Company
Chattanooga, TN 37409-0219 ©2022
Origin India

849975

LOT

EXP

0 41167 41243 5







This label may not be the latest approved by FDA.
For current labeling information, please visit https://www.fda.gov/drugsatfda

This label may not be the latest approved by FDA.
For current labeling information, please visit https://www.fda.gov/drugsatfda

Allegra® Hives 24 HR

NON-DROWSY

5 TABLETS

NDC 41167-4126-8

NON-DROWSY

Allegra® Hives

fexofenadine HCl tablet
180 mg/antihistamine

24 HR

REDUCES HIVES and RELIEVES ITCHING due to hives

5 TABLETS



Actual Size

849982

849982

849982

Drug Facts

Active ingredient (in each tablet)	Purpose
Fexofenadine HCl 180 mg	Antihistamine

Uses reduces hives and relieves itching due to hives (urticaria). This product will not prevent hives or an allergic skin reaction from occurring.

Warnings

Severe Allergy Warning: Get emergency help **immediately** if you have hives along with any of the following symptoms:

- trouble swallowing
- dizziness or loss of consciousness
- swelling of tongue
- swelling in or around mouth
- trouble speaking
- drooling
- wheezing or problems breathing

These symptoms may be signs of anaphylactic shock. This condition can be life threatening if not treated by a health professional **immediately**. Symptoms of anaphylactic shock may occur when hives first appear or up to a few hours later.

Not a Substitute for Epinephrine. If your doctor has prescribed an epinephrine injector for "anaphylaxis" or severe allergy symptoms that could occur with your hives, never use this product as a substitute for the epinephrine injector. If you have been prescribed an epinephrine injector, you should carry it with you at all times.

Do not use

- to **prevent** hives from any known cause such as:
  - foods
  - insect stings
  - medicines
  - latex or rubber gloves

  because this product will not stop hives from occurring. Avoiding the cause of your hives is the only way to prevent them. Hives can sometimes be serious. If you do not know the cause of your hives, see your doctor for a medical exam. Your doctor may be able to help you find a cause.
- if you have ever had an allergic reaction to this product or any of its ingredients

Drug Facts (continued)

- store between 20° and 25°C (68° and 77°F)
- protect from excessive moisture

Inactive ingredients colloidal silicon dioxide, croscarmellose sodium, hypromellose, iron oxide blends, magnesium stearate, microcrystalline cellulose, polyethylene glycol, povidone, pregelatinized starch, titanium dioxide

Drug Facts (continued)

Questions or comments? call toll-free **1-866-844-2798** or www.allegra.com

The makers of Allegra® do not make store brand products.
The trade dress of this Allegra® package is subject to trademark protection.
Dist. By: Chattem, Inc., a Sanofi Company, Chattanooga, TN 37409-0219
©2022 Origin India

Drug Facts (continued)

Ask a doctor before use if you have

- kidney disease. Your doctor should determine if you need a different dose.
- hives that are an unusual color, look bruised or blistered
- hives that do not itch

When using this product

- do not take more than directed
- do not take at the same time as aluminum or magnesium antacids
- do not take with fruit juices (see Directions)

Stop use and ask a doctor if

- an allergic reaction to this product occurs. Seek medical help right away.
- symptoms do not improve after 3 days of treatment
- the hives have lasted more than 6 weeks

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

adults and children 12 years of age and over	take one 180 mg tablet with water once a day; do not take more than 1 tablet in 24 hours
children under 12 years of age	do not use
adults 65 years of age and older	ask a doctor
consumers with kidney disease	ask a doctor

Other information

- safely sealed: do not use if carton is opened or if individual blister units are torn or opened

LOT

EXP

0 41167 41270 1

Allegra® Hives 24 HR

Reference ID: 5218613

This label may not be the latest approved by FDA.
For current labeling information, please visit https://www.fda.gov/drugsatfda

858013

Allegra® Hives 24 HR

NDC 41167-4126-8

NON-DROWSY

5 TABLETS

Allegra® Hives

fexofenadine HCl tablet
180 mg/antihistamine

24 HR

REDUCES HIVES and RELIEVES ITCHING due to hives

5 TABLETS

Actual Size

NON-DROWSY

858013

Drug Facts (continued)

Inactive ingredients colloidal silicon dioxide, croscarmellose sodium, hypromellose, iron oxide blends, magnesium stearate, microcrystalline cellulose, polyethylene glycol, povidone, pregelatinized starch, titanium dioxide

- store between 20° and 25°C (68° and 77°F)
- protect from excessive moisture

Drug Facts (continued)

Questions or comments? call toll-free **1-866-944-2798** or www.allegra.com

The makers of Allegra® do not make store brand products.
The trade dress of this Allegra® package is subject to trademark protection.
Dist. By: Chattem, Inc., a Sanofi Company, Chattanooga, TN 37409-0219
©2022 Origin India

Other information

- safety sealed: do not use if carton is opened or if individual blister units are torn or opened

Directions

adults and children 12 years of age and over	take one 180 mg tablet with water once a day; do not take more than 1 tablet in 24 hours
children under 12 years of age	do not use
adults 65 years of age and older	ask a doctor
consumers with kidney disease	ask a doctor

Drug Facts (continued)

Ask a doctor before use if you have

- kidney disease. Your doctor should determine if you need a different dose.
- hives that are an unusual color, look bruised or blistered
- hives that do not itch

When using this product

- do not take more than directed
- do not take at the same time as aluminum or magnesium antacids
- do not take with fruit juices (see Directions)

Stop use and ask a doctor if

- an allergic reaction to this product occurs. Seek medical help right away.
- symptoms do not improve after 3 days of treatment
- the hives have lasted more than 6 weeks

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Drug Facts

Active ingredient (in each tablet) **Purpose**
Fexofenadine HCl 180 mg ..... Antihistamine

Uses reduces hives and relieves itching due to hives (urticaria). This product will not prevent hives or an allergic skin reaction from occurring.

Warnings

Severe Allergy Warning: Get emergency help **immediately** if you have hives along with any of the following symptoms:

- trouble swallowing
- dizziness or loss of consciousness
- swelling of tongue
- swelling in or around mouth
- drooling
- wheezing or problems breathing
- trouble speaking

These symptoms may be signs of anaphylactic shock. This condition can be life threatening if not treated by a health professional **immediately.** Symptoms of anaphylactic shock may occur when hives first appear or up to a few hours later.

Not a Substitute for Epinephrine. If your doctor has prescribed an epinephrine injector for "anaphylaxis" or severe allergy symptoms that could occur with your hives, never use this product as a substitute for the epinephrine injector. If you have been prescribed an epinephrine injector, you should carry it with you at all times.

Do not use

- **to prevent** hives from any known cause such as:
  - foods
  - insect stings
  - medicines
  - latex or rubber gloves

  because this product will not stop hives from occurring. Avoiding the cause of your hives is the only way to prevent them. Hives can sometimes be serious. If you do not know the cause of your hives, see your doctor for a medical exam. Your doctor may be able to help you find a cause.
- if you have ever had an allergic reaction to this product or any of its ingredients

Allegra® Hives 24 HR

SAMPLE:
NOT FOR RETAIL SALE

LOT
EXP

858013

(b) (4)

Sanofi Regulatory Specifications — Verified Date: 08.15.22

Drug Facts Info

Drug Facts (Title)	Font name: Helvetica Neue 77 Bold Condensed Oblique	10 point type
Headings	Font name: Helvetica Neue 77 Bold Condensed Oblique	8 point type
Subheadings	Font name: Helvetica Neue 77 Bold Condensed	6 point type
Body Text	Font name: Helvetica Neue 57 Condensed	6 point type
Bullets	Font name: Universal News w. Commercial Pi 97	5 point type
Bullets on same lines: end of statement separated from bulleted statement by two ems		YES
Spacing of the hairlines from edge of box – i.e. Minimum of 1 space either side of Drug Facts Box		YES

Tracking:	-16 pt. to -18 pt.	Horizontal Scale:	84%
Leading (Minimum space in body of Drug Facts):	6.5 pt.	Maximum Characters/Inch:	39
Barlines:	2.5 pt.	Hairlines:	0.5 pt.

Principal Display Panel Info

Font size of Net Quantity of Contents (Smallest character height in inches)	0.20 in
PDP dimensions (in millimeters)	H 75.00 mm W 110.00 mm
PDP dimensions (in square inches)	H 2.95 in W 4.33 in (12.77 sq in)
Font size of Statement of Identity (If not live text, to be measured in Helvetica capital "M")	14 pt
Font size of Logo/Largest Copy on PDP (If not live text, to be measured in Helvetica capital "M")	46 pt
Ratio of Statement of Identity to Logo/Largest Copy on PDP	30%

Reference ID: 5218613

This label may not be the latest approved by FDA.
For current labeling information, please visit https://www.fda.gov/drugsatfda

(b) (4)



Sanofi Regulatory Specifications — Verified Date: 07/27/22

Drug Facts Info

Drug Facts (Title)	Font name: Helvetica Neue 77 Bold Condensed Oblique	10 point type
Drug Facts (continued)	Font name: Helvetica Neue 77 Bold Condensed Oblique / Helvetica Neue 57 Condensed	8 point type
Headings	Font name: Helvetica Neue 77 Bold Condensed Oblique	8 point type
Subheadings	Font name: Helvetica Neue 77 Bold Condensed	6 point type
Body Text	Font name: Helvetica Neue 57 Condensed	6 point type
Bullets	Font name: Universal News w. Commercial Pi 97	5 point type
Bullets on same lines: end of statement separated from bulleted statement by two ems		YES
Spacing of the hairlines from edge of box – i.e. Minimum of 1 space either side of Drug Facts Box		YES

Tracking:	-10 pt.	Horizontal Scale:	98%-100%
Leading (Minimum space in body of Drug Facts):	6.5 pt.	Maximum Characters/Inch:	34
Barlines:	2.5 pt.	Hairlines:	0.5 pt.

Principal Display Panel Info

Font size of Net Quantity of Contents (Smallest character height in inches)	.20 in
PDP dimensions (in millimeters)	H 109.99 mm W 75.03 mm
PDP dimensions (in square inches)	H 4.33 in W 2.96 in (12.824 sq in)
Font size of Statement of Identity (If not live text, to be measured in Helvetica capital "M")	13.9 pt
Font size of Logo/Largest Copy on PDP (If not live text, to be measured in Helvetica capital "M")	42.25 pt
Ratio of Statement of Identity to Logo/Largest Copy on PDP Statement of Identity (pt.) (Divided by) Largest Logo Copy (pt.)	30%

Reference ID: 5218613

Reference ID: 5218613



This label may not be the latest approved by FDA.
For current labeling information, please visit https://www.fda.gov/drugsatfda

This label may not be the latest approved by FDA.
For current labeling information, please visit https://www.fda.gov/drugsatfda

849980

NON-DROWSY
Allegra® Hives 24HR
30 TABLETS

Drug Facts
Active ingredient (in each tablet) *Purpose*
Fexofenadine HCl 180 mg ..... Antihistamine

Uses
reduces hives and relieves itching due to hives (urticaria).
This product will not prevent hives or an allergic skin reaction from occurring. ▲

NDC 41167-4126-4

NON-DROWSY

Allegra® Hives

fexofenadine HCl tablet
180 mg/antihistamine

24 HR

REDUCES HIVES and
RELIEVES ITCHING due to hives

30 TABLETS

Actual Size

LOT

EXP

0 41167 41264 0

Allegra® Hives 24HR

Drug Facts (continued)

Other information
- safety sealed: do not use if carton is opened or if printed foil inner seal on bottle is torn or missing
- store between 20° and 25°C (68° and 77°F)
- protect from excessive moisture

Inactive ingredients
colloidal silicon dioxide, croscarmellose sodium, hypromellose, iron oxide blends, magnesium stearate, microcrystalline cellulose, polyethylene glycol, povidone, pregelatinized starch, titanium dioxide

Questions or comments?
call toll-free **1-866-844-2798** or www.allegra.com

The makers of Allegra® do not make store brand products. The trade dress of this Allegra® package is subject to trademark protection.

Dist. By: Chattem, Inc., a Sanofi Company, Chattanooga, TN 37409-0219
©2022 Origin India

849980

Drug Facts (continued)

Warnings
Severe Allergy Warning: Get emergency help **immediately** if you have hives along with any of the following symptoms:
- trouble swallowing
- dizziness or loss of consciousness
- swelling of tongue
- swelling in or around mouth
- trouble speaking
- wheezing or problems breathing
- drooling

These symptoms may be signs of anaphylactic shock. This condition can be life threatening if not treated by a health professional **immediately**. Symptoms of anaphylactic shock may occur when hives first appear or up to a few hours later.
Not a Substitute for Epinephrine. If your doctor has prescribed an epinephrine injector for "anaphylaxis" or severe allergy symptoms that could occur with your hives, never use this product as a substitute for the epinephrine injector. If you have been prescribed an epinephrine injector, you should carry it with you at all times.

Do not use
- to **prevent** hives from any known cause such as:
  - foods
  - insect stings
  - medicines
  - latex or rubber gloves

  because this product will not stop hives from occurring. Avoiding the cause of your hives is the only way to prevent them. Hives can sometimes be serious. If you do not know the cause of your hives, see your doctor for a medical exam. Your doctor may be able to help you find a cause.
- if you have ever had an allergic reaction to this product or any of its ingredients. ▶

Drug Facts (continued)

Ask a doctor before use if you have
- kidney disease. Your doctor should determine if you need a different dose.
- hives that are an unusual color, look bruised or blistered
- hives that do not itch

When using this product
- do not take more than directed
- do not take at the same time as aluminum or magnesium antacids
- do not take with fruit juices (see Directions)

Stop use and ask a doctor if
- an allergic reaction to this product occurs. Seek medical help right away.
- symptoms do not improve after 3 days of treatment
- the hives have lasted more than 6 weeks

If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

adults and children 12 years of age and over	take one 180 mg tablet with water once a day; do not take more than 1 tablet in 24 hours
children under 12 years of age	do not use
adults 65 years of age and older	ask a doctor
consumers with kidney disease	ask a doctor ▶

849980

(b) (4)

This label may not be the latest approved by FDA.
For current labeling information, please visit https://www.fda.gov/drugsatfda

(b) (4)







This label may not be the latest approved by FDA.
For current labeling information, please visit https://www.fda.gov/drugsatfda

849978

NON-DROWSY
Allegra® Hives 24 HR
60 TABLETS

Drug Facts
Active ingredient (in each tablet) Purpose
Fexofenadine HCl 180 mg ..... Antihistamine

Uses
reduces hives and relieves itching due to hives (urticaria).
This product will not prevent hives or an allergic skin reaction from occurring. ▲

NDC 41167-4126-1

NON-DROWSY

Allegra® Hives

fexofenadine HCl tablet
180 mg/antihistamine

24 HR

REDUCES HIVES and
RELIEVES ITCHING due to hives

60 TABLETS

Actual Size

LOT

EXP

0 41167 41261 9

Allegra® Hives 24 HR

Drug Facts (continued)

Other information
- safety sealed: do not use if carton is opened or if printed foil inner seal on bottle is torn or missing
- store between 20° and 25°C (68° and 77°F)
- protect from excessive moisture

Inactive ingredients
colloidal silicon dioxide, croscarmellose sodium, hypromellose, iron oxide blends, magnesium stearate, microcrystalline cellulose, polyethylene glycol, povidone, pregelatinized starch, titanium dioxide

Questions or comments?
call toll-free **1-866-844-2798** or www.allegra.com

The makers of Allegra® do not make store brand products. The trade dress of this Allegra® package is subject to trademark protection.

Dist. By: Chattem, Inc., a Sanofi Company, Chattanooga, TN 37409-0219
©2022 Origin India

849978

Drug Facts (continued)

Warnings
Severe Allergy Warning: Get emergency help **immediately** if you have hives along with any of the following symptoms:
- trouble swallowing
- dizziness or loss of consciousness
- swelling of tongue
- swelling in or around mouth
- trouble speaking
- wheezing or problems breathing
- drooling

These symptoms may be signs of anaphylactic shock. This condition can be life threatening if not treated by a health professional **immediately**. Symptoms of anaphylactic shock may occur when hives first appear or up to a few hours later.
Not a Substitute for Epinephrine. If your doctor has prescribed an epinephrine injector for "anaphylaxis" or severe allergy symptoms that could occur with your hives, never use this product as a substitute for the epinephrine injector. If you have been prescribed an epinephrine injector, you should carry it with you at all times.

Do not use
- to **prevent** hives from any known cause such as:
  - foods
  - insect stings
  - medicines
  - latex or rubber gloves

because this product will not stop hives from occurring. Avoiding the cause of your hives is the only way to prevent them. Hives can sometimes be serious. If you do not know the cause of your hives, see your doctor for a medical exam.Your doctor may be able to help you find a cause.
- if you have ever had an allergic reaction to this product or any of its ingredients. ▶

Drug Facts (continued)

Ask a doctor before use if you have
- kidney disease. Your doctor should determine if you need a different dose.
- hives that are an unusual color, look bruised or blistered
- hives that do not itch

When using this product
- do not take more than directed
- do not take at the same time as aluminum or magnesium antacids
- do not take with fruit juices (see Directions)

Stop use and ask a doctor if
- an allergic reaction to this product occurs. Seek medical help right away.
- symptoms do not improve after 3 days of treatment
- the hives have lasted more than 6 weeks

If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

adults and children 12 years of age and over	take one 180 mg tablet with water once a day; do not take more than 1 tablet in 24 hours
children under 12 years of age	do not use
adults 65 years of age and older	ask a doctor
consumers with kidney disease	ask a doctor ▶

849978

(b) (4)

This label may not be the latest approved by FDA.
For current labeling information, please visit https://www.fda.gov/drugsatfda



Active ingredient (in each tablet) ***Purpose***
Fexofenadine HCl 180 mg.....Antihistamine

Uses reduces hives and relieves itching due to hives (urticaria). This product will not prevent hives or an allergic skin reaction from occurring.

Warnings **Severe Allergy Warning:** Get emergency help **immediately** if you have hives along with any of the following symptoms: ■ trouble swallowing ■ dizziness or loss of consciousness ■ drooling ■ swelling of tongue ■ swelling in or around mouth ■ trouble speaking ■ wheezing or problems breathing ▶

NDC 41167-4126-1

NON-DROWSY
Allegra®
Hives
fexofenadine HCl tablet 24 HR
180 mg/antihistamine
60 TABLETS

Dist. By: Chattem, Inc.,
a Sanofi Company
Chattanooga, TN 37409
©2022
Origin India

Peel Here

849979

These symptoms may be signs of anaphylactic shock. This condition can be life threatening if not treated by a health professional **immediately**. Symptoms of anaphylactic shock may occur when hives first appear or up to a few hours later. **Not a Substitute for Epinephrine.** If your doctor has prescribed an epinephrine injector for "anaphylaxis" or severe allergy symptoms that could occur with your hives, never use this product as a substitute for the epinephrine injector. If you have been prescribed an epinephrine injector, you should carry it with you at all times. **Do not use** ■ to **prevent** hives from any known cause such as: ■ foods ■ insect stings ■ medicines ■ latex or rubber gloves because this product will not stop hives from occurring. Avoiding the cause of your hives is the only way to prevent them. Hives can sometimes be serious. If you do not know the ▶

cause of your hives, see your doctor for a medical exam. Your doctor may be able to help you find a cause. ■ if you have ever had an allergic reaction to this product or any of its ingredients **Ask a doctor before use if you have** ■ kidney disease. Your doctor should determine if you need a different dose. ■ hives that are an unusual color, look bruised or blistered ■ hives that do not itch **When using this product** ■ do not take more than directed ■ do not take at the same time as aluminum or magnesium antacids ■ do not take with fruit juices (see Directions) **Stop use and ask a doctor if** ■ an allergic reaction to this product occurs. Seek medical help right away. ■ symptoms do not improve after 3 days of treatment ■ the hives have lasted ▶

more than 6 weeks **If pregnant or breast-feeding**, ask a health professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. ***Directions*** ■ **adults and children 12 years of age and over** take one 180 mg tablet with water once a day; do not take more than 1 tablet in 24 hours ■ **children under 12 years of age** do not use ■ **adults 65 years of age and older** ask a doctor ■ **consumers with kidney disease** ask a doctor ***Other information*** ■ safety sealed: do not use if carton was opened or if printed foil inner seal on bottle is torn or missing ■ store between 20° and 25°C (68° and 77°F) ■ protect from excessive moisture ***Inactive ingredients*** colloidal silicon dioxide, croscarmellose ▶

sodium, hypromellose, iron oxide blends, magnesium stearate, microcrystalline cellulose, polyethylene glycol, povidone, pregelatinized starch, titanium dioxide ***Questions or comments?*** call toll-free **1-866-844-2798** or www.allegra.com

Allegra® NDC 41167-4126-1
Hives
fexofenadine HCl tablet 24 HR
180 mg/antihistamine 60 TABLETS

Dist. By: Chattem, Inc., a Sanofi Company
Chattanooga, TN 37409-0219
©2022 Origin India

This label may not be the latest approved by FDA.
For current labeling information, please visit https://www.fda.gov/drugsatfda

---

This is a representation of an electronic record that was signed electronically. Following this are manifestations of any and all electronic signatures for this electronic record.

/s/

NUSHIN F TODD
08/01/2023 10:33:39 AM