

APPROVED

JUL 25 2000

0076112 LEX 01 00000000 048302

Abreva

Apply the ointment
to the product ulcer,
abrasion, and
burns.

Drug Facts

Active ingredient	Purpose
Docosanol 1.5%	Cold sores/ulcers and blisters

Use: • Apply to cold sores/ulcers/blisters on the lips or face.
• Shortens healing time and duration of symptoms.
• Reduces pain, burning, and/or itching.

Warnings
For external use only.
Do not use:
• If you are allergic to any ingredients in this product.
When using this product:
• Apply only to affected areas.
• Do not use in or near the eyes.
• Avoid applying directly to the mouth.
• Do not apply this product with lipsticks. This may conceal infection.
Keep out of reach of children.
• Use only on the lips and around the mouth.
• If you feel pain, redness or the sore does not heal, stop use within 10 days.
Keep out of reach of children. If swallowed, get medical help or contact a poison control center right away.

Directions
• Adults and children 12 years of age and over:
• Wash hands before and after applying ointment.
• Apply to affected areas on face or lips at the first sign of cold sores/ulcers/blisters. Early treatment ensures the best results.
• Use as directed.
• Use 5 times a day until healed.
• Children under 12 years of age: Do not use.

Other information • Store at 20°-25°C (68°-77°F). • Do not freeze.
Inactive ingredients • Polyethylene glycol, polyethylene glycol, polyethylene glycol, polyethylene glycol, polyethylene glycol.
Contains • See 1-877-766-0520 website.

tingling, pain, burning, and/or itching.
(this is a sub header under "shortens healing time and duration of symptoms")



Distributed by Bristol-Myers Squibb
Pharmaceutical Research Corporation, New York, NY
© 2000 Bristol-Myers Squibb

Abreva

Abreva

Abreva

Abreva

NEW Cold Sore Treatment

Shortens the healing time and duration of symptoms.
• Reduces pain, burning, and itching.

Apply to cold sores/ulcers/blisters on the lips or face.
• Shortens healing time and duration of symptoms.
• Reduces pain, burning, and/or itching.

Use only on the lips and around the mouth.
• If you feel pain, redness or the sore does not heal, stop use within 10 days.
Keep out of reach of children. If swallowed, get medical help or contact a poison control center right away.

Use as directed.
• Use 5 times a day until healed.
• Children under 12 years of age: Do not use.

Apply to affected areas on face or lips at the first sign of cold sores/ulcers/blisters. Early treatment ensures the best results.
• Use as directed.
• Use 5 times a day until healed.
• Children under 12 years of age: Do not use.

Wash hands before and after applying ointment.
• Apply to affected areas on face or lips at the first sign of cold sores/ulcers/blisters. Early treatment ensures the best results.
• Use as directed.
• Use 5 times a day until healed.
• Children under 12 years of age: Do not use.

Do not use:
• If you are allergic to any ingredients in this product.

When using this product:
• Apply only to affected areas.
• Do not use in or near the eyes.
• Avoid applying directly to the mouth.
• Do not apply this product with lipsticks. This may conceal infection.

Keep out of reach of children.
• Use only on the lips and around the mouth.
• If you feel pain, redness or the sore does not heal, stop use within 10 days.
Keep out of reach of children. If swallowed, get medical help or contact a poison control center right away.

For external use only.
Do not use:
• If you are allergic to any ingredients in this product.

Use: • Apply to cold sores/ulcers/blisters on the lips or face.
• Shortens healing time and duration of symptoms.
• Reduces pain, burning, and/or itching.

Drug Facts

Active ingredient	Purpose
Docosanol 1.5%	Cold sores/ulcers and blisters

Use: • Apply to cold sores/ulcers/blisters on the lips or face.
• Shortens healing time and duration of symptoms.
• Reduces pain, burning, and/or itching.

Warnings

For external use only.
Do not use:
• If you are allergic to any ingredients in this product.

When using this product:
• Apply only to affected areas.
• Do not use in or near the eyes.
• Avoid applying directly to the mouth.
• Do not apply this product with lipsticks. This may conceal infection.

Keep out of reach of children.
• Use only on the lips and around the mouth.
• If you feel pain, redness or the sore does not heal, stop use within 10 days.
Keep out of reach of children. If swallowed, get medical help or contact a poison control center right away.

Directions

• Adults and children 12 years of age and over:
• Wash hands before and after applying ointment.
• Apply to affected areas on face or lips at the first sign of cold sores/ulcers/blisters. Early treatment ensures the best results.
• Use as directed.
• Use 5 times a day until healed.
• Children under 12 years of age: Do not use.

Other information

• Store at 20°-25°C (68°-77°F). • Do not freeze.

Inactive ingredients

• Polyethylene glycol, polyethylene glycol, polyethylene glycol, polyethylene glycol, polyethylene glycol.

Contains

• See 1-877-766-0520 website.

APPROVED

JUL 25 2000

Abreva (Doxycycline Hyclate) Cream 2.0% 7.14.00

200%

Abreva
Doxycycline Hyclate Cream
NET WT 2.0g (0.7 oz)

Retain outer carton for full product uses, directions and warnings.

Distributed for Avair Pharmaceuticals by
SmithKline Beecham
Consumer Healthcare, LP
Pittsburgh, PA 15230, Made in the U.S.A.
© 2000 SmithKline Beecham

Abreva
Doxycycline Hyclate Cream
NET WT 2.0g (0.7 oz)

Retain outer carton for full product uses, directions and warnings.

Distributed for Avair Pharmaceuticals by
SmithKline Beecham
Consumer Healthcare, LP
Pittsburgh, PA 15230, Made in the U.S.A.
© 2000 SmithKline Beecham