



Visit us at www.abreva.com
 If you are not completely satisfied, save receipt and remaining product, and call us at 1-877-709-3539 weekdays.
 Abreva and logo design and overall design elements are trademarks owned by the GSK group of companies.
 Distributed for Avantir Pharmaceuticals by
GlaxoSmithKline Consumer Healthcare, L.P.
 Moon Township, PA 15108, Made in Germany

Drug Facts
 Press Pump completely about 5-8 times to get first dose.
 DOCOSANOL 10% CREAM

Drug Facts (continued)

Keep out of reach of children if swallowed, get medical help or contact a poison control center right away

Purpose
 Docosanol 10%.....Cold sore/fever blister treatment

Uses
 • treats cold sores/fever blisters on the face or lips
 • shortens healing time and duration of symptoms;

Warnings
 This product may cause a severe allergic reaction. Symptoms may include:
 • hives • facial swelling
 • wheezing/difficulty breathing • shock • rash
 If an allergic reaction occurs, stop use and seek medical help right away.

Other information
 • store at 20-25°C (68-77°F)
 • do not freeze
 • do not reuse

When using this product
 • apply only to affected areas • do not use in or near the eyes
 • avoid applying directly inside your mouth
 • do not share this product with anyone. This may spread infection.

Do not use • if you are allergic to any ingredient in this product

For external use only

Ingredients
 Inactive: benzyl alcohol, light mineral oil, propylene glycol, purified water, sucrose disaccharide, sucrose stearate

Questions or comments?
 call toll-free 1-877-709-3539 (English/Spanish) weekdays

Lot 104396XA
Exp ©2015 GSK

abreva
 DOCOSANOL 10% CREAM
 COLD SORE/FEVER BLISTER TREATMENT

Thank you for purchasing ABREVA. We hope that since ABREVA has been demonstrated to shorten the healing time of cold sores, it helps you look and feel better faster. Please follow insert directions for proper use and refer to the information below.

You are not alone
 An estimated one-in-five Americans is affected by cold sores each year. Most people who experience them get two to three cold sore outbreaks each year. Just before and during an outbreak, the cold sore infection can be easily passed on to others. It is especially important to be aware of early warning signs that a cold sore is coming, and to be sure to avoid close physical contact when you have one.

Getting to the Root of the Problem
 A blister, or cluster of blisters around your mouth, is caused by the cold sore virus. After the cold sore has healed, the virus stays dormant in your body but can be reactivated by everyday triggers like stress, sunlight, extreme hot or cold weather, colds and flu, lack of sleep, or menstruation.

The Sooner, the Better
 A typical cold sore outbreak can last as long as seven to ten days. The sooner you begin using ABREVA, the more effective the medicine will be. So for best results, use ABREVA at the very start of the cold sore – when you feel the first sign of the tingle, redness, bump or itch.

ABREVA is different from other over-the-counter cold sore products. Only ABREVA contains 10% docosanol, which helps to shorten the healing time of cold sores and the duration of symptoms including tingling, pain, itching and burning.

How to use your ABREVA Pump

- For Adults and Children 12 years and over
- ABREVA should be used five times a day until the cold sore is healed. Apply it directly to the cold sore at the first sign of a tingle, redness, bump or itch for best results.
- If you wish, you can apply cosmetics such as lipstick over ABREVA. However, use a separate applicator, like a cotton swab, to apply cosmetics over an unhealed cold sore to avoid spreading the infection. For best results, remove any cosmetics prior to applying ABREVA.

STEP 1 - Wash your hands before and after treating the cold sore.
STEP 2 - Remove cap and press pump firmly. When first used, expect to press pump completely about 6-8 times before obtaining first dose.
STEP 3 - Rub in gently but completely 5 times a day until healed.
STEP 4 - Place cap back on pump after each use.

For more information: Call us at 1-877-709-3539 weekdays or visit our web site www.abreva.com.

abreva
 DOCOSANOL 10% CREAM
 COLD SORE/FEVER BLISTER TREATMENT

(b) (4)

*Abreva contains the only OTC medicine approved by the FDA to shorten healing time and duration of symptoms.

We value your feedback
www.abreva.com

Distributed for Avamir Pharmaceuticals by
 GlaxoSmithKline Consumer Healthcare, L.P.
 Moorestown, PA 19380
 Made in Germany
 ©2015 GSK

Press Pump completely about 6-8 times to get first dose.

Drug Facts	Retain this insert for full product uses, directions and warnings.
Active ingredient	Purpose Docosanol 10% Cold sore/fever blister treatment
Uses	<ul style="list-style-type: none"> treats cold sores/fever blisters on the face or lips shortens healing time and duration of symptoms: tingling, pain, burning, and/or itching
Warnings	
Allergy alert	This product may cause a severe allergic reaction. Symptoms may include:
	<ul style="list-style-type: none"> hives facial swelling wheezing/difficulty breathing shock rash If an allergic reaction occurs, stop use and seek medical help right away.
For external use only	Do not use • if you are allergic to any ingredient in this product
When using this product	<ul style="list-style-type: none"> apply only to affected areas do not use in or near the eyes avoid applying directly inside your mouth do not share this product with anyone. This may spread infection.
Stop use and ask a doctor if	<ul style="list-style-type: none"> your cold sore gets worse or the cold sore is not healed within 10 days
Keep out of reach of children	If swallowed, get medical help or contact a poison control center right away
Directions	
	<ul style="list-style-type: none"> adults and children 12 years or over: <ul style="list-style-type: none"> wash hands before and after applying cream apply to affected area on face or lips at the first sign of cold sore/fever blister (tingle). early treatment ensures the best results rub in gently but completely use 5 times a day until healed children under 12 years: ask a doctor
Other information	store at 20°-25°C (68°-77°F) • do not freeze
Inactive ingredients	benzyl alcohol, light mineral oil, propylene glycol, purified water, sucrose distearate, sucrose stearate
Questions or comments?	call toll-free 1-877-709-3539 (English/Spanish) weekdays
	ABREVA and ABREVA logo are registered trademarks of the GSK group of companies.

Barcode: 3 0766-0811-64 0

104417XA

Lot Exp

NDC 0135-0200-03

gsk

abreva
 DOCOSANOL 10% CREAM
 Cold Sore/Fever Blister Treatment

Only FDA Approved Medicine to Shorten Healing Time*

Portable Convenient Pump

Cold Sore Treatment
 NET WT 2g (.07 oz)

(b) (4)



We value your feedback
www.abreva.com

***Abreva contains the only OTC medicine approved by the FDA to shorten healing time and duration of symptoms**
Retain this insert for full product uses, directions and warnings.

abreva
DOCOSANOL 10% CREAM
 COLD SORE/FEVER-BLISTER TREATMENT

Drug Facts	
Active ingredient	Purpose
Docosanol 10%.....	Cold sore/fever blister treatment
Uses	Drug Facts (continued)
<ul style="list-style-type: none"> treats cold sores/fever blisters on the face or lips shortens healing time and duration of symptoms: <ul style="list-style-type: none"> tingling, pain, burning, and/or itching 	contact a poison control center right away Directions <ul style="list-style-type: none"> adults and children 12 years or over: <ul style="list-style-type: none"> wash hands before and after applying cream apply to affected area on face or lips at the first sign of cold sore/fever blister (tingle). early treatment ensures the best results rub in gently but completely use 5 times a day until healed children under 12 years: ask a doctor
Warnings	Other information
Allergy alert This product may cause a severe allergic reaction. Symptoms may include: <ul style="list-style-type: none"> hives facial swelling wheezing/difficulty breathing shock rash If an allergic reaction occurs, stop use and seek medical help right away.	• store at 20°-25°C (68°-77°F) • do not freeze
For external use only	Inactive ingredients
Do not use • if you are allergic to any ingredient in this product When using this product <ul style="list-style-type: none"> apply only to affected areas do not use in or near the eyes avoid applying directly inside your mouth do not share this product with anyone. This may spread infection. 	benzyl alcohol, light mineral oil, propylene glycol, purified water, sucrose distearate, sucrose stearate
Stop use and ask a doctor if <ul style="list-style-type: none"> your cold sore gets worse or the cold sore is not healed within 10 days 	Questions or comments?
Keep out of reach of children If swallowed, get medical help or	call toll-free 1-877-709-3539 (English/Spanish) weekdays

If you are not completely satisfied, save receipt and remaining product, and call us at 1-877-709-3539 weekdays.
 Distributed for **Avanir Pharmaceuticals** by **GlaxoSmithKline** Consumer Healthcare, L.P.
 Moon Twp, PA 15108, Made in Germany 104420XA
 ©2015 GSK

Abreva and logo are trademarks of the GSK group of companies.

3 0766-0801-35 1

LOT
 EXP

NDC 0135-0200-04

gsk

abreva

DOCOSANOL 10% CREAM
Cold Sore/Fever Blister Treatment

Only FDA Approved Medicine
to Shorten Healing Time*

Cold Sore Treatment



104391XA

 2 TUBES of 2g (0.07 oz) | Total NET WT 4g (0.14 oz)

abreva®

DOCOSANOL 10% CREAM
 Cold Sore/Fever Blister Treatment

***Abreva contains the only OTC medicine approved by the FDA to shorten healing time and duration of symptoms**
Retain this insert for full product uses, directions and warnings.

Drug Facts	
Active ingredient	Purpose
Docosanol 10%.....	Cold sore/fever blister treatment
Uses	
<ul style="list-style-type: none"> • treats cold sores/fever blisters on the face or lips • shortens healing time and duration of symptoms: <ul style="list-style-type: none"> • tingling, pain, burning, and/or itching 	
Warnings	
Allergy alert	
This product may cause a severe allergic reaction. Symptoms may include:	
<ul style="list-style-type: none"> • hives • facial swelling • wheezing/difficulty breathing • shock • rash 	
If an allergic reaction occurs, stop use and seek medical help right away.	
For external use only	
Do not use • if you are allergic to any ingredient in this product	
When using this product	
<ul style="list-style-type: none"> • apply only to affected areas • do not use in or near the eyes • avoid applying directly inside your mouth • do not share this product with anyone. This may spread infection. 	
Stop use and ask a doctor if	
• your cold sore gets worse or the cold sore is not healed within 10 days	
Keep out of reach of children	
If swallowed, get medical help or contact a poison control center right away	
Directions	
<ul style="list-style-type: none"> • adults and children 12 years or over: <ul style="list-style-type: none"> • wash hands before and after applying cream • apply to affected area on face or lips at the first sign of cold sore/fever blister (tingle). • early treatment ensures the best results • rub in gently but completely • use 5 times a day until healed • children under 12 years: ask a doctor 	
Other information	
<ul style="list-style-type: none"> • store at 20°-25°C (68°-77°F) • do not freeze 	
Inactive ingredients	
benzyl alcohol, light mineral oil, propylene glycol, purified water, sucrose distearate, sucrose stearate	
Questions or comments?	
call toll-free 1-877-709-3539 (English/Spanish) weekdays	

Thank you for purchasing ABREVA cold sore/fever blister treatment. We hope that since ABREVA cream has been demonstrated to shorten the healing time of cold sores, it helps you look and feel better faster. Please follow insert directions for proper use and refer to the information below.

You are not alone
 An estimated one-in-five Americans is affected by cold sores each year. Most people who experience them get two to three cold sore outbreaks each year. Just before and during an outbreak, the cold sore infection can be easily passed on to others. It is especially important to be aware of early warning signs that a cold sore is coming, and to be sure to avoid close physical contact when you have one.

Getting to the Root of the Problem
 A blister, or cluster of blisters around your mouth, is caused by the cold sore virus. After the cold sore has healed, the virus stays dormant in your body but can be reactivated by everyday triggers like stress, sunlight, extreme hot or cold weather, colds and flu, lack of sleep, or menstruation.

The Sooner, the Better
 A typical cold sore outbreak can last as long as seven to ten days. The sooner you begin using ABREVA cream, the more effective the medicine will be. So for best results, use ABREVA cream at the very start of the cold sore – when you feel the first sign of the tingle, redness, bump or itch.

ABREVA cream is different from other over-the-counter cold sore products. Only ABREVA cream contains 10% docosanol, which helps to shorten the healing time of cold sores and the duration of symptoms including tingling, pain, itching and burning. Abreva and logo design are trademarks of the GSK group of companies.

If you are not completely satisfied, save receipt and remaining product, and call us at 1-877-709-3539 weekdays.

Lot:
Exp:



We value your feedback
www.abreva.com

Distributed for **Avanir Pharmaceuticals** by
 GlaxoSmithKline Consumer Healthcare, L.P.
 Moon Township, PA 15108, Made in Germany
 ©2015 GSK 104392XA

(b) (4)

