


- NOTES:
- 1) ALL DIMENSIONS IN INCHES
 - 2) TOLERANCES UNLESS SPECIFIED
 - 3) DRAWING SHOWS PRINT SIZE UP
 - 4) NO PRINT AREA
 - 5) PART NUMBER AREA MUST USE DECIMAL POINT FOR PART NUMBER
 - 6) NO PRINT, NO COLOR & NO VARNISH AREA
 - 7) "LOT" AND "EXP" PRINT AREA
 - 8) "LOT" AND "EXP" PRINT AREA

SUG. NO.		LB-PS-849	
REVISED PART # 1/8 TO RIGHT	DRUG	Pepcid Complete	
ADDED TOLERANCES	PLAT	ARS	
UPDATED REVISION LEVEL	SIZE	11	
CHG LOWER LEFT CORNER	DATE	12/4/98	
INTL. NO.	DATE	REVISED	
APPROVED		APPROVED	

Johnson & Johnson O MEBCK
CONSUMER PHARMACEUTICALS CO.
CAMP HILL ROAD, FORT WASHINGTON, PA 19034

TITLE: BOOKLET LABEL FOR
FACT 150CC BOTTLE



FREE PEPCID COMPLETE Offer
(Print in offer. See details inside.)

+ =

The benefits of a acid control at Pepcid AC reduce your heartburn acid. **DO NOT USE IF PRINTED FOLIO INNER SEAL UNDER BOTTLE CAP IS BROKEN.**

Drug Facts
Active Ingredients (in each chewable tablet): Purpose
Famotidine 10 mg..... Acid reducer
Calcium carbonate 800 mg..... Antacid
Magnesium hydroxide 165 mg..... Antacid

Questions? 1-800-755-4008
Lift Tab for Warnings and Directions

EXCESS ACID: A BURNING PROBLEM
It is normal for the stomach to produce acid, especially after consuming food and beverages. However, acid in the stomach may move up into the wrong place (the esophagus), causing burning pain and discomfort that interfere with everyday activities.

Heartburn—Caused by acid in the esophagus
Burning pain/discomfort in esophagus
A valve-like muscle called the lower esophageal sphincter (LES) is relaxed in an open position
Acid moves up from stomach

TIPS FOR MANAGING HEARTBURN

- Do not lie flat or bend over soon after eating.
- Do not eat late at night, or just before bedtime.
- Certain foods or drinks are more likely to cause heartburn, such as rich, spicy, fatty, and fried foods, chocolate, caffeine, alcohol, and even some fruits and vegetables.
- Eat slowly and do not eat big meals.
- If you are overweight, lose weight.
- If you smoke, quit smoking.
- Raise the head of your bed.
- Wear loose fitting clothing around your stomach.

FREE PEPCID COMPLETE OFFER
Buy 4 PEPCID® COMPLETE packages 25's or larger and get **FREE PEPCID COMPLETE 15's**
(Mail-In Offer / Expires 9/1/2003)

Here's How You Get A Free PEPCID COMPLETE 15's Package
1. Please include 4 product packages (remove UPC bar code removed from the bottom of outer carton) from any PEPCID COMPLETE or PEPCID AC Package (25's or larger).
2. Complete this request form, place in a stamped envelope along with cash register receipts and 4 proof-of-purchases and mail to:
PEPCID COMPLETE 15's Package
Highland Park, MI 48234-7190
© Johnson & Johnson • MEBCK 2000
00000009

NAME _____ Mr. _____ Mrs. _____ Ms. _____
(please print)
ADDRESS _____
CITY _____ STATE _____ ZIP _____
APT.# _____ TELEPHONE # _____
e-mail address: _____
See back for additional information

3

Drug Facts (continued)

Other information

- read the directions and warnings before use
- keep this booklet. It contains important information.
- store at 25-30°C (77-85°F)
- protect from moisture

Inactive ingredients cellulose acetate, corn starch, croscarmellose, flavors, hydroxypropyl cellulose, hydroxypropyl methylcellulose, lactose, magnesium stearate, pregelatinized starch, red iron oxide, sodium lauryl sulfate, sugar


Questions or comments? call toll-free 1-800-755-4008

A NON-PRESCRIPTION STOMACH MEDICINE

PEPCID COMPLETE combines an acid reducer (famotidine) with antacids (calcium carbonate and magnesium hydroxide) to relieve heartburn in two different ways. Acid reducers decrease the production of new stomach acid, while antacids neutralize acid that is already in the stomach. The active ingredients in PEPCID COMPLETE have been used for years to treat acid-related problems in millions of people safely and effectively.

• Just 1 tablet relieves heartburn due to acid indigestion.

Shown actual size



6

1) How frequently do you use non-prescription antacids or stomach remedies?

A. 2 to 3 days per week or more B. 1 day per week

C. 2 to 3 days per month or less

2) How severe is a typical heartburn or indigestion episode?

A. very severe B. somewhat severe C. not very severe D. not at all severe

3) PEPCID COMPLETE is a combination of a fast-acting chewable antacid and a powerful acid controller like Zantac75® or Pepcid AC. Which products will you replace with PEPCID COMPLETE?

A. _____ B. _____

Please exclude me from all future PEPCID COMPLETE offers or information.

NOTE: Offer good only in U.S.A. and APO/FPO areas. This request form may not be mechanically reproduced. LIMITING PEPCID COMPLETE PER PERSON. See this or companion requests will be honored. Your offer rights may not be transferred or sold. Offer void where prohibited or taxed. Please allow 6 to 8 weeks for receipt of package. © Registered trademark of Merck & Co., Inc. *Registered trademark of Glaxo Wellcome Warner-Lambert, OTC GP.

2

Drug Facts (continued)

Use -relieves heartburn associated with acid indigestion and sour stomach

Warnings

Allergy alert: Do not use if you are allergic to famotidine or other acid reducers

Do not use - if you have trouble swallowing - with other famotidine products or acid reducers

Ask a doctor or pharmacist before use if you are presently taking a prescription drug. Antacids may interact with certain prescription drugs.


Stop use and ask a doctor if - stomach pain continues - you need to take this product for more than 14 days

If pregnant or breastfeeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- adults and children 12 years and over:
 - do not swallow tablet whole; chew completely
 - to relieve symptoms, chew 1 tablet before swallowing
 - do not use more than 2 chewable tablets in 24 hours
- children under 12 years: ask a doctor



Labeling Format Information:

Fonts: Univers condensed, bold and bold italic.

Drug Facts: 9 pt Leading: 6 pt


Header: 8 pt Bullets: 6 pt

Subheader: 8 pt Barlines: 2.5 pt

Body Text: 6 pt Hairlines: .5 pt

Drug Facts (continued): 9 pt

Horizontal Scale: 94% Average Kerning: -4



LOT: _____ EXP: _____

XXXXXXXXXX

NOTES:

- 1) ALL DIMENSIONS, IN INCHES.
- 2) TOLERANCES UNLESS SPECIFIED:
- 3) DRAWING SHOWN PRINT: THE UP
- 4) NO PRINT AREA
- 5) PART NUMBER AREA MUST USE
- 6) DCCR 8 FT. FRONT FOR PART NUMBER
- 7) WITH NO BOX AROUND PART NUMBER
- 8) UPC AREA
- 9) NO PRINT, NO COLOR & NO VARNISH AREA
- 10) LOT AND EXP. PRINT AREA

REV. NO.	REV. DATE	REV. BY	REV. REASON
1	4/19/00	ADD	ADDED PART NO AREA UNDER BASE LABEL DATA
2	1/19/00	ADD	ADDED TOLERANCES
3	2/26/99	CHG	CHG'D LOWER LEFT CORNER
4			
5			
6			
7			
8			
9			
10			
11			
12			
13			
14			
15			
16			
17			
18			
19			
20			
21			
22			
23			
24			
25			
26			
27			
28			
29			
30			
31			
32			
33			
34			
35			
36			
37			
38			
39			
40			
41			
42			
43			
44			
45			
46			
47			
48			
49			
50			
51			
52			
53			
54			
55			
56			
57			
58			
59			
60			
61			
62			
63			
64			
65			
66			
67			
68			
69			
70			
71			
72			
73			
74			
75			
76			
77			
78			
79			
80			
81			
82			
83			
84			
85			
86			
87			
88			
89			
90			
91			
92			
93			
94			
95			
96			
97			
98			
99			
100			

FREE PEPICID COMPLETE OFFER

Buy 4 PEPICID COMPLETE packages
25s or larger and get
FREE PEPICID COMPLETE 15s

Mail-In Offer / Expires 9/1/2003

NAME (Please print) _____
ADDRESS _____
CITY _____
STATE _____
TELEPHONE # _____
E-mail address: _____

1. Please include a proof-of-purchase (UPC bar code removed from the bottom of outer carton) from any PEPICID COMPLETE or PEPICID AC Package (25s or larger).

2. Complete this request form, place in a striped envelope along with cash register receipts and a proof-of-purchase and mail to:
PEPICID COMPLETE 15s Package
P.O. Box 3795
Highland Park, NJ 08520-0795
© Johnson & Johnson • MERCK 2000

See back for additional information

EXCESS ACID: A BURNING PROBLEM

It's normal for the stomach to produce acid, especially after consuming food and beverages. However, acid in the stomach may move up into the wrong place (the esophagus), causing burning pain and discomfort that interfere with everyday activities.

Heartburn—Caused by acid in the esophagus

Burning pain/discomfort in esophagus

A valve-like muscle called the lower esophageal sphincter (LES) is relaxed in an open position

Acid moves up from stomach

TIPS FOR MANAGING HEARTBURN

- Do not lie flat or bend over soon after eating.
- Do not eat late at night, or just before bedtime.
- Certain foods or drinks are more likely to cause heartburn, such as rich, spicy, fatty, and fried foods, chocolate, caffeine, alcohol, and even some fruits and vegetables.
- Eat slowly and do not eat big meals.
- If you are overweight, lose weight.
- If you smoke, quit smoking.
- Raise the head of your bed.
- Wear loose fitting clothing around your stomach.

NEW! **Peppid Complete** NEW! 889 26

WITH DUAL ACTION
Reduces AND Neutralizes Acid

25 Mint Flavored Chewable Tablets

+5 Tablets FREE!

Works with **Just One Tablet!**

Contains Calcium! Relieves Heartburn Due to Acid Indigestion

FREE PEPICID COMPLETE Offer

Mail in offer. See details inside.

DO NOT USE IF PRINTED POLYMER SEAL UNDER BOTTLE CAP IS BROKEN.

The benefits of Peppid Complete are enhanced when taken with Peppid AC.

Drug Facts

Active ingredients	Purpose
(in each chewable tablet)	
Famotidine 10 mg	Acid reducer
Calcium carbonate 800 mg	Antacid
Magnesium hydroxide 165 mg	Antacid

Use reduces heartburn associated with acid indigestion and sour stomach.

Questions? 1-800-755-4008

See the Back Panel for Warnings and Directions



Drug Facts (continued)

Warnings
Allergy alert: Do not use if you are allergic to famotidine or other acid reducers.
Do not use:
• if you have trouble swallowing • with other famotidine products or acid reducers
• if you have ever had a heartburn episode while taking a prescription drug. Antacids may interact with certain prescription drugs.
Ask a doctor or pharmacist before use if you are presently taking a prescription drug. Antacids may interact with certain prescription drugs.
Stop use and ask a doctor if:
• stomach pain continues • you need to take this product for more than 14 days
If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions
• adults and children 12 years and over:
• do not swallow whole; chew completely
• to relieve symptoms, chew 1 tablet before swallowing
• do not use more than 2 chewable tablets in 24 hours
• children under 12 years: ask a doctor


Other information
• read the directions and warnings before use • keep this booklet. It contains important information.
• store at 25-30°C (77-86°F) • protect from moisture

Inactive ingredients: cellulose acetate, corn starch, dextrates, flavors, hydroxypropyl cellulose, hydroxypropyl methylcellulose, lactose, magnesium stearate, pregelatinized starch, red iron oxide, sodium lauryl sulfate, sugar

Questions or comments? call toll-free 1-800-755-4008

ANTACID, NON-PRESCRIPTION
STOMACH MEDICINE

PEPCID COMPLETE combines an acid reducer (famotidine) with antacids (calcium carbonate and magnesium hydroxide) to relieve heartburn in two different ways:
Acid reducers decrease the production of new stomach acid; antacids neutralize acid that is already in the stomach. The active ingredients in PEPCID COMPLETE have been used for years to treat acid-related problems in millions of people safely and effectively.



Shown actual size
• Just 1 tablet relieves heartburn due to acid indigestion.

1) How frequently do you use non-prescription antacids or stomach remedies?
A. 2 to 3 days per week or more B. 1 day per week
C. 2 to 3 days per month or less

2) How severe is a typical heartburn or indigestion episode?
A. very severe B. somewhat severe
C. not very severe D. not at all severe

3) PEPCID COMPLETE is a combination of a fast-acting chewable antacid, and a powerful acid controller like Zantac75® or Pepcid AC. Which product will you replace with PEPCID COMPLETE?
A. _____
B. _____

Please exclude me from all future PEPCID COMPLETE offers or information.

NOTE: Offer good only in U.S.A. and APO/FPO addresses. This request form may not be mechanically reproduced. LIMIT TWO PEPCID COMPLETE OFFERS PER FAMILY AND/OR ADDRESS. No group or organization requests will be honored. Your offer rights may not be transferred or assigned. Offer void where prohibited or taxed. Please allow 6 to 8 weeks for receipt of package.



© Registered trademark of Merck & Co., Inc.
*Registered trademark of Glaxo Wellcome Warner-Lambert OTCGP

Labeling Format Information:

Fonts: Univers condensed, bold and bold italic.

Drug Facts: 9 pt Leading: 6 pt
Header: 8 pt Bullets: 6 pt
Subheader: 8 pt Barlines: 2.5 pt
Body Text: 6 pt Hairlines: .5 pt
Drug Facts (continued): 10 pt

Horizontal Scale: 94% Average Kerning: -4



QUESTIONS OR COMMENTS?

Please call 1-800-755-4008 for more information and a guide on how to help control heartburn.

© Registered trademark of Merck & Co., Inc.
Distributed by: **Johnson & Johnson • MERCK**
CONSUMER PHARMACEUTICALS CO.
FORT WASHINGTON, PA 19374 USA

FORM 1	DATE	2/26/99	APPROVED	DATE	2/26/99
FORM 2	DATE	2/26/99	APPROVED	DATE	2/26/99
FORM 3	DATE	2/26/99	APPROVED	DATE	2/26/99
FORM 4	DATE	2/26/99	APPROVED	DATE	2/26/99
FORM 5	DATE	2/26/99	APPROVED	DATE	2/26/99
FORM 6	DATE	2/26/99	APPROVED	DATE	2/26/99
FORM 7	DATE	2/26/99	APPROVED	DATE	2/26/99
FORM 8	DATE	2/26/99	APPROVED	DATE	2/26/99
FORM 9	DATE	2/26/99	APPROVED	DATE	2/26/99
FORM 10	DATE	2/26/99	APPROVED	DATE	2/26/99
FORM 11	DATE	2/26/99	APPROVED	DATE	2/26/99
FORM 12	DATE	2/26/99	APPROVED	DATE	2/26/99
FORM 13	DATE	2/26/99	APPROVED	DATE	2/26/99
FORM 14	DATE	2/26/99	APPROVED	DATE	2/26/99
FORM 15	DATE	2/26/99	APPROVED	DATE	2/26/99
FORM 16	DATE	2/26/99	APPROVED	DATE	2/26/99
FORM 17	DATE	2/26/99	APPROVED	DATE	2/26/99
FORM 18	DATE	2/26/99	APPROVED	DATE	2/26/99
FORM 19	DATE	2/26/99	APPROVED	DATE	2/26/99
FORM 20	DATE	2/26/99	APPROVED	DATE	2/26/99

Labeling Format Information:
 Fonts: Unifers condensed, bold and bold italic.
 Drug Facts: 9 pt
 Header: 8 pt
 Subheader: 8 pt
 Body Text: 6 pt
 Drug Facts (continued): 10 pt
 and bold italic.
 Leading: 6 pt
 Bullets: 6 pt
 Barlines: 2.5 pt
 Hairlines: .5 pt

PROVEN EFFECTIVE IN CLINICAL STUDIES
 In clinical studies, PEPCID COMPLETE was significantly better than placebo pills in relieving heartburn.

Onset of Relief
Percent of heartburn episodes relieved within 30 minutes

Treatment	Percent of heartburn episodes relieved within 30 minutes
PEPCID COMPLETE	45
Placebo	33
Famotidine	41
Antacid	0

Duration of Relief
Percent of heartburn episodes relieved for at least 7 hours

Treatment	Percent of heartburn episodes relieved for at least 7 hours
PEPCID COMPLETE	70
Placebo	59
Famotidine	61
Antacid	0

QUESTIONS OR COMMENTS?

Please call 1-800-755-4008 for more information and a guide on how to help control heartburn.

Registered trademark of Merck & Co., Inc.
 Distributed by: **Johnson & Johnson • MERCK**
 CONSUMER PHARMACEUTICALS CO.
 FORT WASHINGTON, PA 19034 USA

XXXXXX

LOT: _____
 EXP: _____

Johnson & Johnson - J&J

See back for additional information

Highland Park, NJ 08520-4179
c Johnson & Johnson • HEBECK 2900

PEPCID COMPLETE 15's, PO Box 3719

TELEPHONE # _____
e-mail address: _____

STATE _____
CITY _____
ZIP _____
APT # _____

NAME (please print) _____
ADDRESS _____
Mr. _____ Mrs. _____ Ms. _____

Mail-In Offer / Expires 9/1/2003

Here's How You Get A Free PEPCID COMPLETE 15's Package

1. Please include a proof-of-purchase (UPC bar code removed from any PEPCID COMPLETE or PEPCID AC Package (25's or larger)).

2. Complete this request form, place in a stamped envelope along with cash register receipts and 4 proof-of-purchases and mail to:

Buy 4 PEPCID COMPLETE packages
25's or larger and get
FREE PEPCID COMPLETE 15's

00000000

EXCESS ACID: A BURNING PROBLEM

It is normal for the stomach to produce acid, especially after consuming food and beverages. However, acid in the stomach may move up into the wrong place (the esophagus), causing burning pain and discomfort that interfere with everyday activities.

Heartburn—Caused by acid in the esophagus

Burning pain/discomfort in esophagus

A valve-like muscle called the lower esophageal sphincter (LES) is relaxed in an open position

Acid moves up from stomach

TIPS FOR MANAGING HEARTBURN

- Do not lie flat or bend over soon after eating.
- Do not eat late at night, or just before bedtime.
- Certain foods or drinks are more likely to cause heartburn, such as rich, spicy, fatty, and fried foods, chocolate, caffeine, alcohol, and even some fruits and vegetables.
- Eat slowly and do not eat big meals.
- If you are overweight, lose weight.
- If you smoke, quit smoking.
- Raise the head of your bed.
- Wear loose fitting clothing around your stomach.

NEW! NDC 14837-889-90

Pepcid Complete

Acid Reducer + Antacid

WITH DUAL ACTION

Reduces AND Neutralizes Acid

90 Minute Flavored Chewable Tablets

Works with Just One Tablet!

Contains Calcium! Relieves Heartburn Due to Acid Indigestion!

FREE PEPCID COMPLETE Offer

(When Offer Expires 9/1/2003)

DO NOT USE IF PRINTED FOL LAMINER SEAL UNDER BOTTLE CAP IS BROKEN.

The benefits of a chewable antacid

+ Acid control of Pepcid AC

= A single tablet with antacid, acid reducer and antacid.

Drug Facts

(in each chewable tablet)

Active Ingredients	Purpose
Famotidine 10 mg	Acid reducer
Calcium carbonate 800 mg	Antacid
Magnesium hydroxide 165 mg	Antacid

Use to relieve heartburn associated with acid indigestion and sour stomach.

Questions? 1-800-755-4008

Look for Warnings and Directions

Drug Facts (continued)

Warnings

Allergy alert: Do not use if you are allergic to famotidine or other acid reducers.

Do not use

- if you have trouble swallowing
- with other famotidine products or acid reducers
- if you are presently taking a prescription drug. Antacids may interact with certain prescription drugs.

Ask a doctor or pharmacist before use if you are

- pregnant or breastfeeding, ask a health professional before use.
- taking birth control pills, you need to take this product for more than 14 days.
- taking other medicines, you need to take this product for more than 14 days.
- taking other medicines, you need to take this product for more than 14 days.
- taking other medicines, you need to take this product for more than 14 days.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- adults and children 12 years and over:
 - do not swallow tablet whole; chew completely
 - to relieve symptoms, chew 1 tablet before swallowing
 - do not use more than 2 chewable tablets in 24 hours
- children under 12 years: ask a doctor

Other information


- read the directions and warnings before use
- store at 25-30°C (77-86°F)
- product from moisture

Inactive ingredients: cellulose acetate, corn starch, dextrates, flavors, hydroxypropyl cellulose, hydroxypropyl methylcellulose, lactose, magnesium stearate, pregelatinized starch, red iron oxide, sodium lauryl sulfate, sugar

Questions or comments? call toll-free 1-800-755-4008

A NON-PRESCRIPTION STOMACH MEDICINE

PEPCID COMPLETE combines an acid reducer (famotidine) with antacids (calcium carbonate and magnesium hydroxide) to relieve heartburn in two different ways. Acid reducers decrease the production of new stomach acid; antacids neutralize acid that is already in the stomach. The active ingredients in PEPCID COMPLETE have been used for years to treat acid-related problems in millions of people safely and effectively.



Shown actual size

- Just 1 tablet relieves heartburn due to acid indigestion.

1) How frequently do you use non-prescription antacids or stomach remedies?

A. 2 to 3 days per week or more B. 1 day per week

C. 2 to 3 days per month or less

2) How severe is a typical heartburn or indigestion episode?

A. very severe B. somewhat severe

C. not very severe D. not at all severe

3) PEPCID COMPLETE is a combination of a fast-acting chewable antacid, and a powerful acid controller like Zantac 75® or Pepcid AC. Which products will you replace with PEPCID COMPLETE?

A. _____

B. _____


Please exclude me from all future PEPCID COMPLETE offers or information.

NOTE: Offer good only in U.S.A. and APO/FPO addresses. This request form may not be mechanically reproduced. LIMIT TWO PEPCID COMPLETE OFFERS PER FAMILY AND/OR ADDRESS. No group or organization requests will be honored. Your offer rights may not be transferred or assigned. Offer void where prohibited or taxed. Please allow 6 to 8 weeks for receipt of package.

© Registered trademark of Merck & Co., Inc.
*Registered trademark of Glaxo Wellcome Warner-Lambert OTCGP



PROVEN EFFECTIVE IN CLINICAL STUDIES
 In clinical studies, **PEPCID COMPLETE** was significantly better than placebo pills in relieving heartburn.



QUESTIONS OR COMMENTS?




Please call 1-800-355-4008 for more information and a guide on how to help control heartburn.

Registered trademark of Merck & Co., Inc.
 Distributed by: **Johnson & Johnson • MERCK**
 CONSUMER PHARMACEUTICALS CO.
 FORT WASHINGTON, PA 19034 USA

Labeling Format Information:

Fonts: Univers condensed, bold and bold italic.
 Drug Facts: 9 pt | Leading: 6-6.56 pt
 Header: 8-8.75 pt | Bullets: 6.56 pt
 Subheader: 6-6.56 pt | Barlines: 2.5 pt
 Body Text: 6-6.56 pt | Hairlines: .5 pt
 Drug Facts (continued): 9.84 pt
 Horizontal Scale: 100-105% | Average Kerning: 0



Peppid[®] Complete

Do not use if the individual pouch is open or torn.

INDIVIDUAL POUCHES

Peppid[®] Complete
 Acid control
 of Peppid AC
 neutralizes acid!

A single label

Peppid[®] Complete
 Acid control
 of Peppid AC
 neutralizes acid!

The benefits of Peppid[®] Complete are enhanced when used with Peppid AC.

Drug Facts

Active ingredients (in each chewable tablet)

Famotidine 10 mg
 Calcium carbonate 800 mg
 Magnesium hydroxide 165 mg

Purpose
 Acid reducer
 Antacid

Use: relieves heartburn associated with acid indigestion and sour stomach.

Warnings
 Allergy alert: Do not use if you are allergic to famotidine or other acid reducers.

Do not use
 - if you have trouble swallowing - with other famotidine products or acid reducers.
 Ask a doctor or pharmacist before use if you are presently taking a prescription drug. Antacids may interact with certain prescription drugs.

Stop use and ask a doctor if:
 - stomach pain continues - you need to take this product for more than 14 days.
 - you are pregnant or breastfeeding. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions
 - adults: Chew 1 or 2 chewable tablets 2 or 3 times a day, before or after meals.
 - children 12 years and over: Chew 1 or 2 chewable tablets 2 or 3 times a day, before or after meals.
 - do not use more than 2 chewable tablets in 24 hours.
 - children under 12 years: Ask a doctor.

Other information

- Read the directions and warnings before use.
- Keep the carton and the edge insert. They contain important information.
- Store at 25-30°C (77-86°F) - protect from moisture.

Inactive ingredients: calcium acetate, corn starch, dextrates, flavors, hydroxypropyl cellulose, hydroxypropyl methylcellulose, lactose, magnesium stearate, pregelatinized starch, red iron oxide, sodium lauryl sulfate, sugar.


Questions or comments? call toll-free 1-800-755-4008




02000000

Labeling Format Information:

Fonts:	Univers condensed, bold and bold italic.
Drug Facts:	9 pt
Header:	8 pt
Subheader:	8 pt
Body Text:	6 pt
Drug Facts (continued):	10 pt
Horizontal Scale:	94%
Average Kerning:	-4




The benefits of a chewable antacid



A single tablet reduces and neutralizes acid.

Do not use if the individual pouch is open or torn.



Drug Facts

Active ingredients (in each chewable tablet)

Famotidine 10 mg
 Calcium carbonate 800 mg
 Magnesium hydroxide 165 mg

Purpose
 Acid reducer
 Antacid
 Antacid

Use: relieves heartburn associated with acid indigestion and sour stomach

Warnings
Allergy alert: Do not use if you are allergic to famotidine or other acid reducers

Do not use

- if you have trouble swallowing
- with other famotidine products or acid reducers

Ask a doctor or pharmacist before use if you are presently taking a prescription drug. Antacids may interact with certain prescription drugs.

Stop use and ask a doctor if:

- stomach pain continues
- you need to take this product for more than 14 days

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- adults and children 12 years and over
- do not swallow tablet whole; chew completely
- to relieve symptoms, chew 1 tablet before swallowing
- for more than 2 chewable tablets in 24 hours
- children under 12 years, ask a doctor

Other information

- read the directions and warnings before use
- keep the carton and package intact. They contain important information.
- store at 25-30°C (77-86°F) • protect from moisture

Drug Facts (continued)

Inactive ingredients: cellulose acetate, corn starch, dextrates, flavors, hydroxypropyl cellulose, hydroxypropyl methylcellulose, lactose, magnesium stearate, pregelatinized starch, red iron oxide, sodium lauryl sulfate, sugar

Questions or comments? call toll-free 1-800-755-4008

Distributed by:


Johnson & Johnson • MERCK

CONSUMER PHARMACEUTICALS CO.
 FORT WASHINGTON, PA 19034 USA


® Registered trademark of Merck & Co., Inc.

Horizontal Scale: 94%
 Average Kerning: -4

Labeling Format Information:
 Fonts: Unifers condensed, bold and bold italic.
 Drug Facts: 9 pt
 Header: 8 pt
 Subheader: 8 pt
 Body Text: 6 pt
 Drug Facts (continued): 10 pt
 Hairlines: 5 pt
 Barlines: 2.5 pt
 Bullets: 6 pt
 Leading: 6 pt




XXXXXXXXXX



NEW!
 Works with
 Just One Tablet!

Pepcid Complete
 Acid Reducer + Antacid
 WITH DUAL ACTION
 Reduces AND
 Neutralizes Acid



Take-Along Packs

35 Mint Flavored
 Chewable Tablets

Contains Calcium!
 Relieves Heartburn Due To Acid Indigestion.

NDC 16837-888-35

NEW!

Pepcid Complete
 Acid Reducer + Antacid
 WITH DUAL ACTION
 Reduces AND
 Neutralizes Acid



NOT FOR INDIVIDUAL
 RETAIL SALE



**Take-Along
 Packs**

35 Mint Flavored
 Chewable Tablets

Contains Calcium!
 Relieves Heartburn Due To Acid Indigestion

Pepcid Complete
 Acid Reducer + Antacid
 WITH DUAL ACTION
 Reduces AND
 Neutralizes Acid

Works with
 Just One Tablet!

The ONLY one
 that's two



Starts to
 neutralize AND
 works for
 hours!
 in seconds

Pepcid Complete
 Acid Reducer + Antacid
 WITH DUAL ACTION
 Reduces AND
 Neutralizes Acid
 Works with
 Just One Tablet!

Works with
 Just One Tablet!



Take-Along Pack

Labeling Format Information:

Fonts: Univers condensed, bold and bold italic.	
Drug Facts: 9.72 pt	Leading: 6.58 pt
Header: 8.64 pt	Bullets: 8.05 pt
Subheader: 6.48 pt	Barlines: 2.5 pt
Body Text: 6.48 pt	Hairlines: .5 pt
Drug Facts (continued): NA	

Horizontal Scale: 90% Average Kerning: -20







The benefits of a chewable antacid + Acid control of Pepcid AC = A simple tablet that both reduces and neutralizes acid.

Do not use if the individual pouch is open or torn.

<p>Drug Facts</p> <p>Active ingredients (in each chewable tablet)</p> <p>Famotidine 10 mg..... Calcium carbonate 800 mg..... Magnesium hydroxide 165 mg.....</p> <p><i>Use relieves heartburn associated with acid indigestion and sour stomach.</i></p>	<p>Purpose</p> <p>Acid reducer Antacid</p>	<p>Warnings</p> <p>Allergy alert: Do not use if you are allergic to famotidine or other acid-reducers Do not use if you have trouble swallowing. *with other famotidine products or acid reducers Ask a doctor or pharmacist before use if you are presently taking a prescription drug. Antacids may interact with certain prescription drugs. Stop use and ask a doctor if: stomach pain continues *you need to take this product for more than 14 days If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.</p>	<p>Directions</p> <ul style="list-style-type: none"> * adults and children 12 years and over: * do not swallow tablet whole; chew completely * to relieve symptoms, chew 1 tablet before swallowing * do not use more than 2 chewable tablets in 24 hours * children under 12 years: ask a doctor <p>Other information</p> <ul style="list-style-type: none"> * read the directions and warnings before use * keep the carton and package insert. They contain important information. * store at 25-30°C (77-86°F) * protect from moisture <p>Inactive ingredients: cellulose acetate, corn starch, dextrates, flavors, hydroxypropyl cellulose, hydroxypropyl methylcellulose, lactose, magnesium stearate, pregelatinized starch, red iron oxide, sodium lauryl sulfate, sugar</p> <p>Questions or comments? call toll-free 1-800-755-4008</p>
--	--	---	--



Distributed by:
Johnson & Johnson • MERCK
 CONSUMER PHARMACEUTICALS CO.
 FORT WASHINGTON, PA 19034 USA

Proven effective in clinical studies

Duration of Relief
Percent of heartburn episodes relieved for at least 7 hours

Treatment	Percent of heartburn episodes relieved for at least 7 hours
PEPCID COMPLETE	70
PEPCID	59
Placebo	61

Onset of Relief
Percent of heartburn episodes relieved within 30 minutes

Treatment	Percent of heartburn episodes relieved within 30 minutes
PEPCID COMPLETE	45
PEPCID	33
Placebo	41

In clinical studies, PEPCID COMPLETE was significantly better than placebo pills in relieving heartburn.

Warnings

Heartburn and acid indigestion are common, but you should stop taking PEPCID COMPLETE and see your doctor promptly if:

- You have trouble swallowing or stomach pain continues.
- You need to take this product for more than 14 days.
- You may have a serious condition that may need different treatment.
- Important: Allergy alert:** Do not use if you are allergic to famotidine or other acid reducers. Do not use with other famotidine products or acid reducers.
- Antacids may interact with certain prescription drugs. If you are presently taking a prescription drug, ask a doctor or pharmacist before use.
- This product should not be given to children under 12 years old, unless directed by a doctor.
- If pregnant or breast-feeding, ask a health professional before use.
- Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

Questions or comments?

© Registered trademark of Merck & Co., Inc.
 Distributed by: **Johnson & Johnson • MERCK**
 CONSUMER PHARMACEUTICALS CO.
 FORT WASHINGTON, PA 19044 USA

0000000

PEPCID COMPLETE \$2.00 REBATE OFFER

00000000

Please send my \$2.00 refund to:

Name: _____
 Address: _____
 Apt. # _____
 City: _____
 State: _____
 Zip: _____

LIMIT ONE OFFER PER FAMILY AND/OR ADDRESS
 Requests must be postmarked by December 31, 2003.

Note: Offer good only in U.S.A. and AP/CF PD addresses. Reproduction of this offer in any medium or form without permission from Johnson & Johnson/Consumer Pharmaceuticals Company violates copyright statutes. Subscribers that do not meet requirements or are duplicates will not be honored or returned. Certificate proof-of-purchase or cash register receipts is prohibited. Please allow 6 to 8 weeks for delivery.

Mail-In Certificate
 Offer Expires September 1, 2003
 To receive your \$2.00 refund by mail on PEPCID COMPLETE 25 count or larger, Buy: One PEPCID COMPLETE 25 count or larger between September 1, 2000 and September 1, 2003. Sorry, other sizes indicated are valid on this offer.


Mail: This completed certificate, one UPC symbol from the outer carton and original cash register receipt dated between September 1, 2000, and September 1, 2003, with price paid circle, Sent to: PEPCID COMPLETE, 12701 \$2.00 Refund Offer, P.O. Box 3719 Highland Park, MI 48203-0719.



Labeling Format Information:

Font: Univers condensed, bold and bold italic.
 Drug Facts: NA
 Header: 10 pt
 Subheader: 8 pt
 Body Text: 8 pt
 Drug Facts (continued): NA

Leading: 10 pt
 Bullets: 8 pt
 Barlines: NA pt
 Hairlines: NA pt

Horizontal Scale: 94% Average Kerning: -4


		CONSUMER PHARMACEUTICALS CO. CAMP HILL ROAD, FORT WASHINGTON, PA, 19034	
DIVG. NO. BROCHURE -42		TITLE: BROCHURE, DOUBLE FOLD	
DRAWN ARS		DATE 10/10/97	
PLDT SCALE 1:1		APPROVED	
ARS 1		DATE 11/24/97	
INIT.		REVISIONS	
NO.		DATE	
1		11/24/97	
REMOVED BLEEDER BARS		REVISIONS	

NOTES:
 1) ALL DIMENSIONS IN INCHES.
 2) TOLERANCES UNLESS SPECIFIED:
 FRACTIONS ±1/64
 3)  NO COLOR, NO COPY, NO BLEED
 4)  FOLD LINE


DO NOT USE IF THE INDIVIDUAL POUCH IS OPEN OR TORN.

WHAT YOU SHOULD KNOW ABOUT:

PACKAGE INSERT




Acid Reducer + Antacid



A non-prescription stomach medicine

PEPCID COMPLETE combines an acid reducer (famotidine) with antacids (calcium carbonate and magnesium hydroxide) to relieve heartburn in two different ways: Acid reducers decrease the production of new stomach acid; antacids neutralize acid that is already in the stomach. The active ingredients in PEPCID COMPLETE have been used for years to treat acid-related problems in millions of people safely and effectively.




• Just 1 tablet relieves heartburn due to acid indigestion.

Shown-actual size

How to use PEPCID COMPLETE

Use PEPCID COMPLETE to relieve heartburn due to acid indigestion and sour stomach. Children under 12 years: ask a doctor



TO RELIEVE SYMPTOMS
 Chew 1 tablet completely before swallowing.


PEPCID COMPLETE can be used up to twice daily (up to 2 chewable tablets in 24 hours).

Excess acid: a burning problem

It is normal for the stomach to produce acid, especially after consuming food and beverages. However, acid in the stomach may move up into the wrong place (the esophagus), causing burning pain and discomfort that interfere with everyday activities.

• Heartburn — Caused by acid in the esophagus

A valve-like muscle called the lower esophageal sphincter (LES) is relaxed in an open position




Burning pain/discomfort in esophagus

Acid moves up from stomach

Tips for Managing Heartburn

- Do not lie flat or bend over soon after eating.
- Do not eat late at night, or just before bedtime.
- Certain foods or drinks are more likely to cause heartburn, such as rich, spicy, fatty, and fried foods, chocolate, caffeine, alcohol, and even some fruits and vegetables.
- Eat slowly and do not eat big meals.
- If you are overweight, lose weight.
- If you smoke, quit smoking.
- Raise the head of your bed.
- Wear loose fitting clothing around your stomach.

Labeling Format Information:
 Fonts: Univers condensed, bold and bold italic.
 Drug Facts: NA
 Header: 10 pt
 Subheader: 8 pt
 Body Text: 8 pt
 Drug Facts (continued): NA
 Horizontal Scale: 94%
 Average Kerning: -4



Proven effective in clinical studies

In clinical studies, PEPCID COMPLETE was significantly better than placebo pills in relieving heartburn.

Onset of Relief

Percent of heartburn episodes relieved within 30 minutes

PEPCID COMPLETE	45
Placebo	33
Famotidine	41
Antacid	10

Duration of Relief

Percent of heartburn episodes relieved for at least 7 hours

PEPCID COMPLETE	70
Placebo	59
Famotidine	61
Antacid	10

Warnings

Heartburn and acid indigestion are common, but you should stop taking PEPCID COMPLETE and see your doctor promptly if:

- You have trouble swallowing or stomach pain continues.
- You need to take this product for more than 14 days.
- You may have a serious condition that may need different treatment.

Important:

Allergy alert: Do not use if you are allergic to famotidine or other acid reducers. Do not use with other famotidine products or acid reducers.


Antacids may interact with certain prescription drugs. If you are presently taking a prescription drug, ask a doctor or pharmacist before use.

This product should not be given to children under 12 years old, unless directed by a doctor.

- If pregnant or breast-feeding, ask a health professional before use.
- Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Questions or comments?

Please call 1-800-755-4008 for more information and a guide on how to help control heartburn.



© Registered trademark of Merck & Co., Inc.
 CONSUMER PHARMACEUTICALS CO.
 FORT WASHINGTON, PA 19034 USA
 Distributed by:
Johann-Johnen • MERCK


0000000

© Registered trademark of Merck & Co., Inc.
 CONSUMER PHARMACEUTICALS CO.
 FORT WASHINGTON, PA 19034 USA
 Distributed by:
Johann-Johnen • MERCK

Its unique DUAL ACTION formula makes it the best heartburn reliever.

Congratulations on your purchase of PEPCID COMPLETE. We're sure you'll agree that





Do not use if the individual pouch is open or torn.

	Drug Facts	
	Active ingredients (in each chewable tablet)	Purpose
	Famotidine 10 mg.....	Acid reducer
	Calcium carbonate 800 mg.....	Antacid
	Magnesium hydroxide 165 mg.....	Antacid
	Use relieves heartburn associated with acid indigestion and sour stomach	
	Warnings	
	Allergy alert: Do not use if you are allergic to famotidine or other acid reducers	
	Do not use	
	• if you have trouble swallowing • with other famotidine products or acid reducers	
	Ask a doctor or pharmacist before use if you are presently taking a prescription drug. Antacids may interact with certain prescription drugs.	
	Stop use and ask a doctor if	
	• stomach pain continues • you need to take this product for more than 14 days	
	If pregnant or breast-feeding, ask a health professional before use.	
	Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.	
	Directions	
	• adults and children 12 years and over:	
	• do not swallow tablet whole: chew completely	
	• to relieve symptoms, chew 1 tablet before swallowing	
	• do not use more than 2 chewable tablets in 24 hours	
	• children under 12 years: ask a doctor	
	Other information	
	• read the directions and warnings before use	
	• keep the carton. It contains important information.	
	• store at 25-30°C (77-86°F) • protect from moisture	
	Inactive ingredients cellulose acetate, corn starch, dextrates, flavors, hydroxypropyl cellulose, hydroxypropyl methylcellulose, lactose, magnesium stearate, pregelatinized starch, red iron oxide, sodium lauryl sulfate, sugar	
	Questions or comments? call toll-free 1-800-755-4008	

WHAT PATIENTS SHOULD KNOW ABOUT:

Pepcid Complete

WHAT PATIENTS SHOULD KNOW ABOUT:

Proven effective in relieving heartburn.
In clinical studies, PEPICID COMPLETE was significantly more effective than placebo in relieving heartburn.



Warnings

Heartburn and acid indigestion are common, but you should see your doctor promptly if you have trouble swallowing or stomach pain continues for more than 14 days. You may have a serious condition that may need diagnosis. **Important: Allergy alert:** Do not use if you are allergic to famotidine products or acid reducers. Do not use with other famotidine products or acid reducers. This product should not be given to children under 12 years of age. If pregnant or breast-feeding, ask a health professional for advice. Keep out of reach of children. In case of overdose, call your Poison Control Center right away.

Questions or comments

Please call 1-800-755-4008 for more information and a guide on how to help control your heartburn.



© Registered trademark of Merck & Co., Inc.
Distributed by: **Johnson & Johnson** • MERCK
CONSUMER PHARMACEUTICALS CO.

NEW! **PEPICID Complete**
Acid Reducer + Antacid

WITH DUAL ACTION
Reduces AND Neutralizes Acid

Works with Just One Tablet!

Contains Calcium!
Relieves Heartburn Due to Acid Indigestion

25 Mint Flavored Chewable Tablets

WHAT PATIENTS SHOULD KNOW ABOUT:

Pepcid Complete

A non-prescription stomach medicine

PEPICID COMPLETE combines an acid reducer (famotidine) with antacids (calcium carbonate and magnesium hydroxide) to relieve heartburn in two different ways: Acid reducers decrease the production of new stomach acid; antacids neutralize acid that is already in the stomach. The active ingredients in PEPICID COMPLETE have been used for years to treat acid-related problems in millions of people safely and effectively.

• Just 1 tablet relieves heartburn due to acid indigestion.

Shown actual size

How to use PEPICID COMPLETE

Use PEPICID COMPLETE to relieve heartburn due to acid indigestion and sour stomach. Children under 12 years of age should ask a doctor.




TO RELIEVE SYMPTOMS
Chew 1 tablet completely before swallowing.

PEPICID COMPLETE can be used up to twice daily up to 2 chewable tablets in 24 hours.

Excess acid: a burning problem

It is normal for the stomach to produce acid, especially after consuming food and beverages. However, acid in the stomach may move up into the wrong place (the esophagus), causing burning pain and discomfort that interfere with everyday activities.

• **Heartburn—Caused by acid in the esophagus**



- Tips for Managing Heartburn**
- Do not lie flat or bend over soon after eating.
 - Do not eat late at night, or just before bedtime.
 - Certain foods or drinks are more likely to cause heartburn, such as rich, spicy, fatty, and fried foods, chocolate, caffeine, alcohol, and even some fruits and vegetables.
 - Eat slowly and do not eat big meals.
 - If you are overweight, lose weight.
 - If you smoke, quit smoking.
 - Raise the head of your bed.
 - Wear loose fitting clothing around your stomach.

Pepcid Complete

WHAT PATIENTS SHOULD KNOW ABOUT:

Proven effective in clinical studies

In clinical studies, PEPCID COMPLETE was significantly better than placebo pills in relieving heartburn.



Warnings

Heartburn and acid indigestion are common, but you should stop taking PEPCID COMPLETE and see your doctor promptly if:

- You have trouble swallowing or stomach pain continues.
- You need to take this product for more than 14 days.
- You may have a serious condition that may need different treatment.
- Important: Allergy alert:** Do not use if you are allergic to famotidine or other acid reducers.
- Antacids may interact with certain prescription drugs. If you are presently taking a prescription drug, ask a doctor or pharmacist before use.
- This product should not be given to children under 12 years old, unless directed by a doctor.
- If pregnant or breast-feeding, ask a health professional before use.
- Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Questions or comments?

Please call 1-800-755-4008 for more information and a guide on how to help control heartburn.



© Registered trademark of Merck & Co., Inc.
Distributed by: **Johnson & Johnson** **MERCK**
CONSUMER PHARMACEUTICALS CO., FORT WASHINGTON, PA 19034 USA

NEW!
Pepcid Complete
Acid Reducer + Antacid

WITH **DUAL ACTION**
Reduces AND Neutralizes Acid

Works WITH Just One Tablet!

Contains Calcium!
Relieves Heartburn Due To Acid Indigestion

25 Mint Flavored Chewable Tablets

Pepcid Complete

rescription stomach medicine

as an acid reducer (famotidine) with antacids (calcium hydroxide) to relieve heartburn in two different ways: Acid reduction of new stomach acid; antacids neutralize acid that is a active ingredients in PEPCID COMPLETE have been used for problems in millions of people safely and effectively.

at 1 tablet relieves heartburn due to acid indigestion.

How to use PEPCID COMPLETE

PEPCID COMPLETE can be used up to twice daily up to 2 chewable tablets in 24 hours).




Less acid: a burning problem

ch to produce acid, especially after consuming food and in the stomach may move up into the wrong place (the esophagus) causing pain and discomfort that interfere with everyday activities.

Tips for Managing Heartburn

- Do not lie flat or bend over soon after eating.
- Do not eat late at night, or just before bedtime.
- Certain foods or drinks are more likely to cause heartburn, such as rich, spicy, fatty, and fried foods, chocolate, caffeine, alcohol, and even some fruits and vegetables.
- Eat slowly and do not eat big meals.
- If you are overweight, lose weight.
- If you smoke, quit smoking.
- Raise the head of your bed.
- Wear loose fitting clothing around your stomach.



Warnings
 Allergy alert: Do not use if you are allergic to famotidine or other acid reducers

Dist. by: **Janssen** • **Janssen** • **MERCK CONSUMER PHARMACEUTICALS CO.**
 FORT WASHINGTON, PA 19034 USA. ©Registered trademark of Merck & Co., Inc.

Inactive ingredients cellulose acetate, corn starch, dextrans, flavors, hydroxypropyl cellulose, hydroxypropyl methylcellulose, lactose, magnesium stearate, pregelatinized starch, red iron oxide, sodium lauryl sulfate, sugar

Questions or comments? call toll-free 1-800-755-4008

150% OF SIZE

Labeling Format Information:

Fonts: Univers condensed, bold and bold italic.

Drug Facts: 8 pt
Header: 7 pt
Subheader: 6 pt
Body Text: 6 pt
Barlines: 1.5 pt
Halftones: 5 pt
Drug Facts (continued): 8 pt

Average Kerning: -4

Horizontal Scale: 94%

While folded on line, tear open at slit

Peppid[®] Complete
 WITH DUAL ACTION
 Acid Reducer + Antacid

Reduces AND Neutralizes Acid
Contains Calcium!
Relieves Heartburn Due to Acid Indigestion

Works with Just One Tablet!
Mint Flavored Chewable Tablet

Do not use if pouch is open or torn.

Drug Facts

Active ingredients (for each chewable tablet) Purpose
 Famotidine 10 mg..... Acid reducer
 Calcium carbonate 600 mg..... Antacid
 Magnesium hydroxide 165 mg..... Antacid

Use relieves heartburn associated with acid indigestion and sour stomach

Warnings
Allergy Alert: Do not use if you are allergic to famotidine or other acid reducers

Dist. by **Janssen** • **Janssen** • **MERCK CONSUMER PHARMACEUTICALS CO.**
 FORT WASHINGTON, PA 19034 USA. ©Registered trademark of Merck & Co., Inc.

While folded on line, tear open at slit

Drug Facts (continued)

Warnings (continued)
 Do not use if you have trouble swallowing
 • with other famotidine products or acid reducers

Ask a doctor or pharmacist before use if you are presently taking a prescription drug. Antacids may interact with certain prescription drugs.

Stop use and ask a doctor if: stomach pain continues
 • you need to take this product for more than 14 days

If pregnant or breast feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions
 • adults and children 12 years and over
 • do not swallow tablet whole, chew completely
 • to relieve symptoms, chew 1 tablet before swallowing
 • do not use more than 2 chewable tablets in 24 hours
 • children under 12 years: ask a doctor

Other information
 • store at 25-30°C (77-86°F) • protect from moisture

Inactive ingredients cellulose acetate, corn starch, dextrans, flavors, hydroxypropyl cellulose, hydroxypropyl methylcellulose, lactose, magnesium stearate, pregelatinized starch, red iron oxide, sodium lauryl sulfate, sugar

Questions or comments? call toll-free 1-800-755-4008

