


**Base Label,
Page 3**




Cover Page 1



**Page 2
(Back of Cover)**

Labeling Format Information: Fonts: HelveticaDF Cond Medium, Cond Bold, Cond Bold Italic
 Drug Facts: 9.00 pt | Body Text: 6.00 pt | Bullets: 5.00 pt | Horizontal Scale: 100%
 Headers: 8.00 pt | Leading: 6.50 pt | Barlines: 2.50 pt | Average Kerning: 0
 Subheaders: 6.00 pt | Drug Facts (continued): 8.00 pt | Hairlines: 0.50 pt

McNeil Submission Artwork



Drug Facts (continued)

Other information

- each tablet contains: calcium 320 mg, magnesium 70 mg
- read the directions and warnings before use
- read the bottle. It contains important information.
- store at 20°-25°C (68°-77°F) • protect from moisture
- do not use if foil seal under bottle cap printed "SAFETY SEAL®" is open or torn

Inactive ingredients cellulose acetate, corn starch, croscopolone, D&C yellow no. 10 aluminum lake, dextrose excipient, FD&C blue no. 1 aluminum lake, flavors, gum arabic, hydroxypropyl cellulose, hypromellose, lactose monohydrate, magnesium stearate, maltodextrin, mineral oil, sucralose

Questions or comments? call 1-800-755-4008 (toll-free) or 215-273-8755 (collect)

Pepcid Complete
Acid Reducer + Antacid
Famotidine 10 mg per tablet
Calcium carbonate 800 mg per tablet
Magnesium hydroxide 165 mg per tablet
50 Chewable Tablets

Shown actual size

Tips for Managing Heartburn

- Do not lie flat or bend over after eating
- Do not wear tight-fitting clothing around the stomach
- Do not eat before bedtime
- Raise the head of your bed
- Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol, and certain fruits and vegetables
- Eat slowly and avoid big meals
- If overweight, lose weight
- Quit smoking

Base Label
Page 3

COVER PAGE 1

Cover
Page 1

Drug Facts (continued)

Ask a doctor before use if you have

- had heartburn over 3 months. This may be a sign of a more serious condition.
- heartburn with **lightheadedness, sweating, or dizziness**
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
- frequent chest pain
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain

Ask a doctor or pharmacist before use if you are presently taking a prescription drug. Antacids may interact with certain prescription drugs.

Stop use and ask a doctor if

- your heartburn continues or worsens
- you need to take this product for more than 14 days

If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions


- adults and children 12 years and over:
 - do not swallow tablet whole; chew completely
 - to relieve symptoms, chew 1 tablet before swallowing
 - do not use more than 2 chewable tablets in 24 hours
- children under 12 years: ask a doctor

Page 2
(Back of Cover)

Labeling Format Information: Fonts: HelveticaDF Cond Medium, Cond Bold, Cond Bold Italic

Drug Facts: 9.00 pt	Body Text: 6.00 pt	Bullets: 5.00 pt	Horizontal Scale: 100%
Headers: 8.00 pt	Leading: 6.50 pt	Barlines: 2.50 pt	
Subheaders: 6.00 pt	Drug Facts (continued): 8.00 pt	Hairlines: 0.50 pt	Average Kerning: 0


McNeil Submission Artwork



Base Label,
Page 3




Cover Page 1



Page 2
(Back of Cover)

Labeling Format Information: Fonts: Univers 57 Condensed, 67 Bold, Bold Condensed Oblique
 Drug Facts: 8.50 pt | Body Text: 6.00 pt | Bullets: 5.00 pt | Horizontal Scale: 90%
 Headers: 8.00 pt | Leading: 6.50 pt | Barlines: 2.50 pt
 Subheaders: 6.00 pt | Drug Facts (continued): 8.00 pt | Hairlines: 0.50 pt | Average Kerning: 0

McNeil Submission
Artwork



Drug Facts (continued)

Other information

- each tablet contains: calcium 320 mg, magnesium 70 mg
- read the directions and warnings before use
- read the bottle. It contains important information.
- store at 20°-25°C (68°-77°F) • protect from moisture
- do not use if foil seal under bottle cap printed "SAFETY SEAL®" is open or torn

Inactive ingredients cellulose acetate, corn starch, croscopolone, D&C red no. 7 calcium lake, dextrose excipient, FD&C blue no. 1 aluminum lake, FD&C red no. 40 aluminum lake, flavors, gum arabic, hydroxypropyl cellulose, hypromellose, lactose monohydrate, magnesium stearate, maltodextrin, mineral oil, modified starch, sucralose

Questions or comments? call 1-800-755-4008 (toll-free) or 215-273-8755 (collect)

Pepcid Complete

Acid Reducer + Antacid
Famotidine 10 mg per tablet
Calcium carbonate 800 mg per tablet
Magnesium hydroxide 165 mg per tablet
50 Chewable Tablets

Shown actual size

Tips for Managing Heartburn

- Do not lie flat or bend over after eating
- Do not wear tight-fitting clothing around the stomach
- Do not eat before bedtime
- Raise the head of your bed
- Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol, and certain fruits and vegetables
- Eat slowly and avoid big meals
- If overweight, lose weight
- Quit smoking

Distributed by: **JOHNSON & JOHNSON CONSUMER INC.**
McNeil Consumer Healthcare Division
Fort Washington, PA 19034 USA
©McNeil Consumer Pharmaceuticals Co. 2016
PEPCID® and PEPCID COMPLETE® are registered trademarks of McNeil CPC

Cover Page 1

Drug Facts (continued)

Ask a doctor before use if you have

- had heartburn over 3 months. This may be a sign of a more serious condition.
- heartburn with **lightheadedness, sweating, or dizziness**
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
- frequent chest pain
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain

Ask a doctor or pharmacist before use if you are presently taking a prescription drug. Antacids may interact with certain prescription drugs.

Stop use and ask a doctor if

- your heartburn continues or worsens
- you need to take this product for more than 14 days

If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions


- adults and children 12 years and over:
 - do not swallow tablet whole; **chew completely**
 - to relieve symptoms, chew 1 tablet before swallowing
 - do not use more than 2 chewable tablets in 24 hours
- children under 12 years: ask a doctor

**Page 2
(Back of Cover)**


Labeling Format Information: Fonts: HelveticaDF Cond Medium, Cond Bold, Cond Bold Italic

Drug Facts: 9.00 pt	Body Text: 6.00 pt	Bullets: 5.00 pt	Horizontal Scale: 100%
Headers: 8.00 pt	Leading: 6.50 pt	Barlines: 2.50 pt	
Subheaders: 6.00 pt	Drug Facts (continued): 8.00 pt	Hairlines: 0.50 pt	Average Kerning: 0

McNeil Submission Artwork




Cover Page 1



Base Label, Page 3



Cover Page 1




Page 2 (Back of Cover)

Labeling Format Information: Fonts: Univers 57 Condensed, 67 Bold Condensed, Oblique

Drug Facts: 9 pt	Body Text: 6 pt	Bullets: 5 pt	Horizontal Scale: 90%
Headers: 8 pt	Leading: 6.5 pt	Barlines: 2.5 pt	
Subheaders: 6 pt	Drug Facts (continued): 8 pt	Hairlines: 0.5 pt	Average Kerning: 0


McNeil Submission Artwork



**Base Label,
Page 3**




Cover Page 1



**Page 2
(Back of Cover)**

Labeling Format Information: Fonts: HelveticaDF Cond Medium, Cond Bold, Cond Bold Italic
 Drug Facts: 9.00 pt | Body Text: 6.00 pt | Bullets: 5.00 pt | Horizontal Scale: 100%
 Headers: 8.00 pt | Leading: 6.50 pt | Barlines: 2.50 pt
 Subheaders: 6.00 pt | Drug Facts (continued): 8.00 pt | Hairlines: 0.50 pt | Average Kerning: 0

McNeil Submission Artwork



Base Label Page 3

Base Label Page 3

Cover Page 1

Page 2 (Back of Cover)

This is a representation of an electronic record that was signed electronically and this page is the manifestation of the electronic signature.

/s/

VALERIE S PRATT
06/08/2016