

Drug Facts Format Information					
Drug Facts:	NA pt	Subheadings/Body Text:	4.25 pt	Barlines: NA pt	Horizontal Scale: 80%
Drug Facts (continued):	NA pt	Bullets:	4.25 pt	Hairlines: NA pt	Tracking: 0
Headings:	4.25 pt	Leading:	4.75 pt		

DUAL ACTION

Pepcid Complete

Acid Reducer + Antacid
Just One Tablet!
Berry Flavor


Relieves Heartburn
Due to Acid Indigestion **1 Chewable Tablet**

Do not use if speech is hoarse or changed.

Active ingredients (in each chewable tablet)

Famotidine 10 mg	Pepceon
Calcium carbonate 100 mg	Acid reducer
Magnesium hydroxide 100 mg	Antacid

Relieves heartburn associated with acid indigestion and sour stomach

To Open: White Folded on Line, Tear At Slit 

Warnings: Always alert: Do not use if you are allergic to famotidine or other acid reducers. Do not use if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.

Do not use if you have: Ask a doctor before use if you have heartburn over 3 weeks. This may be a sign of a more serious condition. • heartburn with **lightheadedness, swelling, or dizziness** • chest pain or shoulder pain with shortness of breath, sweating, pain spreading to arms, neck or shoulders, or light-headedness • frequent chest pain • frequent wheezing, particularly with heartburn • unexplained weight loss • nausea or vomiting • stomach pain • kidney disease. Ask a doctor or pharmacist before use if you are taking a prescription drug. Antacids and acid reducers may interact with certain prescription drugs. **Stop use and ask a doctor if:** • your heartburn continues or worsens • you need to take this product for more than 14 days, if pregnant or breast-feeding, ask a health professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-235-0223). **Directions:** • adults and children 12 years and over: • **do not swallow tablet whole; chew completely** • to relieve symptoms, chew 1 tablet before swallowing • do not use more than 2 chewable tablets in 24 hours • children under 12 years: ask a doctor. **Other information:** • each tablet contains: calcium 100 mg, magnesium 100 mg • read the directions and warnings before use • keep the carton. It contains important information. • store at 20°-25°C (68°-77°F) • protect from moisture • do not use if powder is loose or damaged. **Inactive ingredients:** cellulose acetate, corn starch, croscarmellose, D,L-lysine, 7-calcium beta, dextrin, croscarmellose, FD&C blue no. 1 aluminum lake, FD&C red no. 40 aluminum lake, flavors, gum arabic, hydroxypropyl methylcellulose, hypromellose, lactose monohydrate, magnesium stearate, methylcellulose, mineral oil, modified starch, sucralose. **†may contain mineral oil.**

Directions or comments: 1-800-305-4000 (toll-free) or 215-279-4700 (collect)

Dist. by: **Kanva Brands LLC**
Somers, NJ 07083
© Kanva Brands LLC 2024

NNNNNN EXP: YYYMM

0000000

Net Content Statement		NET CONTENT	AREA OF PDP
PDP Area: (width x height)	24.8 in ²	1/16 IN.	5 SQ. IN. OR LESS
Net Content Size:	0.102"	1/8 IN.	MORE THAN 5 SQ. IN. BUT NOT MORE THAN 25 SQ. IN.
		3/16 IN.	MORE THAN 25 SQ. IN. BUT NOT MORE THAN 100 SQ. IN.
		1/4 IN.	MORE THAN 100 SQ. IN. BUT NOT MORE THAN 400 SQ. IN.
		1/2 IN.	OVER 400 SQ. IN.

Drug Facts Format Information			
Drug Facts:	10.00 pt	Subheadings/Body Text:	6.00 pt
Drug Facts (continued):	8.00 pt	Bullets:	5.00 pt
Headings:	8.00 pt	Leadings:	6.50 pt
		Barlines:	2.5 pt
		Horizontal Scale:	90%
		Hairlines:	0.5 pt
		Tracking:	:20



Net Content Statement		NET CONTENT	AREA OF PDP
PDP Area: (width x height)	5.9 in ²	1/16 IN.	5 SQ. IN. OR LESS
Net Content Size:	0.078"	1/8 IN.	MORE THAN 5 SQ. IN. BUT NOT MORE THAN 25 SQ. IN.
		3/16 IN.	MORE THAN 25 SQ. IN. BUT NOT MORE THAN 100 SQ. IN.
		1/4 IN.	MORE THAN 100 SQ. IN. BUT NOT MORE THAN 400 SQ. IN.
		1/2 IN.	OVER 400 SQ. IN.

Drug Facts Format Information					
Drug Facts:	8.50 pt	Subheadings/Body Text:	6.00 pt	Barlines:	2.5 pt
Drug Facts (continued):	8.00 pt	Bullets:	5.00 pt	Horizontal Scale:	85/100%
Headings:	8.00 pt	Leading:	6.50 pt	Trailings:	0

7 168371291501 3
EXP NNNNNN
YYYYMM
00000000

Drug Facts (continued)
Other information
• each tablet contains: calcium 520 mg, magnesium 70 mg
• read the directions and warnings before use
• read the bottle. It contains important information.
• store at 20°-25°C (68°-77°F) • protect from moisture
• do not use if foil inner seal imprinted with "Sealed For Your Safety" is broken or missing
Inactive ingredients cellulose acetate, corn starch, croscopolone, D&C red no. 7 calcium lake, dextrose excipient, FD&C blue no. 1 aluminum lake, FD&C red no. 40 aluminum lake, flavors, gum arabic, hydroxypropyl cellulose, hypromellose, lactose monohydrate, magnesium stearate, methylcellulose, mineral oil*, modified starch, sucralose *may contain mineral oil
Questions or comments? call 1-800-755-8100 (toll-free) or 215-272-8285 (collect)

Pepcid Complete
Acid Reducer + Antacid
Famotidine 10 mg per tablet
Calcium carbonate 800 mg per tablet
Magnesium hydroxide 165 mg per tablet
50 Chewable Tablets



Shown actual size

Tips for Managing Heartburn
• Do not lie flat or bend over after eating
• Do not wear tight-fitting clothing around the stomach
• Do not eat before bedtime
• Raise the head of your bed
• Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol, and certain fruits and vegetables
• Eat slowly and avoid big meals
• If overweight, lose weight
• Quit smoking

DUAL ACTION NDC 16837-298-50
Pepcid Complete
Famotidine 10 mg—Acid reducer
Calcium carbonate 800 mg—Antacid
Magnesium hydroxide 165 mg—Antacid
Just One Tablet!
Relieves Heartburn Due to Acid Indigestion
Berry Flavor
50 Chewable Tablets

For Complete Directions and Warnings
Do not use if foil inner seal imprinted with "Sealed For Your Safety" is broken or missing.

Drug Facts
Active ingredients (in each chewable tablet)
Famotidine 10 mg..... Acid reducer
Calcium carbonate 800 mg..... Antacid
Magnesium hydroxide 165 mg..... Antacid
Use relieves heartburn associated with acid indigestion and sour stomach
Warnings
Allergy alert Do not use if you are allergic to famotidine or other acid reducers
Do not use
• if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.
• with other acid reducers

Manufactured by: Kenova Brands LLC
Summit, NJ 07901
©Kenova Brands LLC 2024


Drug Facts (continued)
Ask a doctor before use if you have
• had heartburn over 2 months. This may be a sign of a more serious condition.
• heartburn with lightheadedness, sweating, or dizziness
• chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
• frequent chest pain
• frequent wheezing, particularly with heartburn
• unexplained weight loss
• nausea or vomiting
• stomach pain
• kidney disease
Ask a doctor or pharmacist before use if you are taking a prescription drug. Antacids and acid reducers may interact with certain prescription drugs.
Stop use and ask a doctor if
• your heartburn continues or worsens
• you need to take this product for more than 14 days
If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)
Directions
• adults and children 12 years and over:
• do not swallow tablet whole; chew completely
• to relieve symptoms, chew 1 tablet before swallowing
• do not use more than 2 chewable tablets in 24 hours
• children under 12 years: ask a doctor

Net Content Statement		NET CONTENT	AREA OF PDP
1/16 IN.		5 SQ. IN. OR LESS	
1/8 IN.		MORE THAN 5 SQ. IN. BUT NOT MORE THAN 25 SQ. IN.	
3/16 IN.		MORE THAN 25 SQ. IN. BUT NOT MORE THAN 100 SQ. IN.	
1/4 IN.		MORE THAN 100 SQ. IN. BUT NOT MORE THAN 400 SQ. IN.	
1/2 IN.		OVER 400 SQ. IN.	

PDP Area: (width x height) 5.9 in²
Net Content Size: 0.076"

Drug Facts Format Information					
Drug Facts:	8.50 pt	Subheadings/Body Text:	6.00 pt	Barlines:	2.5 pt
Drug Facts (continued):	8.00 pt	Bullets:	5.00 pt	Hairlines:	0.5 pt
Headings:	8.00 pt	Leading:	6.50 pt	Horizontal Scale:	85/100%
				Tracking:	0

7 16377 29172 5



EXP NNNNNN
YYYYMM
00000000

Drug Facts (continued)

Other information


- each **Net Content**: calcium 320 mg, magnesium 70 mg
- read the directions and warnings before use
- read the bottle. It contains important information.
- store at 20°-25°C (68°-77°F) • protect from moisture
- do not use if foil inner seal imprinted with "Sealed For Your Safety" is broken or missing

Inactive ingredients: cellulose acetate, corn starch, croscopolone, D&C red no. 7 calcium lake, dextrose excipient, FD&C blue no. 1 aluminum lake, FDC red no. 40 aluminum lake, flavors, gum arabic, hydroxypropyl cellulose, hypromellose, lactose monohydrate, magnesium stearate, maltodextrin, mineral oil*, modified starch, sucralose *may contain mineral oil

Questions or comments? call 1-800-755-4008 (toll-free) or 215-273-8755 (collect)

Pepcid Complete
Acid Reducer + Antacid
Famotidine 10 mg per tablet
Calcium carbonate 800 mg per tablet
Magnesium hydroxide 165 mg per tablet

70 Chewable Tablets
Distributed by: Kenvue Brands LLC
Summit, NJ 07901
© Kenvue Brands LLC 2024



Shown actual size

Tips for Managing Heartburn

- Do not lie flat or bend over after eating
- Do not wear tight-fitting clothing around the stomach
- Do not eat before bedtime
- Raise the head of your bed
- Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol, and certain fruits and vegetables
- Eat slowly and avoid big meals
- If overweight, lose weight
- Quit smoking

VALUE SIZE NDC 16827-208-70

DUAL ACTION

Pepcid Complete

Famotidine 10 mg—Acid reducer
Calcium carbonate 800 mg—Antacid
Magnesium hydroxide 165 mg—Antacid

Just One Tablet!
Relieves Heartburn Due to Acid Indigestion
Berry Flavor

70 Chewable Tablets

actual size

For Complete Directions and Warnings

Do not use if foil inner seal imprinted with "Sealed For Your Safety" is broken or missing.

Drug Facts

Active ingredients (in each chewable tablet)	Purpose
Famotidine 10 mg	Acid reducer
Calcium carbonate 800 mg	Antacid
Magnesium hydroxide 165 mg	Antacid

Use relieves heartburn associated with acid indigestion and sour stomach

Warnings

Allergy alert: Do not use if you are allergic to famotidine or other acid reducers

Do not use

- if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.
- with other acid reducers

Distributed by:
Kenvue Brands LLC
Summit, NJ 07901
© Kenvue Brands LLC 2024

Drug Facts (continued)

Ask a doctor before use if you have

- had heartburn over 3 months. This may be a sign of a more serious condition.
- heartburn with light-headedness, sweating, or dizziness
- chest pain or shoulder pain with shortness of breath, sweating, pain spreading to arms, neck or shoulders, or lightheadedness
- frequent chest pain
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain
- kidney disease

Ask a doctor or pharmacist before use if you are taking a prescription drug. Antacids and acid reducers may interact with certain prescription drugs.

Stop use and ask a doctor if

- your heartburn continues or worsens
- you need to take this product for more than 14 days

If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions

- adults and children 12 years and over:
 - do not swallow tablet whole: chew completely
 - to relieve symptoms, chew 1 tablet before swallowing
 - do not use more than 2 chewable tablets in 24 hours
- children under 12 years: ask a doctor

Net Content Statement		NET CONTENT	AREA OF PDP
		1/10 IN.	5 SQ. IN. OR LESS
PDP Area: (width x height)	7.34 in ²	1/8 IN.	MORE THAN 5 SQ. IN. BUT NOT MORE THAN 25 SQ. IN.
Net Content Size:	0.075"	3/16 IN.	MORE THAN 25 SQ. IN. BUT NOT MORE THAN 100 SQ. IN.
		1/4 IN.	MORE THAN 100 SQ. IN. BUT NOT MORE THAN 400 SQ. IN.
		1/2 IN.	OVER 400 SQ. IN.

Drug Facts Format Information			
Drug Facts:	8.50 pt	Subheadings/Body Text:	6.00 pt
Drug Facts (continued):	8.00 pt	Bullets:	5.00 pt
Headings:	8.00 pt	Leading:	6.50 pt
		Barlines:	2.5 pt
		Hairlines:	0.5 pt
		Horizontal Scale:	95/100%
		Tracking:	0



NNNNNN
EXP. YYYY/MM

00000000

Drug Facts (continued)
Other information
<ul style="list-style-type: none"> each tablet contains: calcium 220 mg, magnesium 70 mg read the directions and warnings before use read the bottle. It contains important information. store at 20°-25°C (68°-77°F) protect from moisture do not use if foil inner seal imprinted with "Sealed For Your Safety" is broken or missing
Inactive ingredients cellulose acetate, corn starch, croscopolone, D&C red no. 7 calcium lake, dextrose excipient, FD&C blue no. 1 aluminum lake, FD&C red no. 40 aluminum lake, flavors, gum arabic, hydroxypropyl cellulose, hypromellose, lactose monohydrate, magnesium stearate, maltoedextrin, mineral oil*, modified starch, sucralose *may contain mineral oil
Questions or comments? call 1-800-755-4008 (toll-free) or 215-273-8755 (collect)

Pepcid Complete
Acid Reducer + Antacid
Famotidine 10 mg per tablet
Calcium carbonate 800 mg per tablet
Magnesium hydroxide 165 mg per tablet
100 Chewable Tablets
Distributed by Kenvue Brands LLC
Summit, NJ 07901
©Kenvue Brands LLC 2024



Shown actual size

Tips for Managing Heartburn

- Do not lie flat or bend over after eating
- Do not wear tight-fitting clothing around the stomach
- Do not eat before bedtime
- Raise the head of your bed
- Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol, and certain fruits and vegetables
- Eat slowly and avoid big meals
- If overweight, lose weight
- Quit smoking

NDC 16837-298-12

DUAL ACTION

Pepcid Complete

Famotidine 10 mg—Acid reducer
Calcium carbonate 800 mg—Antacid
Magnesium hydroxide 165 mg—Antacid

Just One Tablet!
Relieves Heartburn Due to Acid Indigestion
Berry Flavor

100 Chewable Tablets

actual size

For Complete Directions and Warnings

Do not use if foil inner seal imprinted with "Sealed For Your Safety" is broken or missing.

Active ingredients (in each chewable tablet)	Purposes
Famotidine 10 mg	Acid reducer
Calcium carbonate 800 mg	Antacid
Magnesium hydroxide 165 mg	Antacid

Use relieves heartburn associated with acid indigestion and sour stomach

Warnings

Allergy alert: Do not use if you are allergic to famotidine or other acid reducers

Do not use

- if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.
- with other acid reducers

Distributed by: Kenvue Brands LLC
Summit, NJ 07901
©Kenvue Brands LLC 2024

Drug Facts (continued)
Ask a doctor before use if you have
<ul style="list-style-type: none"> had heartburn over 3 months. This may be a sign of a more serious condition. heartburn with lightheadedness, sweating, or dizziness chest pain or shoulder pain with shortness of breath, sweating, pain spreading to arms, neck or shoulders; or lightheadedness frequent chest pain frequent wheezing, particularly with heartburn unexplained weight loss nausea or vomiting stomach pain kidney disease
Ask a doctor or pharmacist before use if you are taking a prescription drug. Antacids and acid reducers may interact with certain prescription drugs.
Stop use and ask a doctor if
<ul style="list-style-type: none"> your heartburn continues or worsens you need to take this product for more than 14 days
If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)
Directions
<ul style="list-style-type: none"> adults and children 12 years and over. do not swallow tablet whole; chew completely to relieve symptoms, chew 1 tablet before swallowing do not use more than 2 chewable tablets in 24 hours children under 12 years: ask a doctor

Net Content Statement		NET CONTENT	AREA OF PDP
PDP Area: (width x height)	5.93 in ²	1/16 IN.	MORE THAN 5 SQ. IN. BUT NOT MORE THAN 25 SQ. IN.
Net Content Size:	0.078"	1/8 IN.	MORE THAN 25 SQ. IN. BUT NOT MORE THAN 100 SQ. IN.
		3/16 IN.	MORE THAN 100 SQ. IN. BUT NOT MORE THAN 400 SQ. IN.
		1/4 IN.	MORE THAN 400 SQ. IN. BUT NOT MORE THAN 400 SQ. IN.
		1/2 IN.	OVER 400 SQ. IN.

Drug Facts Format Information					
Drug Facts:	8.50 pt	Subheadings/Body Text:	6.00 pt	Barlines:	2.5 pt
Drug Facts (continued):	8.00 pt	Bullets:	5.00 pt	Hairlines:	0.5 pt
Headings:	8.00 pt	Leading:	6.50 pt	Tracking:	0
				Horizontal Scale:	85/100%



NNNNNN
EXP YYY/YY

0000000

Drug Facts (continued)	
Other information	<ul style="list-style-type: none"> each tablet contains: calcium 220 mg, magnesium 70 mg read the directions and warnings before use read the bottle. It contains important information. store at 20°-25°C (68°-77°F) • protect from moisture do not use if foil inner seal imprinted with "Sealed For Your Safety" is broken or missing
Inactive ingredients	cellulose acetate, corn starch, croscopolidone, dextrose excipient, FD&C blue no. 1 aluminum lake, flavor, gum arabic, hydroxypropyl cellulose, hypromellose, lactose monohydrate, magnesium stearate, malto-dextrin, mineral oil*, sucralose, yellow iron oxide *may contain mineral oil
Questions or comments?	call 1-800-755-4088 (toll-free) or 215-273-8756 (collect)

Pepcid Complete
Acid Reducer + Antacid
Famotidine 10 mg per tablet
Calcium carbonate 800 mg per tablet
Magnesium hydroxide 165 mg per tablet
50 Chewable Tablets



Shown actual size

Distributed by: Kaneva Brands LLC
Sumner, NJ 07901
©Kaneva Brands LLC 2024

- Tips for Managing Heartburn**
- Do not lie flat or bend over after eating
 - Do not wear tight-fitting clothing around the stomach
 - Do not eat before bedtime
 - Raise the head of your bed
 - Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol, and certain fruits and vegetables
 - Eat slowly and avoid big meals
 - If overweight, lose weight
 - Quit smoking

NDC 14837-891-66

DUAL ACTION

Pepcid Complete

Famotidine 10 mg—Acid reducer
Calcium carbonate 800 mg—Antacid
Magnesium hydroxide 165 mg—Antacid

Just One Tablet!
Relieves Heartburn Due to Acid Indigestion
Cool Mint Flavor

50 Chewable Tablets

actual size

For Complete Directions and Warnings

Do not use if foil inner seal imprinted with "Sealed For Your Safety" is broken or missing

Drug Facts	
Active ingredients	Purposes
<i>(in each chewable tablet)</i>	
Famotidine 10 mg.....	Acid reducer
Calcium carbonate 800 mg.....	Antacid
Magnesium hydroxide 165 mg.....	Antacid

Use relieves heartburn associated with acid indigestion and sour stomach

Warnings
Allergy alert: Do not use if you are allergic to famotidine or other acid reducers


Do not use
• if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.
• with other acid reducers

Distributed by: Kaneva Brands LLC
Sumner, NJ 07901
©Kaneva Brands LLC 2024

Drug Facts (continued)	
Ask a doctor before use if you have	<ul style="list-style-type: none"> had heartburn over 2 months. This may be a sign of a more serious condition. heartburn with lightheadedness, sweating, or dizziness chest pain or shoulder pain with shortness of breath, sweating, pain spreading to arms, neck or shoulders; or lightheadedness frequent chest pain frequent wheezing, particularly with heartburn unexplained weight loss nausea or vomiting stomach pain kidney disease
Ask a doctor or pharmacist before use if you are taking a prescription drug. Antacids and acid reducers may interact with certain prescription drugs.	
Stop use and ask a doctor if	<ul style="list-style-type: none"> your heartburn continues or worsens you need to take this product for more than 14 days
If pregnant or breast-feeding, ask a health professional before use.	
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)	
Directions	<ul style="list-style-type: none"> adults and children 12 years and over: <ul style="list-style-type: none"> do not swallow tablet whole: chew completely to relieve symptoms, chew 1 tablet before swallowing do not use more than 2 chewable tablets in 24 hours children under 12 years: ask a doctor

Net Content Statement		NET CONTENT	AREA OF PDP
PDP Area: (width x height)	5.9 in ²	1/16 IN.	5 SQ. IN. OR LESS
Net Content Size:	0.076"	1/8 IN.	MORE THAN 5 SQ. IN. BUT NOT MORE THAN 25 SQ. IN.
		3/16 IN.	MORE THAN 25 SQ. IN. BUT NOT MORE THAN 100 SQ. IN.
		1/4 IN.	MORE THAN 100 SQ. IN. BUT NOT MORE THAN 400 SQ. IN.
		1/2 IN.	OVER 400 SQ. IN.

Drug Facts Format Information			
Drug Facts:	8.50 pt	Subheadings/Body Text:	6.00 pt
Drug Facts (continued):	8.00 pt	Bullets:	5.00 pt
Headings:	8.00 pt	Leading:	6.50 pt
Barlines:	2.5 pt	Horizontal Scale:	85/100%
Hairlines:	0.5 pt	Tracking:	0



7 4337 00 00 21 7

NINNNN
YYYYMM
0000000

Drug Facts (continued)

Other information

- each tablet contains: calcium 200 mg, magnesium 70 mg
- read the directions and warnings before use
- read the bottle. It contains important information.
- store at 20°-25°C (68°-77°F)
- protect from moisture
- do not use if foil inner seal imprinted with "Sealed For Your Safety" is broken or missing

Inactive ingredients cellulose acetate, corn starch, croscopolone, dextrose excipient, FD&C blue no. 1 aluminum lake, flavors, gum arabic, hydroxypropyl cellulose, hypromellose, lactose monohydrate, magnesium stearate, maltodextrin, mineral oil*, sucralose, yellow iron oxide *may contain mineral oil

Questions or comments? call 1-800-755-4000 (toll-free) or 215-273-8755 (collect)


Pepcid Complete

Acid Reducer + Antacid

Famotidine 10 mg per tablet
Calcium carbonate 800 mg per tablet
Magnesium hydroxide 165 mg per tablet

70 Chewable Tablets

Developed by: Kenova Brands LLC
Kenova, NJ 07091
© Kenova Brands LLC 2024



Shown actual size

Tips for Managing Heartburn

- Do not lie flat or bend over after eating
- Do not wear tight-fitting clothing around the stomach
- Do not eat before bedtime
- Raise the head of your bed
- Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol, and certain fruits and vegetables
- Eat slowly and avoid big meals
- If overweight, lose weight
- Quit smoking

VALUE SIZE NDC 16837-991-70

DUAL ACTION

Pepcid Complete

Famotidine 10 mg—Acid reducer
Calcium carbonate 800 mg—Antacid
Magnesium hydroxide 165 mg—Antacid

Just One Tablet!

Relieves Heartburn Due to Acid Indigestion

Cool Mint Flavor

70 Chewable Tablets

actual size

For Complete Directions and Warnings

Do not use if foil inner seal imprinted with "Sealed For Your Safety" is broken or missing

Drug Facts

Active ingredients	Purposes
Famotidine 10 mg	Acid reducer
Calcium carbonate 800 mg	Antacid
Magnesium hydroxide 165 mg	Antacid

Use relieves heartburn associated with acid indigestion and sour stomach

Warnings

Allergy alert: Do not use if you are allergic to famotidine or other acid reducers

Do not use

- if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.
- with other acid reducers

Developed by:
Kenova Brands LLC
Kenova, NJ 07091
© Kenova Brands LLC 2024

Drug Facts (continued)

Ask a doctor before use if you have

- bad heartburn over 2 months. This may be a sign of a more serious condition.
- heartburn with lightheadedness, sweating, or dizziness
- chest pain or shoulder pain with shortness of breath, sweating, pain spreading to arms, neck or shoulders, or lightheadedness
- frequent chest pain
- frequent belching, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain
- kidney disease

Ask a doctor or pharmacist before use if you are taking a prescription drug. Antacids and acid reducers may interact with certain prescription drugs.

Stop use and ask a doctor if

- your heartburn continues or worsens
- you need to take this product for more than 14 days

If pregnant or breast-feeding, ask a health professional before use.


Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions

- adults and children 12 years and over:
 - do not swallow tablet whole; chew completely
 - to relieve symptoms, chew 1 tablet before swallowing
 - do not use more than 2 chewable tablets in 24 hours
- children under 12 years: ask a doctor

Net Content Statement		NET CONTENT	AREA OF PDP
PDP Area: (width x height)	4.09 in ²	1/16 IN.	5 SQ. IN. OR LESS
Net Content Size:	0.057"	1/8 IN.	MORE THAN 5 SQ. IN. BUT NOT MORE THAN 25 SQ. IN.
		3/16 IN.	MORE THAN 25 SQ. IN. BUT NOT MORE THAN 100 SQ. IN.
		1/4 IN.	MORE THAN 100 SQ. IN. BUT NOT MORE THAN 400 SQ. IN.
		1/2 IN.	OVER 400 SQ. IN.

Drug Facts Format Information					
Drug Facts:	8.50 pt	Subheadings/Body Text:	6.00 pt	Barlines:	2.5 pt
Drug Facts (continued):	8.00 pt	Bullets:	5.00 pt	Horizontal Scale:	85%
Headings:	8.00 pt	Leading:	6.50 pt	Tracking:	0



Drug Facts (continued)
Other information
• each tablet contains: calcium 320 mg, magnesium 70 mg
• contains FD&C yellow no. 5 (tartrazine) as a color additive
• read the directions and warnings before use
• read the bottle. It contains important information.
• store at 20°-25°C (68°-77°F)
• protect from moisture
• do not use if foil inner seal imprinted with "Sealed For Your Safety" is broken or missing
Inactive ingredients cellulose acetate, corn starch, corn syrup solids, croscopolone, dextrose excipient, FD&C yellow no. 5 aluminum lake (tartrazine), FD&C yellow no. 6 aluminum lake, flavors, gum arabic, hydroxypropyl cellulose, hypromellose, lactose monohydrate, magnesium stearate, maltodextrin, mineral oil*, saccharose, triacetin *may contain mineral oil
Questions or comments?
call 1-800-755-4008 (toll-free) or 215-273-4755 (collect)

Tips for Managing Heartburn
• Do not lie flat or bend over after eating • Do not wear tight-fitting clothing around the stomach • Do not eat before bedtime • Raise the head of your bed • Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol, and certain fruits and vegetables • Eat slowly and avoid big meals • If overweight, lose weight • Quit smoking
Pepcid Complete
Acid Reducer + Antacid
Famotidine 10 mg per tablet
Calcium carbonate 800 mg per tablet
25 Chewable Tablets
Distributed by: Karme Brands LLC
Slovak, NJ 07093
©Karme Brands LLC 2024



NDC 10837-246-25

DUAL ACTION

Pepcid Complete

Famotidine 10 mg—Acid reducer
Calcium carbonate 800 mg—Antacid
Magnesium hydroxide 165 mg—Antacid

Just One Tablet!
Relieves Heartburn Due to Acid Indigestion
Tropical Fruit Flavor

25 Chewable Tablets

actual size

For Complete Directions and Warnings

Do not use if foil inner seal imprinted with "Sealed For Your Safety" is broken or missing

Active Ingredients (in each chewable tablet)	Purposes
Famotidine 10 mg	Acid reducer
Calcium carbonate 800 mg	Antacid
Magnesium hydroxide 165 mg	Antacid

Use relieves heartburn associated with acid indigestion and sour stomach

Distributed by: Karme Brands LLC
Slovak, NJ 07093
©Karme Brands LLC 2024

<p>Drug Facts (continued)</p> <p>Warnings Always alert Do not use if you are allergic to famotidine or other acid reducers</p> <p>Do not use • if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor. • with other acid reducers • had heartburn over 2 months. This may be a sign of a more serious condition. • heartburn with lightheadedness, sweating, or dizziness • chest pain or shoulder pain with shortness of breath, sweating, pain spreading to arms, neck or shoulders; or lightheadedness • frequent chest pain • frequent wheezing, particularly with heartburn • unexplained weight loss • hoarseness or vomiting • stomach pain • kidney disease</p>	<p>Drug Facts (continued)</p> <p>Ask a doctor or pharmacist before use if you are taking a prescription drug. Antacids and acid reducers may interact with certain prescription drugs.</p> <p>Stop use and ask a doctor if • your heartburn continues or worsens • you need to take this product for more than 14 days</p> <p>If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-322-1222)</p> <p>Directions • Adults and children 12 years and over: • do not swallow tablet whole; chew completely • to relieve symptoms, chew 1 tablet before swallowing • do not use more than 2 chewable tablets in 24 hours. • children under 12 years: ask a doctor</p>
----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

Net Content Statement		NET CONTENT	AREA OF PDP
PDP Area: (width x height)	5.9 in ²	1/16 IN.	5 SQ. IN. OR LESS
Net Content Size:	0.078"	1/8 IN.	MORE THAN 5 SQ. IN. BUT NOT MORE THAN 25 SQ. IN.
		3/16 IN.	MORE THAN 25 SQ. IN. BUT NOT MORE THAN 100 SQ. IN.
		1/4 IN.	MORE THAN 100 SQ. IN. BUT NOT MORE THAN 400 SQ. IN.
		1/2 IN.	OVER 400 SQ. IN.

Drug Facts Format Information			
Drug Facts:	8.50 pt	Subheadings/Body Text:	6.00 pt
Drug Facts (continued):	8.00 pt	Bullets:	5.00 pt
Headings:	8.00 pt	Leading:	6.50 pt
Barlines:	2.5 pt	Horizontal Scale:	100%
Hairlines:	0.5 pt	Tracking:	0



NNNNNN
EXP YYYMM
0000000

Drug Facts (continued)	
Other information	<ul style="list-style-type: none"> each tablet contains: calcium 320 mg, magnesium 70 mg contains FD&C yellow no. 5 (tartrazine) as a color additive read the directions and warnings before use store at 20°-25°C (68°-77°F) do not use if foil inner seal imprinted with "Sealed For Your Safety" is broken or missing
Inactive ingredients	cellulose acetate, corn starch, corn syrup solids, croscopolone, dextrose excipient, FD&C yellow no. 5 aluminum lake (tartrazine), FD&C yellow no. 6 aluminum lake, flavors, gum arabic, hydroxypropyl cellulose, hypromellose, lactose monohydrate, magnesium stearate, methylcellulose, mineral oil, sucralose, triacetin
Questions or comments?	call 1-800-755-4000 (toll-free) or 215-273-8755 (collect)

Pepcid Complete
Acid Reducer + Antacid
Famotidine 10 mg per tablet
Calcium carbonate 800 mg per tablet
Magnesium hydroxide 165 mg per tablet
50 Chewable Tablets



Shown actual size

- Tips for Managing Heartburn**
- Do not lie flat or bend over after eating
 - Do not wear tight-fitting clothing around the stomach
 - Do not eat before bedtime
 - Raise the head of your bed
 - Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol, and certain fruits and vegetables
 - Eat slowly and avoid big meals
 - If overweight, lose weight
 - Quit smoking

DUAL ACTION NDC 16837-248-50

Pepcid Complete

Famotidine 10 mg—Acid reducer
Calcium carbonate 800 mg—Antacid
Magnesium hydroxide 165 mg—Antacid

Just One Tablet!
Believes Heartburn Due to Acid Indigestion
Tropical Fruit Flavor

50 Chewable Tablets

actual size

For Complete Directions and Warnings

Do not use if foil inner seal imprinted with "Sealed For Your Safety" is broken or missing

Drug Facts

Active ingredients Purposes (in each chewable tablet)

Famotidine 10 mg Acid reducer
Calcium carbonate 800 mg Antacid
Magnesium hydroxide 165 mg Antacid

Use relieves heartburn associated with acid indigestion and sour stomach

Warnings
Allergy alert: Do not use if you are allergic to famotidine or other acid reducers

Do not use

- if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.
- with other acid reducers

Distributed by: Kenova Brands LLC
Summit, NJ 07901
©Kenova Brands LLC 2024

Drug Facts (continued)	
Ask a doctor before use if you have	<ul style="list-style-type: none"> had heartburn over 3 months. This may be a sign of a more serious condition. heartburn with lightheadedness, sweating, or dizziness chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulder; or lightheadedness frequent chest pain frequent wheezing, particularly with heartburn unexplained weight loss nausea or vomiting stomach pain kidney disease
Ask a doctor or pharmacist before use if you are taking a prescription drug.	Antacids and acid reducers may interact with certain prescription drugs.
Stop use and ask a doctor if	<ul style="list-style-type: none"> your heartburn continues or worsens you need to take this product for more than 14 days
If pregnant or breast-feeding, ask a health professional before use.	
Keep out of reach of children.	In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)
Directions	<ul style="list-style-type: none"> adults and children 12 years and over: <ul style="list-style-type: none"> do not swallow tablet whole; chew completely to relieve symptoms, chew 1 tablet before swallowing do not use more than 2 chewable tablets in 24 hours children under 12 years: ask a doctor

This is a representation of an electronic record that was signed electronically. Following this are manifestations of any and all electronic signatures for this electronic record.

/s/

NUSHIN F TODD
01/17/2025 03:11:51 PM