


Drug Facts Format Information					
Drug Facts:	NA pt	Subheaders/body text:	4.25 pt	Barlines:	NA pt
Drug Facts (continued):	NA pt	Bullets:	4.25 pt	Headlines:	NA pt
Headings:	4.25 pt	Leadings:	4.75 pt	Tracking:	0



**Warning: Stop Use** if you experience severe stomach pain, persistent vomiting, or difficulty swallowing. Stop use if you experience severe dizziness, lightheadedness, or fainting. Stop use if you experience severe constipation or diarrhea. Stop use if you experience severe headache or dizziness. Stop use if you experience severe nausea or vomiting. Stop use if you experience severe bloating or gas. Stop use if you experience severe heartburn or acid reflux. Stop use if you experience severe indigestion or stomach pain. Stop use if you experience severe constipation or diarrhea. Stop use if you experience severe dizziness, lightheadedness, or fainting. Stop use if you experience severe headache or dizziness. Stop use if you experience severe nausea or vomiting. Stop use if you experience severe bloating or gas. Stop use if you experience severe heartburn or acid reflux. Stop use if you experience severe indigestion or stomach pain.

**Directions:** Chew 2 tablets 4 times a day after meals and at bedtime. Do not take more than 8 tablets in 24 hours. Do not take more than 2 weeks of treatment without consulting your doctor. Do not take if you are pregnant or breastfeeding. Do not take if you are taking other medications. Do not take if you have a history of stomach ulcers or other stomach conditions. Do not take if you have a history of kidney or liver disease. Do not take if you have a history of heart disease or high blood pressure. Do not take if you have a history of seizures or other neurological conditions. Do not take if you have a history of allergic reactions to any of the ingredients. Do not take if you are allergic to any of the ingredients. Do not take if you are taking other medications. Do not take if you have a history of stomach ulcers or other stomach conditions. Do not take if you have a history of kidney or liver disease. Do not take if you have a history of heart disease or high blood pressure. Do not take if you have a history of seizures or other neurological conditions. Do not take if you have a history of allergic reactions to any of the ingredients. Do not take if you are allergic to any of the ingredients.

**Other information:** Contains 30 chewable tablets. Each tablet contains 375 mg of famotidine and 10 mg of ranitidine. Do not take if you are pregnant or breastfeeding. Do not take if you are taking other medications. Do not take if you have a history of stomach ulcers or other stomach conditions. Do not take if you have a history of kidney or liver disease. Do not take if you have a history of heart disease or high blood pressure. Do not take if you have a history of seizures or other neurological conditions. Do not take if you have a history of allergic reactions to any of the ingredients. Do not take if you are allergic to any of the ingredients.

**Contains:** Famotidine, Ranitidine, and other inactive ingredients.

**Contains 30 Chewable Tablets.**


**© 2010 Watson Pharmaceuticals, Inc.**





Net Content Statement		NET CONTENT	AREA OF PDP
PDP Area: (width x height)	4.09 in <sup>2</sup>	1/16 IN.	5 SQ. IN. OR LESS
Net Content Size:	0.057"	1/8 IN.	MORE THAN 5 SQ. IN. BUT NOT MORE THAN 25 SQ. IN.
		3/16 IN.	MORE THAN 25 SQ. IN. BUT NOT MORE THAN 100 SQ. IN.
		1/4 IN.	MORE THAN 100 SQ. IN. BUT NOT MORE THAN 400 SQ. IN.
		1/2 IN.	OVER 400 SQ. IN.

Drug Facts Format Information					
Drug Facts:	8.50 pt	Subheadings/Body Text:	6.00 pt	Barlines:	2.5 pt
Drug Facts (continued):	8.00 pt	Bullets:	5.00 pt	Hairlines:	0.5 pt
Headings:	8.00 pt	Leading:	6.50 pt	Tracking:	0
				Horizontal Scale:	85%



NNNNNN  
 YYYMM  
 EXP  
 30059923

**Drug Facts (continued)**  
**Other information**  
 • each tablet contains: calcium 320 mg, magnesium 70 mg  
 • read the directions and warnings before use  
 • read the bottle. It contains important information.  
 • store at 20°-25°C (68°-77°F)  
 • protect from moisture  
 • do not use if foil inner seal imprinted with "Sealed For Your Safety" is broken or missing  
**Inactive ingredients** cellulose acetate, corn starch, croscopolone, DSC red no. 7, calcium lake, dextrose excipient, FD&C blue no. 1 aluminum lake, FD&C red no. 40 aluminum lake, fibers, gum arabic, hydroxypropyl cellulose, hypromellose, lactose monohydrate, magnesium stearate, polydextrose, mineral oil\*, modified starch, sucralose  
 \*may contain mineral oil  
**Questions or comments?**  
 call 1-800-755-4008 (toll-free) or 215-273-8755 (collect)

**Tips for Managing Heartburn**  
 • Do not lie flat or bend over after eating  
 • Do not wear tight-fitting clothing around the stomach  
 • Do not eat before bedtime  
 • Raise the head of your bed  
 • Eat slowly and avoid big meals  
 • If overweight, lose weight  
 • Quit smoking  
 • Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol, and certain fruits and vegetables



**Pepcid Complete**  
 Acid Reducer + Antacid  
 25 Chewable Tablets  
 Distributed by Kenue Brands LLC, Summit, NJ 07901  
 © Kenue Brands LLC 2025

NDC 18437-298-25

**DUAL ACTION**

# Pepcid Complete

Famotidine 10 mg—Acid reducer  
 Calcium carbonate 800 mg—Antacid  
 Magnesium hydroxide 165 mg—Antacid

**Just One Tablet!**  
 Relieves Heartburn Due to Acid Indigestion  
 Berry Flavor

**25 Chewable Tablets**

actual size

**For Complete Directions and Warnings**

Do not use if foil inner seal imprinted with "Sealed For Your Safety" is broken or missing.

Active ingredients (in each chewable tablet)	Purposes
Famotidine 10 mg	Acid reducer
Calcium carbonate 800 mg	Antacid
Magnesium hydroxide 165 mg	Antacid

**Use** relieves heartburn associated with acid indigestion and sour stomach

Distributed by Kenue Brands LLC, Summit, NJ 07901  
 © Kenue Brands LLC 2025

<p><b>Drug Facts (continued)</b></p> <p><b>Warnings</b>  <b>Allergy alert:</b> Do not use if you are allergic to famotidine or other acid reducers</p> <p><b>Do not use</b>            • if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.            • with other acid reducers</p> <p><b>Ask a doctor before use if you have</b>            • had heartburn over 3 months. This may be a sign of a more serious condition.            • heartburn with <b>lightheadedness, sweating, or dizziness</b>            • chest pain or shoulder pain with shortness of breath, sweating, pain spreading to arms, neck or shoulders, or lightheadedness            • frequent chest pain            • frequent wheezing, particularly with heartburn            • unexplained weight loss            • stomach pain</p>	<p><b>Drug Facts (continued)</b></p> <p><b>Ask a doctor or pharmacist before use if you are taking a prescription drug.</b> Antacids and acid reducers may interact with certain prescription drugs.</p> <p><b>Stop use and ask a doctor if</b>            • your heartburn continues or worsens            • you need to take this product for more than 14 days</p> <p><b>If pregnant or breast-feeding,</b> ask a health professional before use. <b>Keep out of reach of children.</b> In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)</p> <p><b>Directions</b>            • adults and children 12 years and over:            • do not swallow tablet whole; <b>chew completely</b>            • to relieve symptoms, <b>chew 1 tablet</b> before swallowing            • do not use more than 2 chewable tablets in 24 hours            • children under 12 years: ask a doctor.</p>
-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

Net Content Statement		NET CONTENT	AREA OF PDP
PDP Area: (width x height)	5.9 in <sup>2</sup>	1/8 IN.	5 SQ. IN. OR LESS
Net Content Size:	0.078"	1/8 IN.	MORE THAN 5 SQ. IN. BUT NOT MORE THAN 25 SQ. IN.
		3/16 IN.	MORE THAN 25 SQ. IN. BUT NOT MORE THAN 100 SQ. IN.
		1/4 IN.	MORE THAN 100 SQ. IN. BUT NOT MORE THAN 400 SQ. IN.
		1/2 IN.	OVER 400 SQ. IN.

Drug Facts Format Information			
Drug Facts:	8.50 pt	Subheadings/Body Text:	6.00 pt
Drug Facts (continued):	8.00 pt	Bullets:	5.00 pt
Headings:	8.00 pt	Barlines:	2.5 pt
		Hairlines:	0.5 pt
		Horizontal Scale:	85/100%
		Tracking:	0



NNNNNN  
EXP YYY/MM

30059924

Drug Facts (continued)
<b>Other information</b>
• each tablet contains: calcium 200 mg, magnesium 70 mg
• read the directions and warnings before use
• read the bottle. It contains important information.
• store at 20°-25°C (68°-77°F) — protect from moisture
• do not use if foil inner seal imprinted with "Sealed For Your Safety" is broken or missing
<b>Inactive ingredients:</b> cellulose acetate, corn starch, croscopolidone, DSC red no. 7 calcium lake, dextrose excipient, FD&C blue no. 1 aluminum lake, FD&C red no. 40 aluminum lake, flavors, gum arabic, hydroxypropyl cellulose, hypromellose, lactose monohydrate, magnesium stearate, maltodextrin, mineral oil*, modified starch, sucralose *may contain mineral oil
<b>Questions or comments?</b> call 1-800-755-4008 (toll-free) or 215-273-8755 (collect)

**Pepcid Complete**  
Acid Reducer + Antacid  
50 Chewable Tablets

**Tips for Managing Heartburn**

- Do not lie flat or bend over after eating
- Do not wear tight-fitting clothing around the stomach
- Do not eat before bedtime
- Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol, and certain fruits and vegetables
- Raise the head of your bed
- Eat slowly and avoid big meals
- If overweight, lose weight
- Quit smoking

NDC 15837-298-50

# DUAL ACTION Pepcid Complete

Famotidine 10 mg—Acid reducer  
Calcium carbonate 800 mg—Antacid  
Magnesium hydroxide 165 mg—Antacid

**Just One Tablet!**  
Relieves Heartburn Due to Acid Indigestion  
Berry Flavor

**50 Chewable Tablets**

actual size

**For Complete Directions and Warnings**

Do not use if foil inner seal imprinted with "Sealed For Your Safety" is broken or missing.

**Drug Facts**

Active ingredients	Purposes (in each chewable tablet)
Famotidine 10 mg	Acid reducer
Calcium carbonate 800 mg	Antacid
Magnesium hydroxide 165 mg	Antacid

**Use** relieves heartburn associated with acid indigestion and sour stomach.

**Warnings**  
Always alert. Do not use if you are allergic to famotidine or other acid reducers.

**Do not use**

- if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.
- with other acid reducers


Distributed by: Kenvue Brands LLC, Summit, NJ 07901  
© Kenvue Brands LLC 2025

Drug Facts (continued)
<b>Ask a doctor before use if you have</b>
• had heartburn over 3 months. This may be a sign of a more serious condition.
• heartburn with lightheadedness, sweating, or dizziness
• chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
• frequent chest pain
• frequent wheezing, particularly with heartburn
• unexplained weight loss
• nausea or vomiting
• stomach pain
• kidney disease
<b>Ask a doctor or pharmacist before use if you are taking a prescription drug.</b> Antacids and acid reducers may interact with certain prescription drugs.
<b>Stop use and ask a doctor if</b>
• your heartburn continues or worsens
• you need to take this product for more than 14 days
<b>If pregnant or breast-feeding,</b> ask a health professional before use.
<b>Keep out of reach of children.</b> In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)
<b>Directions</b>
• adults and children 12 years and over:
• do not swallow tablet whole; chew completely
• to relieve symptoms, chew 1 tablet before swallowing
• do not use more than 2 chewable tablets in 24 hours
• children under 12 years: ask a doctor

Net Content Statement		NET CONTENT	AREA OF PDP
PDP Area: (width x height)	5.9 in <sup>2</sup>	1/16 IN.	5 SQ. IN. OR LESS
Net Content Size:	0.076"	1/8 IN.	MORE THAN 5 SQ. IN. BUT NOT MORE THAN 25 SQ. IN.
		3/16 IN.	MORE THAN 25 SQ. IN. BUT NOT MORE THAN 100 SQ. IN.
		1/4 IN.	MORE THAN 100 SQ. IN. BUT NOT MORE THAN 400 SQ. IN.
		1/2 IN.	OVER 400 SQ. IN.

Drug Facts Format Information					
Drug Facts:	8.50 pt	Subheadings/Body Text:	6.00 pt	Barlines:	2.5 pt
Drug Facts (continued):	8.00 pt	Bullets:	5.00 pt	Hairlines:	0.5 pt
Headings:	8.00 pt	Leading:	6.50 pt	Horizontal Scale:	85/100%
				Tracking:	0

7 1 6 8 7 2 9 1 7 2 1 5



EXP YYYYMM  
NNNNNN

30057644

**Drug Facts (continued)**

**Other information**

- each tablet contains: calcium 220 mg, magnesium 70 mg
- read the directions and warnings before use
- read the bottle. It contains important information.
- store at 20°-25° C (68°-77° F)
  - protect from moisture
- do not use if foil inner seal imprinted with "Sealed For Your Safety" is broken or missing

**Inactive ingredients** cellulose acetate, corn starch, croscopolone, D&C red no. 7 calcium lake, dextrose excipient, FD&C blue no. 1 aluminum lake, FD&C red no. 40 aluminum lake, flavors, gum arabic, hydroxypropyl cellulose, hypromellose, lactose monohydrate, magnesium stearate, methylcellulose, mineral oil, modified starch, sucralose  
\*may contain mineral oil

**Questions or comments?** call 1-800-755-4008 (toll-free) or 215-273-8755 (collect)

**Tips for Managing Heartburn**

- Do not lie flat or bend over after eating
- Do not wear tight-fitting clothing around the stomach
- Do not eat before bedtime
- Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol, and certain fruits and vegetables
- Raise the head of your bed
- Eat slowly and avoid big meals
- If overweight, lose weight
- Quit smoking

**Pepcid Complete**  
Acid Reducer + Antacid  
70 Chewable Tablets

Distributed by: Kenvue Brands LLC, Summit, NJ 07901  
© Kenvue Brands LLC 2025

VALUE SIZE

NDC 19237-298-70

**DUAL ACTION**

# Pepcid Complete

Famotidine 10 mg—Acid reducer  
Calcium carbonate 800 mg—Antacid  
Magnesium hydroxide 165 mg—Antacid

**Just One Tablet!**  
Relieves Heartburn Due to Acid Indigestion  
Berry Flavor

**70 Chewable Tablets**

actual size

**For Complete Directions and Warnings**

Do not use if foil inner seal imprinted with "Sealed For Your Safety" is broken or missing.

**Drug Facts**

Active ingredients (in each chewable tablet)	Purposes
Famotidine 10 mg	Acid reducer
Calcium carbonate 800 mg	Antacid
Magnesium hydroxide 165 mg	Antacid

**Use** relieves heartburn associated with acid indigestion and sour stomach

**Warnings**  
Allergy alert: Do not use if you are allergic to famotidine or other acid reducers

**Do not use**

- if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.
- with other acid reducers

Distributed by: Kenvue Brands LLC, Summit, NJ 07901  
© Kenvue Brands LLC 2025  
www.kenvuepeppid.com

**Drug Facts (continued)**

**Ask a doctor before use if you have**

- had heartburn over 3 months. This may be a sign of a more serious condition.
- heartburn with lightheadedness, sweating, or dizziness
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
- frequent chest pain
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain
- kidney disease

**Ask a doctor or pharmacist before use if you are** taking a prescription drug. Antacids and acid reducers may interact with certain prescription drugs.

**Stop use and ask a doctor if**

- your heartburn continues or worsens
- you need to take this product for more than 14 days

**If pregnant or breast-feeding,** ask a health professional before use.  
**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

**Directions**

- adults and children 12 years and over:
  - do not swallow tablet whole: chew completely
  - to relieve symptoms, chew 1 tablet before swallowing
  - do not use more than 2 chewable tablets in 24 hours
- children under 12 years: ask a doctor

Net Content Statement		NET CONTENT	AREA OF PDP
PDP Area: (width x height)	7.34 in <sup>2</sup>	1/16 IN.	5 SQ. IN. OR LESS
Net Content Size:	0.075"	1/8 IN.	MORE THAN 5 SQ. IN. BUT NOT MORE THAN 25 SQ. IN.
		3/16 IN.	MORE THAN 25 SQ. IN. BUT NOT MORE THAN 100 SQ. IN.
		1/4 IN.	MORE THAN 100 SQ. IN. BUT NOT MORE THAN 400 SQ. IN.
		1/2 IN.	OVER 400 SQ. IN.

Drug Facts Format Information							
Drug Facts:	8.50 pt	Subheadings/Body Text:	6.00 pt	Barlines:	2.5 pt	Horizontal Scale:	95/100%
Drug Facts (continued):	8.00 pt	Bullets:	5.00 pt	Hairlines:	0.5 pt	Tracking:	0
Headings:	8.00 pt	Leading:	6.50 pt				



EXP. NNNNNN  
YYYY/MM

30059925

<b>Drug Facts (continued)</b>
<b>Other information</b>
<ul style="list-style-type: none"> <li>each tablet contains: calcium 320 mg, magnesium 70 mg</li> <li>read the directions and warnings before use</li> <li>read the bottle. It contains important information.</li> <li>store at 20°-25° C (68°-77° F)</li> <li>protect from moisture</li> <li>do not use if foil inner seal imprinted with "Sealed For Your Safety" is broken or missing</li> </ul>
<b>Inactive ingredients</b> cellulose acetate, corn starch, croscopolone, D&C red no. 7 calcium lake, dextrose excipient, FD&C blue no. 1 aluminum lake, FD&C red no. 40 aluminum lake, flavors, gum arabic, hydroxypropyl cellulose, hypromellose, lactose monohydrate, magnesium stearate, maltodextrin, mineral oil, modified starch, sucralose
*may contain mineral oil
<b>Questions or comments?</b> call 1-800-755-4008 (toll-free) or 215-273-8795 (collect)

**Pepcid Complete**  
Acid Reducer + Antacid  
100 Chewable Tablets

Distributed by: Kenvue Brands LLC, Summit, NJ 07901  
© Kenvue Brands LLC 2025

- Tips for Managing Heartburn**
- Do not lie flat or bend over after eating
  - Do not wear tight-fitting clothing around the stomach
  - Do not eat before bedtime
  - Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol, and certain fruits and vegetables
  - Raise the head of your bed
  - Eat slowly and avoid big meals
  - If overweight, lose weight
  - Quit smoking

NDC 16837-296-12

**DUAL ACTION**

# Pepcid Complete

Famotidine 10 mg—Acid reducer  
Calcium carbonate 300 mg—Antacid  
Magnesium hydroxide 165 mg—Antacid

**Just One Tablet!**  
Relieves Heartburn Due to Acid Indigestion  
Berry Flavor

**100 Chewable Tablets**

actual size

**For Complete Directions and Warnings**  
Do not use if foil inner seal imprinted with "Sealed For Your Safety" is broken or missing.

Drug Facts	Purposes
<b>Active ingredients (in each chewable tablet)</b>	
Famotidine 10 mg..... Acid reducer	
Calcium carbonate 300 mg..... Antacid	
Magnesium hydroxide 165 mg..... Antacid	
<b>Use</b>	relieves heartburn associated with acid indigestion and sour stomach
<b>Warnings</b>	
<b>Allergy alert:</b> Do not use if you are allergic to famotidine or other acid reducers	
<b>Do not use</b>	<ul style="list-style-type: none"> <li>if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.</li> <li>with other acid reducers</li> </ul>

Distributed by: Kenvue Brands LLC, Summit, NJ 07901  
© Kenvue Brands LLC 2025

<b>Drug Facts (continued)</b>
<b>Ask a doctor before use if you have</b>
<ul style="list-style-type: none"> <li>had heartburn over 3 months. This may be a sign of a more serious condition.</li> <li>heartburn with lightheadedness, sweating, or dizziness</li> <li>chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness</li> <li>frequent chest pain</li> <li>frequent wheezing, particularly with heartburn</li> <li>unexplained weight loss</li> <li>nausea or vomiting</li> <li>stomach pain</li> <li>kidney disease</li> </ul>
<b>Ask a doctor or pharmacist before use if you are taking a prescription drug.</b> Antacids and acid reducers may interact with certain prescription drugs.
<b>Stop use and ask a doctor if</b>
<ul style="list-style-type: none"> <li>your heartburn continues or worsens</li> <li>you need to take this product for more than 14 days</li> </ul>
<b>If pregnant or breast-feeding,</b> ask a health professional before use.
<b>Keep out of reach of children.</b> In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)
<b>Directions</b>
<ul style="list-style-type: none"> <li>adults and children 12 years and over: <ul style="list-style-type: none"> <li>do not swallow tablet whole: chew completely</li> <li>to relieve symptoms, chew 1 tablet before swallowing</li> <li>do not use more than 2 chewable tablets in 24 hours</li> </ul> </li> <li>children under 12 years: ask a doctor</li> </ul>

Net Content Statement		NET CONTENT	AREA OF PDP
PDP Area: (width x height)	4.09 in <sup>2</sup>	1/16 IN.	5 SQ. IN. OR LESS
Net Content Size:	0.057"	1/8 IN.	MORE THAN 5 SQ. IN. BUT NOT MORE THAN 25 SQ. IN.
		3/16 IN.	MORE THAN 25 SQ. IN. BUT NOT MORE THAN 100 SQ. IN.
		1/4 IN.	MORE THAN 100 SQ. IN. BUT NOT MORE THAN 400 SQ. IN.
		1/2 IN.	OVER 400 SQ. IN.

Drug Facts Format Information			
Drug Facts:	8.50 pt	Subheadings/Body Text:	6.00 pt
Drug Facts (continued):	8.00 pt	Bullets:	5.00 pt
Headings:	8.00 pt	Leading:	6.50 pt
		Barlines:	2.5 pt
		Hairlines:	0.5 pt
		Horizontal Scale:	85%
		Tracking:	0



**Drug Facts (continued)**

**Other information**

- each tablet contains calcium 120 mg, magnesium 70 mg
- read the directions and warnings before use
- read the bottle. It contains important information.
- store at 20°-25°C (68°-77°F)
- protect from moisture
- do not use if foil inner seal imprinted with "Sealed For Your Safety" is broken or missing

**Inactive ingredients** cellulose acetate, corn starch, croscopolone, dextrose excipient, FD&C blue no. 1 aluminum lake, flavors, gum arabic, hydroxypropyl cellulose, hypromellose, lactose monohydrate, magnesium stearate, microdextrin, mineral oil\*, sucralose, yellow iron oxide \*may contain mineral oil

**Questions or comments?**  
call 1-800-755-4008 (toll-free) or 215-273-8755 (collect)

**Tips for Managing Heartburn**

- Do not lie flat or bend over after eating
- Do not wear tight-fitting clothing around the stomach
- Do not eat before bedtime
- Raise the head of your bed
- Eat slowly and avoid big meals
- If overweight, lose weight
- Quit smoking
- Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolates, caffeine, alcohol, and certain fruits and vegetables

**Pepcid Complete**  
Acid Reducer + Antacid  
25 Chewable Tablets

Distributed by Kenvue Brands LLC, Summit, NJ 07901  
© Kenvue Brands LLC 2025



NDC 18827-891-45

**DUAL ACTION**

# Pepcid Complete

Famotidine 10 mg - Acid reducer  
Calcium carbonate 800 mg - Antacid  
Magnesium hydroxide 165 mg - Antacid

**Just One Tablet!**  
Relieves Heartburn Due to Acid Indigestion  
Cool Mint Flavor

25 Chewable Tablets

actual size

**For Complete Directions and Warnings**

Do not use if foil inner seal imprinted with "Sealed For Your Safety" is broken or missing

Active ingredients (in each chewable tablet)	Purposes
Famotidine 10 mg.....	Acid reducer
Calcium carbonate 800 mg.....	Antacid
Magnesium hydroxide 165 mg.....	Antacid

**Use** relieves heartburn associated with acid indigestion and sour stomach

Distributed by Kenvue Brands LLC, Summit, NJ 07901  
© Kenvue Brands LLC 2025

<p><b>Drug Facts (continued)</b></p> <p><b>Warnings</b> Allergy alert: Do not use if you are allergic to famotidine or other acid reducers</p> <p><b>Do not use</b></p> <ul style="list-style-type: none"> <li>if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.</li> <li>with other acid reducers</li> </ul> <p><b>Ask a doctor before use if you have</b></p> <ul style="list-style-type: none"> <li>had heartburn over 3 months. This may be a sign of a more serious condition.</li> <li>heartburn with <b>lightheadedness, sweating, or dizziness</b></li> <li>chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness</li> <li>frequent <b>chest pain</b></li> <li>frequent wheezing, particularly with heartburn</li> <li>unexplained weight loss</li> <li>nausea or vomiting</li> <li>stomach pain</li> <li>kidney disease</li> </ul>	<p><b>Drug Facts (continued)</b></p> <p><b>Ask a doctor or pharmacist before use if you are taking a prescription drug.</b> Antacids and acid reducers may interact with certain prescription drugs.</p> <p><b>Stop use and ask a doctor if</b></p> <ul style="list-style-type: none"> <li>your heartburn continues or worsens</li> <li>you need to take this product for more than 14 days.</li> </ul> <p><b>If pregnant or breast-feeding,</b> ask a health professional before use.</p> <p><b>Keep out of reach of children.</b> In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)</p> <p><b>Directions</b></p> <ul style="list-style-type: none"> <li>adults and children 12 years and over:</li> <li>do not swallow tablet whole; <b>chew completely</b></li> <li>to relieve symptoms, <b>chew 1 tablet</b> before swallowing</li> <li>do not use more than 2 chewable tablets in 24 hours</li> <li>children under 12 years: ask a doctor</li> </ul>
--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

Net Content Statement		NET CONTENT	AREA OF PDP
PDP Area: (width x height) <b>5.93 in<sup>2</sup></b>		1/16 IN.	5 SQ. IN. OR LESS
Net Content Size: <b>0.078"</b>		1/8 IN.	MORE THAN 5 SQ. IN. BUT NOT MORE THAN 25 SQ. IN.
		3/16 IN.	MORE THAN 25 SQ. IN. BUT NOT MORE THAN 100 SQ. IN.
		1/4 IN.	MORE THAN 100 SQ. IN. BUT NOT MORE THAN 400 SQ. IN.
		1/2 IN.	OVER 400 SQ. IN.

Drug Facts Format Information					
Drug Facts:	8.50 pt	Subheadings/Body Text:	6.00 pt	Barlines:	2.5 pt
Drug Facts (continued):	8.00 pt	Bullets:	5.00 pt	Hairlines:	0.5 pt
Headings:	8.00 pt	Leading:	6.50 pt	Tracking:	0
				Horizontal Scale:	85/100%



**Drug Facts (continued)**

**Other information**

- each tablet contains: calcium 320 mg, magnesium 70 mg
- read the directions and warnings before use
- read the bottle. It contains important information.
- store at 20°-25° C (68°-77° F) protect from moisture
- do not use if foil inner seal imprinted with "Sealed for Your Safety" is broken or missing

**Inactive ingredients:** cellulose acetate, corn starch, croscopolone, dextrose, excipient, FD&C blue no. 1 aluminum lake, flavors, gum arabic, hydroxypropyl cellulose, hypromellose, lactose monohydrate, magnesium stearate, maltodextrin, mineral oil\*, sucralose, yellow iron oxide  
\*may contain mineral oil

**Questions or comments?** call 1-800-795-9008 (toll-free) or 215-273-8795 (collect)

**Pepcid Complete**  
Acid Reducer + Antacid  
50 Chewable Tablets

**Tips for Managing Heartburn**

- Do not lie flat or bend over after eating
- Do not wear tight-fitting clothing around the stomach
- Do not eat before bedtime
- Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol, and certain fruits and vegetables
- Raise the head of your bed
- Eat slowly and avoid big meals
- If overweight, lose weight
- Quit smoking

Distributed by: Kenvee Brands LLC, Summit, NJ 07901  
© Kenvee Brands LLC 2025

NDC 16837-091-06

DUAL ACTION

# Pepcid Complete

Famotidine 10 mg—Acid reducer  
Calcium carbonate 800 mg—Antacid  
Magnesium hydroxide 165 mg—Antacid

Just One Tablet!

Relieves Heartburn Due to Acid Indigestion  
Cool Mint Flavor

50 Chewable Tablets

actual size

**For Complete Directions and Warnings**

**Do not use if foil inner seal imprinted with "Sealed for Your Safety" is broken or missing**

**Drug Facts**

Active ingredients (in each chewable tablet)	Purposes
Famotidine 10 mg.....	Acid reducer
Calcium carbonate 800 mg.....	Antacid
Magnesium hydroxide 165 mg.....	Antacid

**Use** relieves heartburn associated with acid indigestion and sour stomach

**Warnings**

**Allergy alert:** Do not use if you are allergic to famotidine or other acid reducers

**Do not use**

- if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.
- with other acid reducers

Distributed by: Kenvee Brands LLC, Summit, NJ 07901  
© Kenvee Brands LLC 2025

**Drug Facts (continued)**

**Ask a doctor before use if you have**

- had heartburn over 3 months. This may be a sign of a more serious condition.
- heartburn with **lightheadedness, sweating, or dizziness**
- chest pain or shoulder pain with shortness of breath, sweating; pain spreading to arms, neck or shoulders; or lightheadedness
- frequent chest pain
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain
- kidney disease

**Ask a doctor or pharmacist before use if you are taking a prescription drug.** Antacids and acid reducers may interact with certain prescription drugs.

**Stop use and ask a doctor if**

- your heartburn continues or worsens
- you need to take this product for more than 14 days

**If pregnant or breast-feeding, ask a health professional before use.**

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

**Directions**

- adults and children 12 years and over:
  - do not swallow tablet whole; chew completely
  - to relieve symptoms, chew 1 tablet before swallowing
  - do not use more than 2 chewable tablets in 24 hours
- children under 12 years: ask a doctor

Net Content Statement		NET CONTENT	AREA OF PDP
PDP Area: (width x height)	5.9 in <sup>2</sup>	1/16 IN.	5 SQ. IN. OR LESS
Net Content Size:	0.076"	1/8 IN.	MORE THAN 5 SQ. IN. BUT NOT MORE THAN 25 SQ. IN.
		3/16 IN.	MORE THAN 25 SQ. IN. BUT NOT MORE THAN 100 SQ. IN.
		1/4 IN.	MORE THAN 100 SQ. IN. BUT NOT MORE THAN 400 SQ. IN.
		1/2 IN.	OVER 400 SQ. IN.

Drug Facts Format Information					
Drug Facts:	8.50 pt	Subheadings/Body Text:	6.00 pt	Barlines:	2.5 pt
Drug Facts (continued):	8.00 pt	Bullets:	5.00 pt	Horizontal Scale:	85/100%
Headings:	8.00 pt	Leading:	6.50 pt	Hairlines:	0.5 pt
				Tracking:	0

7  
0 8 5 7 1 8 0 7 2 1 7  
EXP YYYYYM  
NNNNNN  
30057645

**Drug Facts (continued)**

**Other information**

- each tablet contains: calcium 320 mg, magnesium 70 mg
- read the directions and warnings before use
- read the bottle. It contains important information.
- store at 20°-25°C (68°-77°F) • protect from moisture
- do not use if foil inner seal imprinted with "Sealed For Your Safety" is broken or missing

**Inactive ingredients:** cellulose acetate, corn starch, croscopolone, dextrose excipient, FD&C blue no. 1 aluminum lake, flavors, gum arabic, hydroxypropyl cellulose, hypromellose, lactose monohydrate, magnesium stearate, maltodextrin, mineral oil\*, sucralose, yellow iron oxide  
\*may contain mineral oil

**Questions or comments?** call 1-800-755-4008 (toll-free) or 215-273-8755 (collect)

**Pepcid Complete**  
Acid Reducer + Antacid  
70 Chewable Tablets

Distributed by: Kenvue Brands LLC, Summit, NJ 07901  
© Kenvue Brands LLC 2025

**Tips for Managing Heartburn**

- Do not lie flat or bend over after eating
- Do not wear tight-fitting clothing around the stomach
- Do not eat before bedtime
- Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol, and certain fruits and vegetables
- Raise the head of your bed
- Eat slowly and avoid big meals
- If overweight, lose weight
- Quit smoking

**VALUE SIZE** NDC 16837-891-70

**DUAL ACTION**

# Pepcid Complete

Famotidine 10 mg—Acid reducer  
Calcium carbonate 800 mg—Antacid  
Magnesium hydroxide 165 mg—Antacid

**Just One Tablet!**  
Relieves Heartburn Due to Acid Indigestion  
Cool Mint Flavor

**70 Chewable Tablets**

**For Complete Directions and Warnings**

Do not use if foil inner seal imprinted with "Sealed For Your Safety" is broken or missing

Active ingredients	Purposes (in each chewable tablet)
Famotidine 10 mg	Acid reducer
Calcium carbonate 800 mg	Antacid
Magnesium hydroxide 165 mg	Antacid

**Use** relieves heartburn associated with acid indigestion and sour stomach

**Warnings**  
Allergy alert: Do not use if you are allergic to famotidine or other acid reducers

**Do not use**  
if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.

**Do not use**  
with other acid reducers

Distributed by:  
Kenvue Brands LLC, Summit, NJ 07901  
© Kenvue Brands LLC 2025

**Drug Facts (continued)**

**Ask a doctor before use if you have**

- had heartburn over 3 months. This may be a sign of a more serious condition.
- heartburn with lightheadedness, sweating, or dizziness
- chest pain or shoulder pain with shortness of breath, sweating, pain spreading to arms, neck or shoulders; or lightheadedness
- frequent chest pain
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain
- kidney disease

**Ask a doctor or pharmacist before use if you are** taking a prescription drug. Antacids and acid reducers may interact with certain prescription drugs.

**Step use and ask a doctor if**

- your heartburn continues or worsens
- you need to take this product for more than 14 days

**If pregnant or breast-feeding,** ask a health professional before use.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

**Directions**

- adults and children 12 years and over:
  - do not swallow tablet whole: chew completely
  - to relieve symptoms, chew 1 tablet before swallowing
  - do not use more than 2 chewable tablets in 24 hours
- children under 12 years: ask a doctor

Net Content Statement		NET CONTENT	AREA OF PDP
PDP Area: (width x height)		1/16 IN.	5 SQ. IN. OR LESS
Net Content Size:		1/8 IN.	MORE THAN 5 SQ. IN. BUT NOT MORE THAN 25 SQ. IN.
	4.09 in <sup>2</sup>	3/16 IN.	MORE THAN 25 SQ. IN. BUT NOT MORE THAN 100 SQ. IN.
	0.057"	1/4 IN.	MORE THAN 100 SQ. IN. BUT NOT MORE THAN 400 SQ. IN.
		1/2 IN.	OVER 400 SQ. IN.

Drug Facts Format Information			
Drug Facts:	8.50 pt	Subheadings/Body Text:	6.00 pt
Drug Facts (continued):	8.00 pt	Bullets:	5.00 pt
Headings:	8.00 pt	Leading:	6.50 pt
		Barlines:	2.5 pt
		Hairlines:	0.5 pt
		Horizontal Scale:	85%
		Tracking:	0



**Drug Facts (continued)**

**Other information**

- each tablet contains: calcium 320 mg, magnesium 70 mg
- contains FD&C yellow no. 5 (tartrazine) as a color additive
- read the directions and warnings before use
- read the bottle. It contains important information.
- store at 20°-25°C (68°-77°F) protect from moisture
- do not use if foil inner seal imprinted with "Sealed For Your Safety" is broken or missing

**Inactive ingredients** cellulose acetate, corn starch, corn syrup solids, croscopolone, dextrose equivalent, FD&C yellow no. 5 aluminum lake (tartrazine), FD&C yellow no. 8 aluminum lake, flavors, gum arabic, hydroxypropyl cellulose, hypromellose, lactose monohydrate, magnesium stearate, methylcellulose, mineral oil\*, saccharin, triacetin  
\*may contain mineral oil

**Questions or comments?**  
call 1-800-755-4068 (toll-free) or 215-273-8755 (collect)

**Tips for Managing Heartburn**

- Do not lie flat or bend over after eating
- Do not wear tight-fitting clothing around the stomach
- Do not eat before bedtime
- Raise the head of your bed
- Eat slowly and avoid big meals
- If overweight, lose weight
- Quit smoking
- Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol, and certain fruits and vegetables

**Pepcid Complete**  
Acid Reducer • Antacid  
25 Chewable Tablets

Distributed by: Kenvue Brands LLC, Summit, NJ 07901  
© Kenvue Brands LLC 2025



NDC 16837-246-25

**DUAL ACTION**

# Pepcid Complete

Famotidine 10 mg—Acid reducer  
Calcium carbonate 800 mg—Antacid  
Magnesium hydroxide 165 mg—Antacid

**Just One Tablet!**  
Relieves Heartburn, Gas, Acid Indigestion  
Tropical Fruit Flavor

**25 Chewable Tablets**

ACTUAL SIZE

**For Complete Directions and Warnings**

Do not use if foil inner seal imprinted with "Sealed For Your Safety" is broken or missing

Drug Facts	Purposes
<b>Active ingredients (in each chewable tablet)</b>	
Famotidine 10 mg	Acid reducer
Calcium carbonate 800 mg	Antacid
Magnesium hydroxide 165 mg	Antacid

**Use** relieves heartburn associated with acid indigestion and sour stomach

Distributed by: Kenvue Brands LLC, Summit, NJ 07901  
© Kenvue Brands LLC 2025

<b>Drug Facts (continued)</b>	<b>Drug Facts (continued)</b>
<b>Warnings</b> Allergy alert: Do not use if you are allergic to famotidine or other acid reducers <b>Do not use</b> • if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor. • with other acid reducers <b>Ask a doctor before use if you have</b> • had heartburn over 3 months. This may be a sign of a more serious condition. • heartburn with <b>lightheadedness, sweating, or dizziness</b> • chest pain or shoulder pain with shortness of breath, sweating, pain spreading to arms, neck, or shoulders; or lightheadedness • frequent <b>chest pain</b> • frequent wheezing, particularly with heartburn • unexplained weight loss      • nose or vomiting • stomach pain      • kidney disease	<b>Ask a doctor or pharmacist before use if you are taking a prescription drug.</b> Antacids and acid reducers may interact with certain prescription drugs. <b>Stop use and ask a doctor if</b> • your heartburn continues or worsens • you need to take this product for more than 14 days <b>If pregnant or breast-feeding,</b> ask a health professional before use. <b>Keep out of reach of children.</b> In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).
	<b>Directions</b> • adults and children 12 years and over: • <b>do not swallow tablet whole; chew completely</b> • to relieve symptoms, <b>chew 1 tablet</b> before swallowing • do not use more than 2 chewable tablets in 24 hours • children under 12 years: ask a doctor

Net Content Statement		NET CONTENT	AREA OF PDP
PDP Area: (width x height)	5.9 in <sup>2</sup>	1/16 IN.	5 SQ. IN. OR LESS
Net Content Size:	0.078"	1/8 IN.	MORE THAN 5 SQ. IN. BUT NOT MORE THAN 25 SQ. IN.
		3/16 IN.	MORE THAN 25 SQ. IN. BUT NOT MORE THAN 100 SQ. IN.
		1/4 IN.	MORE THAN 100 SQ. IN. BUT NOT MORE THAN 400 SQ. IN.
		1/2 IN.	OVER 400 SQ. IN.

Drug Facts Format Information			
Drug Facts:	8.50 pt	Subheadings/Body Text:	6.00 pt
Drug Facts (continued):	8.00 pt	Bullets:	5.00 pt
Headings:	8.00 pt	Leading:	6.50 pt
		Barlines:	2.5 pt
		Hairlines:	0.5 pt
		Horizontal Scale:	100%
		Tracking:	0



EXP NNNNNN  
YYYYMM  
30059922

Drug Facts (continued)	
<b>Other information</b>	
<ul style="list-style-type: none"> <li>each tablet contains: calcium 320 mg, magnesium 70 mg</li> <li>contains FD&amp;C yellow no. 5 (tartrazine) as a color additive</li> <li>read the directions and warnings before use</li> <li>store at 20°-25°C (68°-77°F)</li> <li>do not use if foil inner seal imprinted with "Sealed For Your Safety" is broken or missing</li> </ul>	<ul style="list-style-type: none"> <li>read the bottle. It contains important information.</li> <li>protect from moisture</li> </ul>
<b>Inactive ingredients</b> cellulose acetate, corn starch, corn syrup solids, croscopolone, dextrose excipient, FD&C yellow no. 5 aluminum lake (tartrazine), FD&C yellow no. 6 aluminum lake, flavors, gum arabic, hydroxypropyl cellulose, hypromellose, lactose monohydrate, magnesium stearate, maltodextrin, mineral oil*, sucralose, triacetin	
*may contain mineral oil	
<b>Questions or comments?</b> call 1-800-765-4008 (toll-free) or 215-273-8785 (collect)	

**Pepcid Complete**  
Acid Reducer - Antacid  
50 Chewable Tablets

Distributed by Kenwa Brands LLC, Summit, NJ 07901  
© Kenwa Brands LLC 2025

- Tips for Managing Heartburn**
- Do not lie flat or bend over after eating
  - Do not wear tight-fitting clothing around the stomach
  - Do not eat before bedtime
  - Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol, and certain fruits and vegetables
  - Raise the head of your bed
  - Eat slowly and avoid big meals
  - If overweight, lose weight
  - Quit smoking

**DUAL ACTION**  
**Pepcid Complete**  
Famotidine 10 mg - Acid reducer  
Calcium carbonate 800 mg - Antacid  
Magnesium hydroxide 165 mg - Antacid

**Just One Tablet!**  
Relieves Heartburn Due to Acid Indigestion  
Tropical Fruit Flavor

50 Chewable Tablets

**For Complete Directions and Warnings**

Do not use if foil inner seal imprinted with "Sealed For Your Safety" is broken or missing

**Drug Facts**

**Active ingredients Purposes (in each chewable tablet)**

Famotidine 10 mg.....Acid reducer  
Calcium carbonate 800 mg.....Antacid  
Magnesium hydroxide 165 mg.....Antacid

**Use** relieves heartburn associated with acid indigestion and sour stomach

**Warnings**  
**Allergy alert:** Do not use if you are allergic to famotidine or other acid reducers

**Do not use**

- if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.
- with other acid reducers.

Distributed by Kenwa Brands LLC, Summit, NJ 07901  
© Kenwa Brands LLC 2025

Drug Facts (continued)	
<b>Ask a doctor before use if you have</b>	
<ul style="list-style-type: none"> <li>had heartburn over 3 months. This may be a sign of a more serious condition.</li> <li>heartburn with lightheadedness, sweating, or dizziness</li> <li>chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness</li> <li>frequent chest pain</li> <li>frequent wheezing, particularly with heartburn</li> <li>unexplained weight loss</li> <li>nausea or vomiting</li> <li>stomach pain</li> <li>kidney disease</li> </ul>	
<b>Ask a doctor or pharmacist before use if you are taking a prescription drug.</b> Antacids and acid reducers may interact with certain prescription drugs.	
<b>Stop use and ask a doctor if</b>	
<ul style="list-style-type: none"> <li>your heartburn continues or worsens</li> <li>you need to take this product for more than 14 days</li> </ul>	
<b>If pregnant or breast-feeding,</b> ask a health professional before use.	
<b>Keep out of reach of children.</b> In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)	
<b>Directions</b>	
<ul style="list-style-type: none"> <li>adults and children 12 years and over: <ul style="list-style-type: none"> <li>do not swallow tablet whole; chew completely</li> <li>to relieve symptoms, chew 1 tablet before swallowing</li> <li>do not use more than 2 chewable tablets in 24 hours</li> </ul> </li> <li>children under 12 years: ask a doctor</li> </ul>	

-----  
**This is a representation of an electronic record that was signed electronically. Following this are manifestations of any and all electronic signatures for this electronic record.**  
-----

/s/  
-----

NUSHIN F TODD  
09/30/2025 02:55:47 PM