

HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use RELENZA safely and effectively. See full prescribing information for RELENZA.

RELENZA® (zanamivir) Inhalation Powder, for oral inhalation
Initial U.S. Approval: 1999

-----**RECENT MAJOR CHANGES**-----

Warnings and Precautions
Neuropsychiatric Events (5.3) February 2008

-----**INDICATIONS AND USAGE**-----

RELENZA, an influenza neuraminidase inhibitor, is indicated for:
Treatment of influenza in patients 7 years of age and older who have been symptomatic for no more than 2 days. (1.1)

Prophylaxis of influenza in patients 5 years of age and older. (1.2)

Important Limitations on Use of RELENZA:

Not recommended for treatment or prophylaxis of influenza in:

- Individuals with underlying airways disease. (5.1)

Not proven effective for:

- Treatment in individuals with underlying airways disease. (1.3)
- Prophylaxis in nursing home residents. (1.3)

Not a substitute for annual influenza vaccination. (1.3)

-----**DOSAGE AND ADMINISTRATION**-----

Indication	Dose
Treatment of Influenza (2.2)	10 mg twice daily for 5 days
Prophylaxis: (2.3)	
Household Setting	10 mg once daily for 10 days
Community Outbreaks	10 mg once daily for 28 days

Note: The 10 mg dose is provided by 2 inhalations (one 5 mg blister per inhalation). (2.1)

-----**DOSAGE FORMS AND STRENGTHS**-----

Four 5 mg blisters of powder on a ROTADISK® for oral inhalation via DISKHALER®. Packaged in carton containing 5 ROTADISKS (total of 10 doses) and 1 DISKHALER inhalation device. (3)

-----**CONTRAINDICATIONS**-----

Do not use in patients with history of allergic reaction to any ingredient of RELENZA, including lactose (which contains milk proteins). (4)

-----**WARNINGS AND PRECAUTIONS**-----

- **Bronchospasm:** Serious, sometimes fatal, cases have occurred. Not recommended in individuals with underlying airways disease. Discontinue RELENZA if bronchospasm or decline in respiratory function develops. (5.1)
- **Allergic Reactions:** Discontinue RELENZA and initiate appropriate treatment if an allergic reaction occurs or is suspected. (5.2)
- **Neuropsychiatric Events:** Patients with influenza, particularly pediatric patients, may be at an increased risk of seizures, confusion, or abnormal behavior early in their illness. Monitor for signs of abnormal behavior. (5.3)
- **High-risk underlying medical conditions:** Safety and effectiveness have not been demonstrated in these patients. (5.4)

-----**ADVERSE REACTIONS**-----

The most common adverse events reported in >1.5% of patients treated with RELENZA and more commonly than in patients treated with placebo are:

- Treatment Studies – sinusitis, dizziness.
- Prophylaxis studies – fever and/or chills, arthralgia and articular rheumatism. (6.1)

To report SUSPECTED ADVERSE REACTIONS, contact GlaxoSmithKline at 1-888-825-5249 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

-----**DRUG INTERACTIONS**-----

- Live attenuated influenza vaccine, intranasal (7):
 - Do not administer until 48 hours following cessation of RELENZA.
 - Do not administer RELENZA until 2 weeks following administration of the live attenuated influenza vaccine, unless medically indicated.

See 17 for PATIENT COUNSELING INFORMATION and FDA-approved patient labeling.

Revised: February 2008
RLZ:3PI

FULL PRESCRIBING INFORMATION: CONTENTS*

<p>1 INDICATIONS AND USAGE</p> <p>1.1 Treatment of Influenza</p> <p>1.2 Prophylaxis of Influenza</p> <p>1.3 Important Limitations on Use of RELENZA</p> <p>2 DOSAGE AND ADMINISTRATION</p> <p>2.1 Dosing Considerations</p> <p>2.2 Treatment of Influenza</p> <p>2.3 Prophylaxis of Influenza</p> <p>3 DOSAGE FORMS AND STRENGTHS</p> <p>4 CONTRAINDICATIONS</p> <p>5 WARNINGS AND PRECAUTIONS</p> <p>5.1 Bronchospasm</p> <p>5.2 Allergic Reactions</p> <p>5.3 Neuropsychiatric Events</p> <p>5.4 Limitations of Populations Studied</p> <p>5.5 Bacterial Infections</p> <p>5.6 Importance of Proper Use of DISKHALER</p> <p>6 ADVERSE REACTIONS</p> <p>6.1 Clinical Trials Experience</p> <p>6.2 Postmarketing Experience</p> <p>7 DRUG INTERACTIONS</p> <p>8 USE IN SPECIFIC POPULATIONS</p> <p>8.1 Pregnancy</p> <p>8.3 Nursing Mothers</p>	<p>8.4 Pediatric Use</p> <p>8.5 Geriatric Use</p> <p>10 OVERDOSAGE</p> <p>11 DESCRIPTION</p> <p>12 CLINICAL PHARMACOLOGY</p> <p>12.1 Mechanism of Action</p> <p>12.3 Pharmacokinetics</p> <p>12.4 Microbiology</p> <p>13 NONCLINICAL TOXICOLOGY</p> <p>13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility</p> <p>14 CLINICAL STUDIES</p> <p>14.1 Treatment of Influenza</p> <p>14.2 Prophylaxis of Influenza</p> <p>16 HOW SUPPLIED/STORAGE AND HANDLING</p> <p>17 PATIENT COUNSELING INFORMATION</p> <p>17.1 Bronchospasm</p> <p>17.2 Concomitant Bronchodilator Use</p> <p>17.3 Neuropsychiatric Events</p> <p>17.4 Instructions for Use</p> <p>17.5 Risk of Influenza Transmission to Others</p> <p>17.6 FDA-Approved Patient Labeling and Instructions for Use</p>
---	--

*Sections or subsections omitted from the full prescribing information are not listed.

1

2 **FULL PRESCRIBING INFORMATION**

3 **1 INDICATIONS AND USAGE**

4 **1.1 Treatment of Influenza**

5 RELENZA is indicated for treatment of uncomplicated acute illness due to influenza A
6 and B virus in adults and pediatric patients 7 years of age and older who have been symptomatic
7 for no more than 2 days.

8 **1.2 Prophylaxis of Influenza**

9 RELENZA is indicated for prophylaxis of influenza in adults and pediatric patients
10 5 years of age and older.

11 **1.3 Important Limitations on Use of RELENZA**

- 12 • RELENZA is not recommended for treatment or prophylaxis of influenza in individuals with
13 underlying airways disease (such as asthma or chronic obstructive pulmonary disease) due to
14 risk of serious bronchospasm [*see Warnings and Precautions (5.1)*].
- 15 • RELENZA has not been proven effective for treatment of influenza in individuals with
16 underlying airways disease.
- 17 • RELENZA has not been proven effective for prophylaxis of influenza in the nursing home
18 setting.
- 19 • RELENZA is not a substitute for early influenza vaccination on an annual basis as
20 recommended by the Centers for Disease Control's Immunization Practices Advisory
21 Committee.
- 22 • There is no evidence for efficacy of zanamivir in any illness caused by agents other than
23 influenza virus A and B.
- 24 • Patients should be advised that the use of RELENZA for treatment of influenza has not been
25 shown to reduce the risk of transmission of influenza to others.

26 **2 DOSAGE AND ADMINISTRATION**

27 **2.1 Dosing Considerations**

- 28 • RELENZA is for administration to the respiratory tract by oral inhalation only, using the
29 DISKHALER device provided.
- 30 • The 10 mg dose is provided by 2 inhalations (one 5 mg blister per inhalation).
- 31 • Patients should be instructed in the use of the delivery system. Instructions should include a
32 demonstration whenever possible. If RELENZA is prescribed for children, it should be used
33 only under adult supervision and instruction, and the supervising adult should first be
34 instructed by a healthcare professional [*see Patient Counseling Information (17.3)*].
- 35 • Patients scheduled to use an inhaled bronchodilator at the same time as RELENZA should
36 use their bronchodilator before taking RELENZA [*see Patient Counseling Information*
37 (*17.2*)].

38 **2.2 Treatment of Influenza**

- 39 • The recommended dose of RELENZA for treatment of influenza in adults and pediatric
40 patients ages 7 years of age and older is 10 mg twice daily (approximately 12 hours apart) for
41 5 days.
42 • Two doses should be taken on the first day of treatment whenever possible provided there is
43 at least 2 hours between doses.
44 • On subsequent days, doses should be about 12 hours apart (e.g., morning and evening) at
45 approximately the same time each day.
46 • The safety and efficacy of repeated treatment courses have not been studied.

47 **2.3 Prophylaxis of Influenza**

48 Household Setting:

- 49 • The recommended dose of RELENZA for prophylaxis of influenza in adults and pediatric
50 patients 5 years of age and older in a household setting is 10 mg once daily for 10 days.
51 • The dose should be administered at approximately the same time each day.
52 • There are no data on the effectiveness of prophylaxis with RELENZA in a household setting
53 when initiated more than 1.5 days after the onset of signs or symptoms in the index case.

54 Community Outbreaks:

- 55 • The recommended dose of RELENZA for prophylaxis of influenza in adults and adolescents
56 in a community setting is 10 mg once daily for 28 days.
57 • The dose should be administered at approximately the same time each day.
58 • There are no data on the effectiveness of prophylaxis with RELENZA in a community
59 outbreak when initiated more than 5 days after the outbreak was identified in the community.
60 • The safety and effectiveness of prophylaxis with RELENZA have not been evaluated for
61 longer than 28 days' duration.

62 **3 DOSAGE FORMS AND STRENGTHS**

63 Four 5 mg blisters of powder on a ROTADISK for oral inhalation via DISKHALER.
64 Packaged in carton containing 5 ROTADISKS (total of 10 doses) and 1 DISKHALER inhalation
65 device [see *How Supplied/Storage and Handling (16)*].

66 **4 CONTRAINDICATIONS**

67 Do not use in patients with history of allergic reaction to any ingredient of RELENZA
68 including lactose (which contains milk proteins) [see *Warnings and Precautions (5.2)*,
69 *Description (11)*].

70 **5 WARNINGS AND PRECAUTIONS**

71 **5.1 Bronchospasm**

72 RELENZA is not recommended for treatment or prophylaxis of influenza in individuals
73 with underlying airways disease (such as asthma or chronic obstructive pulmonary disease).

74 Serious cases of bronchospasm, including fatalities, have been reported during treatment
75 with RELENZA in patients with and without underlying airways disease. Many of these cases
76 were reported during postmarketing and causality was difficult to assess.

77 RELENZA should be discontinued in any patient who develops bronchospasm or decline
78 in respiratory function; immediate treatment and hospitalization may be required.

79 Some patients without prior pulmonary disease may also have respiratory abnormalities
80 from acute respiratory infection that could resemble adverse drug reactions or increase patient
81 vulnerability to adverse drug reactions.

82 Bronchospasm was documented following administration of zanamivir in 1 of 13 patients
83 with mild or moderate asthma (but without acute influenza-like illness) in a Phase I study. In a
84 Phase III study in patients with acute influenza-like illness superimposed on underlying asthma
85 or chronic obstructive pulmonary disease, 10% (24 of 244) of patients on zanamivir and 9% (22
86 of 237) on placebo experienced a greater than 20% decline in FEV₁ following treatment for
87 5 days.

88 If use of RELENZA is considered for a patient with underlying airways disease, the
89 potential risks and benefits should be carefully weighed. If a decision is made to prescribe
90 RELENZA for such a patient, this should be done only under conditions of careful monitoring of
91 respiratory function, close observation, and appropriate supportive care including availability of
92 fast-acting bronchodilators.

93 **5.2 Allergic Reactions**

94 Allergic-like reactions, including oropharyngeal edema, serious skin rashes, and
95 anaphylaxis have been reported in postmarketing experience with RELENZA. RELENZA
96 should be stopped and appropriate treatment instituted if an allergic reaction occurs or is
97 suspected.

98 **5.3 Neuropsychiatric Events**

99 Influenza can be associated with a variety of neurologic and behavioral symptoms which
100 can include events such as seizures, hallucinations, delirium, and abnormal behavior, in some
101 cases resulting in fatal outcomes. These events may occur in the setting of encephalitis or
102 encephalopathy but can occur without obvious severe disease.

103 There have been postmarketing reports (mostly from Japan) of delirium and abnormal
104 behavior leading to injury in patients with influenza who were receiving neuraminidase
105 inhibitors, including RELENZA. Because these events were reported voluntarily during clinical
106 practice, estimates of frequency cannot be made, but they appear to be uncommon based on
107 usage data for RELENZA. These events were reported primarily among pediatric patients and
108 often had an abrupt onset and rapid resolution. The contribution of RELENZA to these events
109 has not been established. Patients with influenza should be closely monitored for signs of
110 abnormal behavior. If neuropsychiatric symptoms occur, the risks and benefits of continuing
111 treatment should be evaluated for each patient.

112 **5.4 Limitations of Populations Studied**

113 Safety and efficacy have not been demonstrated in patients with high-risk underlying
114 medical conditions. No information is available regarding treatment of influenza in patients with
115 any medical condition sufficiently severe or unstable to be considered at imminent risk of
116 requiring inpatient management.

117 **5.5 Bacterial Infections**

118 Serious bacterial infections may begin with influenza-like symptoms or may coexist with
119 or occur as complications during the course of influenza. RELENZA has not been shown to
120 prevent such complications.

121 **5.6 Importance of Proper Use of DISKHALER**

122 Effective and safe use of RELENZA requires proper use of the DISKHALER to inhale
123 the drug. Prescribers should carefully evaluate the ability of young children to use the delivery
124 system if use of RELENZA is considered [*see Use in Specific Populations (8.4)*].

125 **6 ADVERSE REACTIONS**

126 See Warnings and Precautions for information about risk of serious adverse events such
127 as bronchospasm (5.1) and allergic-like reactions (5.2), and for safety information in patients
128 with underlying airways disease (5.1).

129 **6.1 Clinical Trials Experience**

130 Because clinical trials are conducted under widely varying conditions, adverse reaction
131 rates observed in the clinical trials of a drug cannot be directly compared with rates in the
132 clinical trials of another drug and may not reflect the rates observed in practice.

133 The placebo used in clinical studies consisted of inhaled lactose powder, which is also the
134 vehicle for the active drug; therefore, some adverse events occurring at similar frequencies in
135 different treatment groups could be related to lactose vehicle inhalation.

136 Treatment of Influenza: Clinical Trials in Adults and Adolescents: Adverse events
137 that occurred with an incidence $\geq 1.5\%$ in treatment studies are listed in Table 1. This table shows
138 adverse events occurring in patients ≥ 12 years of age receiving RELENZA 10 mg inhaled twice
139 daily, RELENZA in all inhalation regimens, and placebo inhaled twice daily (where placebo
140 consisted of the same lactose vehicle used in RELENZA).

141

142 **Table 1. Summary of Adverse Events $\geq 1.5\%$ Incidence During Treatment in Adults and**
143 **Adolescents**

Adverse Event	RELENZA		Placebo (Lactose Vehicle) (n = 1,520)
	10 mg b.i.d. Inhaled (n = 1,132)	All Dosing Regimens* (n = 2,289)	
Body as a whole			
Headaches	2%	2%	3%
Digestive			
Diarrhea	3%	3%	4%
Nausea	3%	3%	3%
Vomiting	1%	1%	2%
Respiratory			
Nasal signs and symptoms	2%	3%	3%
Bronchitis	2%	2%	3%
Cough	2%	2%	3%
Sinusitis	3%	2%	2%
Ear, nose, and throat infections	2%	1%	2%
Nervous system			
Dizziness	2%	1%	<1%

144 * Includes studies where RELENZA was administered intranasally (6.4 mg 2 to 4 times per day
145 in addition to inhaled preparation) and/or inhaled more frequently (q.i.d.) than the currently
146 recommended dose.

147
148 Additional adverse reactions occurring in less than 1.5% of patients receiving RELENZA
149 included malaise, fatigue, fever, abdominal pain, myalgia, arthralgia, and urticaria.

150 The most frequent laboratory abnormalities in Phase III treatment studies included elevations
151 of liver enzymes and CPK, lymphopenia, and neutropenia. These were reported in similar
152 proportions of zanamivir and lactose vehicle placebo recipients with acute influenza-like illness.

153 *Clinical Trials in Pediatric Patients:* Adverse events that occurred with an incidence
154 $\geq 1.5\%$ in children receiving treatment doses of RELENZA in 2 Phase III studies are listed in
155 Table 2. This table shows adverse events occurring in pediatric patients 5 to 12 years old
156 receiving RELENZA 10 mg inhaled twice daily and placebo inhaled twice daily (where placebo
157 consisted of the same lactose vehicle used in RELENZA).

158

159 **Table 2. Summary of Adverse Events $\geq 1.5\%$ Incidence During Treatment in Pediatric**
160 **Patients***

Adverse Event	RELENZA 10 mg b.i.d. Inhaled (n = 291)	Placebo (Lactose Vehicle) (n = 318)
Respiratory		
Ear, nose, and throat infections	5%	5%
Ear, nose, and throat hemorrhage	<1%	2%
Asthma	<1%	2%
Cough	<1%	2%
Digestive		
Vomiting	2%	3%
Diarrhea	2%	2%
Nausea	<1%	2%

161 * Includes a subset of patients receiving RELENZA for treatment of influenza in a prophylaxis
162 study.

163

164 In 1 of the 2 studies described in Table 2, some additional information is available from
165 children (5 to 12 years old) without acute influenza-like illness who received an investigational
166 prophylaxis regimen of RELENZA; 132 children received RELENZA and 145 children received
167 placebo. Among these children, nasal signs and symptoms (zanamivir 20%, placebo 9%), cough
168 (zanamivir 16%, placebo 8%), and throat/tonsil discomfort and pain (zanamivir 11%, placebo
169 6%) were reported more frequently with RELENZA than placebo. In a subset with chronic
170 pulmonary disease, lower respiratory adverse events (described as asthma, cough, or viral
171 respiratory infections which could include influenza-like symptoms) were reported in 7 of 7
172 zanamivir recipients and 5 of 12 placebo recipients.

173 Prophylaxis of Influenza: Family/Household Prophylaxis Studies: Adverse events
174 that occurred with an incidence of $\geq 1.5\%$ in the 2 prophylaxis studies are listed in Table 3. This
175 table shows adverse events occurring in patients ≥ 5 years of age receiving RELENZA 10 mg
176 inhaled once daily for 10 days.

177

178 **Table 3. Summary of Adverse Events $\geq 1.5\%$ Incidence During 10-Day Prophylaxis Studies**
179 **in Adults, Adolescents, and Children***

Adverse Event	Contact Cases	
	RELENZA (n = 1,068)	Placebo (n = 1,059)
Lower respiratory		
Viral respiratory infections	13%	19%
Cough	7%	9%
Neurologic		
Headaches	13%	14%
Ear, nose, and throat		
Nasal signs and symptoms	12%	12%
Throat and tonsil discomfort and pain	8%	9%
Nasal inflammation	1%	2%
Musculoskeletal		
Muscle pain	3%	3%
Endocrine and metabolic		
Feeding problems (decreased or increased appetite and anorexia)	2%	2%
Gastrointestinal		
Nausea and vomiting	1%	2%
Non-site specific		
Malaise and fatigue	5%	5%
Temperature regulation disturbances (fever and/or chills)	5%	4%

180 * In prophylaxis studies, symptoms associated with influenza-like illness were captured as
181 adverse events; subjects were enrolled during a winter respiratory season during which time
182 any symptoms that occurred were captured as adverse events.

183
184 *Community Prophylaxis Studies:* Adverse events that occurred with an incidence of
185 $\geq 1.5\%$ in 2 prophylaxis studies are listed in Table 4. This table shows adverse events occurring
186 in patients ≥ 5 years of age receiving RELENZA 10 mg inhaled once daily for 28 days.

187

188 **Table 4. Summary of Adverse Events $\geq 1.5\%$ Incidence During 28-Day Prophylaxis Studies**
189 **in Adults, Adolescents, and Children***

Adverse Event	RELENZA (n = 2,231)	Placebo (n = 2,239)
Neurologic		
Headaches	24%	26%
Ear, nose, and throat		
Throat and tonsil discomfort and pain	19%	20%
Nasal signs and symptoms	12%	13%
Ear, nose, and throat infections	2%	2%
Lower respiratory		
Cough	17%	18%
Viral respiratory infections	3%	4%
Musculoskeletal		
Muscle pain	8%	8%
Musculoskeletal pain	6%	6%
Arthralgia and articular rheumatism	2%	<1%
Endocrine and metabolic		
Feeding problems (decreased or increased appetite and anorexia)	4%	4%
Gastrointestinal		
Nausea and vomiting	2%	3%
Diarrhea	2%	2%
Non-site specific		
Temperature regulation disturbances (fever and/or chills)	9%	10%
Malaise & fatigue	8%	8%

190 * In prophylaxis studies, symptoms associated with influenza-like illness were captured as
191 adverse events; subjects were enrolled during a winter respiratory season during which time
192 any symptoms that occurred were captured as adverse events.

193

194 **6.2 Postmarketing Experience**

195 In addition to adverse events reported from clinical trials, the following events have been
196 identified during postmarketing use of zanamivir (RELENZA). Because they are reported
197 voluntarily from a population of unknown size, estimates of frequency cannot be made. These
198 events have been chosen for inclusion due to a combination of their seriousness, frequency of
199 reporting, or potential causal connection to zanamivir (RELENZA).

200 Allergic Reactions: Allergic or allergic-like reaction, including oropharyngeal edema
201 [see Warnings and Precautions (5.2)].

202 **Psychiatric:** Delirium, including symptoms such as altered level of consciousness,
203 confusion, abnormal behavior, delusions, hallucinations, agitation, anxiety, nightmares [*see*
204 *Warnings and Precautions (5.3)*].

205 **Cardiac:** Arrhythmias, syncope.

206 **Neurologic:** Seizures.

207 **Respiratory:** Bronchospasm, dyspnea [*see Warnings and Precautions (5.1)*].

208 **Skin:** Facial edema; rash, including serious cutaneous reactions; urticaria [*see Warnings*
209 *and Precautions (5.2)*].

210 **7 DRUG INTERACTIONS**

211 Zanamivir is not a substrate nor does it affect cytochrome P450 (CYP) isoenzymes
212 (CYP1A1/2, 2A6, 2C9, 2C18, 2D6, 2E1, and 3A4) in human liver microsomes. No clinically
213 significant pharmacokinetic drug interactions are predicted based on data from in vitro studies.

214 The concurrent use of RELENZA with live attenuated influenza vaccine (LAIV)
215 intranasal has not been evaluated. However, because of potential interference between these
216 products, LAIV should not be administered within 2 weeks before or 48 hours after
217 administration of RELENZA, unless medically indicated. The concern about possible
218 interference arises from the potential for antiviral drugs to inhibit replication of live vaccine
219 virus.

220 Trivalent inactivated influenza vaccine can be administered at any time relative to use of
221 RELENZA [*see Clinical Pharmacology (12.4)*].

222 **8 USE IN SPECIFIC POPULATIONS**

223 **8.1 Pregnancy**

224 Pregnancy Category C. There are no adequate and well-controlled studies of zanamivir in
225 pregnant women. Zanamivir should be used during pregnancy only if the potential benefit
226 justifies the potential risk to the fetus.

227 Embryo/fetal development studies were conducted in rats (dosed from days 6 to 15 of
228 pregnancy) and rabbits (dosed from days 7 to 19 of pregnancy) using the same IV doses (1, 9,
229 and 90 mg/kg/day). Pre- and post-natal developmental studies were performed in rats (dosed
230 from day 16 of pregnancy until litter day 21 to 23). No malformations, maternal toxicity, or
231 embryotoxicity were observed in pregnant rats or rabbits and their fetuses. Because of
232 insufficient blood sampling timepoints in rat and rabbit reproductive toxicity studies, AUC
233 values were not available. In a subchronic study in rats at the 90 mg/kg/day IV dose, the AUC
234 values were greater than 300 times the human exposure at the proposed clinical dose.

235 An additional embryo/fetal study, in a different strain of rat, was conducted using
236 subcutaneous administration of zanamivir, 3 times daily, at doses of 1, 9, or 80 mg/kg during
237 days 7 to 17 of pregnancy. There was an increase in the incidence rates of a variety of minor
238 skeleton alterations and variants in the exposed offspring in this study. Based on AUC
239 measurements, the 80 mg/kg dose produced an exposure greater than 1,000 times the human
240 exposure at the proposed clinical dose. However, in most instances, the individual incidence rate

241 of each skeletal alteration or variant remained within the background rates of the historical
242 occurrence in the strain studied.

243 Zanamivir has been shown to cross the placenta in rats and rabbits. In these animals, fetal
244 blood concentrations of zanamivir were significantly lower than zanamivir concentrations in the
245 maternal blood.

246 **8.3 Nursing Mothers**

247 Studies in rats have demonstrated that zanamivir is excreted in milk. However, nursing
248 mothers should be instructed that it is not known whether zanamivir is excreted in human milk.
249 Because many drugs are excreted in human milk, caution should be exercised when RELENZA
250 is administered to a nursing mother.

251 **8.4 Pediatric Use**

252 Treatment of Influenza: Safety and effectiveness of RELENZA for treatment of
253 influenza have not been assessed in pediatric patients less than 7 years of age, but were studied in
254 a Phase III treatment study in pediatric patients, where 471 children 5 to 12 years of age received
255 zanamivir or placebo [see *Clinical Studies 14.1*]. Adolescents were included in the three
256 principal Phase III adult treatment studies. In these studies, 67 patients were 12 to 16 years of
257 age. No definite differences in safety and efficacy were observed between these adolescent
258 patients and young adults.

259 In a Phase I study of 16 children ages 6 to 12 years with signs and symptoms of
260 respiratory disease, 4 did not produce a measurable peak inspiratory flow rate (PIFR) through the
261 DISKHALER (3 with no adequate inhalation on request, 1 with missing data), 9 had measurable
262 PIFR on each of 2 inhalations, and 3 achieved measurable PIFR on only 1 of 2 inhalations.
263 Neither of two 6-year-olds and one of two 7-year-olds produced measurable PIFR. Overall, 8 of
264 the 16 children (including all those under 8 years old) either did not produce measurable
265 inspiratory flow through the DISKHALER or produced peak inspiratory flow rates below the
266 60 L/min considered optimal for the device under standardized in vitro testing; lack of
267 measurable flow rate was related to low or undetectable serum concentrations [see *Clinical*
268 *Pharmacology (12.3)*, *Clinical Studies (14.1)*]. Prescribers should carefully evaluate the ability
269 of young children to use the delivery system if prescription of RELENZA is considered.

270 Prophylaxis of Influenza: The safety and effectiveness of RELENZA for prophylaxis of
271 influenza have been studied in 4 Phase III studies where 273 children 5 to 11 years of age and
272 239 adolescents 12 to 16 years of age received RELENZA. No differences in safety and
273 effectiveness were observed between pediatric and adult subjects [see *Clinical Studies (14.2)*].

274 **8.5 Geriatric Use**

275 Of the total number of patients in 6 clinical studies of RELENZA for treatment of
276 influenza, 59 patients were 65 years of age and older, while 24 patients were 75 years of age and
277 older. Of the total number of patients in 4 clinical studies of RELENZA for prophylaxis of
278 influenza in households and community settings, 954 patients were 65 years of age and older,
279 while 347 patients were 75 years of age and older. No overall differences in safety or
280 effectiveness were observed between these patients and younger patients, and other reported

281 clinical experience has not identified differences in responses between the elderly and younger
282 patients, but greater sensitivity of some older individuals cannot be ruled out. Elderly patients
283 may need assistance with use of the device.

284 In 2 additional studies of RELENZA for prophylaxis of influenza in the nursing home
285 setting, efficacy was not demonstrated [see *Indications and Usage (1.3)*].


286 **10 OVERDOSAGE**

287 There have been no reports of overdosage from administration of RELENZA.

288 **11 DESCRIPTION**

289 The active component of RELENZA is zanamivir. The chemical name of zanamivir is 5-
290 (acetylamino)-4-[(aminoiminomethyl)-amino]-2,6-anhydro-3,4,5-trideoxy-D-glycero-D-galacto-
291 non-2-enonic acid. It has a molecular formula of $C_{12}H_{20}N_4O_7$ and a molecular weight of 332.3. It
292 has the following structural formula:

293



294

295 Zanamivir is a white to off-white powder for oral inhalation with a solubility of
296 approximately 18 mg/mL in water at 20°C.

297 RELENZA is for administration to the respiratory tract by oral inhalation only. Each
298 RELENZA ROTADISK contains 4 regularly spaced double-foil blisters with each blister
299 containing a powder mixture of 5 mg of zanamivir and 20 mg of lactose (which contains milk
300 proteins). The contents of each blister are inhaled using a specially designed breath-activated
301 plastic device for inhaling powder called the DISKHALER. After a RELENZA ROTADISK is
302 loaded into the DISKHALER, a blister that contains medication is pierced and the zanamivir is
303 dispersed into the air stream created when the patient inhales through the mouthpiece. The
304 amount of drug delivered to the respiratory tract will depend on patient factors such as
305 inspiratory flow. Under standardized in vitro testing, RELENZA ROTADISK delivers 4 mg of
306 zanamivir from the DISKHALER device when tested at a pressure drop of 3 kPa (corresponding
307 to a flow rate of about 62 to 65 L/min) for 3 seconds.

308 **12 CLINICAL PHARMACOLOGY**

309 **12.1 Mechanism of Action**

310 Zanamivir is an antiviral drug [see *Clinical Pharmacology (12.4)*].

311 **12.3 Pharmacokinetics**

312 Absorption and Bioavailability: Pharmacokinetic studies of orally inhaled zanamivir
313 indicate that approximately 4% to 17% of the inhaled dose is systemically absorbed. The peak

314 serum concentrations ranged from 17 to 142 ng/mL within 1 to 2 hours following a 10 mg dose.
315 The area under the serum concentration versus time curve (AUC_{∞}) ranged from 111 to
316 1,364 ng•hr/mL.

317 Distribution: Zanamivir has limited plasma protein binding (<10%).

318 Metabolism: Zanamivir is renally excreted as unchanged drug. No metabolites have
319 been detected in humans.

320 Elimination: The serum half-life of zanamivir following administration by oral inhalation
321 ranges from 2.5 to 5.1 hours. It is excreted unchanged in the urine with excretion of a single dose
322 completed within 24 hours. Total clearance ranges from 2.5 to 10.9 L/hr. Unabsorbed drug is
323 excreted in the feces.

324 Impaired Hepatic Function: The pharmacokinetics of zanamivir have not been studied
325 in patients with impaired hepatic function.

326 Impaired Renal Function: After a single intravenous dose of 4 mg or 2 mg of zanamivir
327 in volunteers with mild/moderate or severe renal impairment, respectively, significant decreases
328 in renal clearance (and hence total clearance: normals 5.3 L/hr, mild/moderate 2.7 L/hr, and
329 severe 0.8 L/hr; median values) and significant increases in half-life (normals 3.1 hr,
330 mild/moderate 4.7 hr, and severe 18.5 hr; median values) and systemic exposure were observed.
331 Safety and efficacy have not been documented in the presence of severe renal insufficiency. Due
332 to the low systemic bioavailability of zanamivir following oral inhalation, no dosage adjustments
333 are necessary in patients with renal impairment. However, the potential for drug accumulation
334 should be considered.

335 Pediatric Patients: The pharmacokinetics of zanamivir were evaluated in pediatric
336 patients with signs and symptoms of respiratory illness. Sixteen patients, 6 to 12 years of age,
337 received a single dose of 10 mg zanamivir dry powder via DISKHALER. Five patients had either
338 undetectable zanamivir serum concentrations or had low drug concentrations (8.32 to
339 10.38 ng/mL) that were not detectable after 1.5 hours. Eleven patients had C_{max} median values of
340 43 ng/mL (range 15 to 74) and AUC_{∞} median values of 167 ng•hr/mL (range 58 to 279). Low or
341 undetectable serum concentrations were related to lack of measurable PIFR in individual patients
342 [see *Use in Specific Populations (8.4), Clinical Studies (14.1)*].

343 Geriatric Patients: The pharmacokinetics of zanamivir have not been studied in patients
344 over 65 years of age [see *Use in Specific Populations (8.5)*].

345 Gender, Race, and Weight: In a population pharmacokinetic analysis in patient
346 studies, no clinically significant differences in serum concentrations and/or pharmacokinetic
347 parameters (V/F , CL/F , k_a , AUC_{0-3} , C_{max} , T_{max} , CL_r , and % excreted in urine) were observed
348 when demographic variables (gender, age, race, and weight) and indices of infection (laboratory
349 evidence of infection, overall symptoms, symptoms of upper respiratory illness, and viral titers)
350 were considered. There were no significant correlations between measures of systemic exposure
351 and safety parameters.

352 **12.4 Microbiology**

353 Mechanism of Action: Zanamivir is an inhibitor of influenza virus neuraminidase
354 affecting release of viral particles.

355 Antiviral Activity: The antiviral activity of zanamivir against laboratory and clinical
356 isolates of influenza virus was determined in cell culture assays. The concentrations of zanamivir
357 required for inhibition of influenza virus were highly variable depending on the assay method
358 used and virus isolate tested. The 50% and 90% effective concentrations (EC₅₀ and EC₉₀) of
359 zanamivir were in the range of 0.005 to 16.0 μM and 0.05 to >100 μM, respectively
360 (1 μM = 0.33 mcg/mL). The relationship between the cell culture inhibition of influenza virus by
361 zanamivir and the inhibition of influenza virus replication in humans has not been established.

362 Resistance: Influenza viruses with reduced susceptibility to zanamivir have been
363 selected in cell culture by multiple passages of the virus in the presence of increasing
364 concentrations of the drug. Genetic analysis of these viruses showed that the reduced
365 susceptibility in cell culture to zanamivir is associated with mutations that result in amino acid
366 changes in the viral neuraminidase or viral hemagglutinin or both. Resistance mutations selected
367 in cell culture which result in neuraminidase amino acid substitutions include E119G/A/D and
368 R292K. Mutations selected in cell culture in hemagglutinin include: K68R, G75E, E114K,
369 N145S, S165N, S186F, N199S, and K222T.

370 In an immunocompromised patient infected with influenza B virus, a variant virus
371 emerged after treatment with an investigational nebulized solution of zanamivir for 2 weeks.
372 Analysis of this variant showed a hemagglutinin substitution (T198I) which resulted in a reduced
373 affinity for human cell receptors, and a substitution in the neuraminidase active site (R152K)
374 which reduced the enzyme's activity to zanamivir by 1,000-fold. Insufficient information is
375 available to characterize the risk of emergence of zanamivir resistance in clinical use.

376 Cross-Resistance: Cross-resistance has been observed between some
377 zanamivir-resistant and some oseltamivir-resistant influenza virus mutants generated in cell
378 culture. However, some of the in cell culture zanamivir-induced resistance mutations,
379 E119G/A/D and R292K, occurred at the same neuraminidase amino acid positions as in the
380 clinical isolates resistant to oseltamivir, E119V and R292K. No studies have been performed to
381 assess risk of emergence of cross-resistance during clinical use.

382 Influenza Vaccine Interaction Study: An interaction study (n = 138) was conducted to
383 evaluate the effects of zanamivir (10 mg once daily) on the serological response to a single dose
384 of trivalent inactivated influenza vaccine, as measured by hemagglutination inhibition titers.
385 There was no difference in hemagglutination inhibition antibody titers at 2 weeks and 4 weeks
386 after vaccine administration between zanamivir and placebo recipients.

387 Influenza Challenge Studies: Antiviral activity of zanamivir was supported for
388 infection with influenza A virus, and to a more limited extent for infection with influenza B
389 virus, by Phase I studies in volunteers who received intranasal inoculations of challenge strains
390 of influenza virus, and received an intranasal formulation of zanamivir or placebo starting before
391 or shortly after viral inoculation.

392 **13 NONCLINICAL TOXICOLOGY**

393 **13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility**

394 Carcinogenesis: In 2-year carcinogenicity studies conducted in rats and mice using a
395 powder formulation administered through inhalation, zanamivir induced no statistically
396 significant increases in tumors over controls. The maximum daily exposures in rats and mice
397 were approximately 23 to 25 and 20 to 22 times, respectively, greater than those in humans at the
398 proposed clinical dose based on AUC comparisons.

399 Mutagenesis: Zanamivir was not mutagenic in in vitro and in vivo genotoxicity assays
400 which included bacterial mutation assays in *S. typhimurium* and *E. coli*, mammalian mutation
401 assays in mouse lymphoma, chromosomal aberration assays in human peripheral blood
402 lymphocytes, and the in vivo mouse bone marrow micronucleus assay.

403 Impairment of Fertility: The effects of zanamivir on fertility and general reproductive
404 performance were investigated in male (dosed for 10 weeks prior to mating, and throughout
405 mating, gestation/lactation, and shortly after weaning) and female rats (dosed for 3 weeks prior
406 to mating through Day 19 of pregnancy, or Day 21 post partum) at IV doses 1, 9, and
407 90 mg/kg/day. Zanamivir did not impair mating or fertility of male or female rats, and did not
408 affect the sperm of treated male rats. The reproductive performance of the F1 generation born to
409 female rats given zanamivir was not affected. Based on a subchronic study in rats at a
410 90 mg/kg/day IV dose, AUC values ranged between 142 and 199 mcg•hr/mL (>300 times the
411 human exposure at the proposed clinical dose).

412 **14 CLINICAL STUDIES**

413 **14.1 Treatment of Influenza**

414 Adults and Adolescents: The efficacy of RELENZA 10 mg inhaled twice daily for
415 5 days in the treatment of influenza has been evaluated in placebo-controlled studies conducted
416 in North America, the Southern Hemisphere, and Europe during their respective influenza
417 seasons. The magnitude of treatment effect varied between studies, with possible relationships to
418 population-related factors including amount of symptomatic relief medication used.

419 Populations Studied: The principal Phase III studies enrolled 1,588 patients ages
420 12 years and older (median age 34 years, 49% male, 91% Caucasian), with uncomplicated
421 influenza-like illness within 2 days of symptom onset. Influenza was confirmed by culture,
422 hemagglutination inhibition antibodies, or investigational direct tests. Of 1,164 patients with
423 confirmed influenza, 89% had influenza A and 11% had influenza B. These studies served as the
424 principal basis for efficacy evaluation, with more limited Phase II studies providing supporting
425 information where necessary. Following randomization to either zanamivir or placebo (inhaled
426 lactose vehicle), all patients received instruction and supervision by a healthcare professional for
427 the initial dose.

428 Principal Results: The definition of time to improvement in major symptoms of
429 influenza included no fever and self-assessment of “none” or “mild” for headache, myalgia,
430 cough, and sore throat. A Phase II and a Phase III study conducted in North America (total of

431 over 600 influenza-positive patients) suggested up to 1 day of shortening of median time to this
432 defined improvement in symptoms in patients receiving zanamivir compared with placebo,
433 although statistical significance was not reached in either of these studies. In a study conducted
434 in the Southern Hemisphere (321 influenza-positive patients), a 1.5-day difference in median
435 time to symptom improvement was observed. Additional evidence of efficacy was provided by
436 the European study.

437 *Other Findings:* There was no consistent difference in treatment effect in patients
438 with influenza A compared with influenza B; however, these trials enrolled smaller numbers of
439 patients with influenza B and thus provided less evidence in support of efficacy in influenza B.

440 In general, patients with lower temperature (e.g., 38.2°C or less) or investigator-rated as
441 having less severe symptoms at entry derived less benefit from therapy.

442 No consistent treatment effect was demonstrated in patients with underlying chronic
443 medical conditions, including respiratory or cardiovascular disease [*see Warnings and*
444 *Precautions (5.3)*].

445 No consistent differences in rate of development of complications were observed
446 between treatment groups.

447 Some fluctuation of symptoms was observed after the primary study endpoint in both
448 treatment groups.

449 **Pediatric Patients:** The efficacy of RELENZA 10 mg inhaled twice daily for 5 days in
450 the treatment of influenza in pediatric patients has been evaluated in a placebo-controlled study
451 conducted in North America and Europe, enrolling 471 patients, ages 5 to 12 years (55% male,
452 90% Caucasian), within 36 hours of symptom onset. Of 346 patients with confirmed influenza,
453 65% had influenza A and 35% had influenza B. The definition of time to improvement included
454 no fever and parental assessment of no or mild cough and absent/minimal muscle and joint aches
455 or pains, sore throat, chills/feverishness, and headache. Median time to symptom improvement
456 was 1 day shorter in patients receiving zanamivir compared with placebo. No consistent
457 differences in rate of development of complications were observed between treatment groups.
458 Some fluctuation of symptoms was observed after the primary study endpoint in both treatment
459 groups.

460 Although this study was designed to enroll children ages 5 to 12 years, the product is
461 indicated only for children 7 years of age and older. This evaluation is based on the combination
462 of lower estimates of treatment effect in 5- and 6-year-olds compared with the overall study
463 population, and evidence of inadequate inhalation through the DISKHALER in a
464 pharmacokinetic study [*see Use in Specific Populations (8.4), Clinical Pharmacology (12.3)*].

465 **14.2 Prophylaxis of Influenza**

466 The efficacy of RELENZA in preventing naturally occurring influenza illness has been
467 demonstrated in 2 post-exposure prophylaxis studies in households and 2 seasonal prophylaxis
468 studies during community outbreaks of influenza. The primary efficacy endpoint in these studies
469 was the incidence of symptomatic, laboratory-confirmed influenza, defined as the presence of 2
470 or more of the following symptoms: oral temperature $\geq 100^{\circ}\text{F}/37.8^{\circ}\text{C}$ or feverishness, cough,

471 headache, sore throat, and myalgia; and laboratory confirmation of influenza A or B by culture,
472 PCR, or seroconversion (defined as a 4-fold increase in convalescent antibody titer from
473 baseline).

474 **Household Prophylaxis Studies:** Two studies assessed post-exposure prophylaxis in
475 household contacts of an index case. Within 1.5 days of onset of symptoms in an index case,
476 each household (including all family members ≥ 5 years of age) was randomized to RELENZA
477 10 mg inhaled once daily or placebo inhaled once daily for 10 days. In the first study only, each
478 index case was randomized to RELENZA 10 mg inhaled twice daily for 5 days or inhaled
479 placebo twice daily for 5 days. In this study, the proportion of households with at least 1 new
480 case of symptomatic laboratory-confirmed influenza was reduced from 19.0% (32 of
481 168 households) for the placebo group to 4.1% (7 of 169 households) for the group receiving
482 RELENZA.

483 In the second study, index cases were not treated. The incidence of symptomatic
484 laboratory-confirmed influenza was reduced from 19.0% (46 of 242 households) for the placebo
485 group to 4.1% (10 of 245 households) for the group receiving RELENZA.

486 **Seasonal Prophylaxis Studies:** Two seasonal prophylaxis studies assessed RELENZA
487 10 mg inhaled once daily versus placebo inhaled once daily for 28 days during community
488 outbreaks. The first study enrolled subjects 18 years of age or greater (mean age 29 years) from 2
489 university communities. The majority of subjects were unvaccinated (86%). In this study, the
490 incidence of symptomatic laboratory-confirmed influenza was reduced from 6.1% (34 of 554)
491 for the placebo group to 2.0% (11 of 553) for the group receiving RELENZA.

492 The second seasonal prophylaxis study enrolled subjects 12 to 94 years of age (mean age
493 60 years) with 56% of them older than 65 years of age. Sixty-seven percent of the subjects were
494 vaccinated. In this study, the incidence of symptomatic laboratory-confirmed influenza was
495 reduced from 1.4% (23 of 1,685) for the placebo group to 0.2% (4 of 1,678) for the group
496 receiving RELENZA.

497 **16 HOW SUPPLIED/STORAGE AND HANDLING**

498 RELENZA is supplied in a circular double-foil pack (a ROTADISK) containing 4 blisters
499 of the drug. Five ROTADISKS are packaged in a white polypropylene tube. The tube is
500 packaged in a carton with 1 blue and gray DISKHALER inhalation device (NDC 0173-0681-01).

501 **Store at 25°C (77°F); excursions permitted to 15° to 30°C (59° to 86°F) (see USP**
502 **Controlled Room Temperature).** Keep out of reach of children. Do not puncture any
503 RELENZA ROTADISK blister until taking a dose using the DISKHALER.

504 **17 PATIENT COUNSELING INFORMATION**

505 *See FDA-Approved Patient Labeling (17.6).*

506 **17.1 Bronchospasm**

507 **Patients should be advised of the risk of bronchospasm, especially in the setting of**
508 **underlying airways disease, and should stop RELENZA and contact their physician if they**
509 **experience increased respiratory symptoms during treatment such as worsening wheezing,**

510 **shortness of breath, or other signs or symptoms of bronchospasm [see Warnings and**
511 **Precautions (5.1)]. If a decision is made to prescribe RELENZA for a patient with asthma**
512 **or chronic obstructive pulmonary disease, the patient should be made aware of the risks**
513 **and should have a fast-acting bronchodilator available.**

514 **17.2 Concomitant Bronchodilator Use**

515 Patients scheduled to take inhaled bronchodilators at the same time as RELENZA should
516 be advised to use their bronchodilators before taking RELENZA.

517 **17.3 Neuropsychiatric Events**

518 Patients with influenza (the flu), particularly children and adolescents, may be at an
519 increased risk of seizures, confusion, or abnormal behavior early in their illness. These events
520 may occur after beginning RELENZA or may occur when flu is not treated. These events are
521 uncommon but may result in accidental injury to the patient. Therefore, patients should be
522 observed for signs of unusual behavior and a healthcare professional should be contacted
523 immediately if the patient shows any signs of unusual behavior [see Warnings and Precautions
524 (5.3)].

525 **17.4 Instructions for Use**

526 Patients should be instructed in use of the delivery system. Instructions should include a
527 demonstration whenever possible. For the proper use of RELENZA, the patient should read and
528 follow carefully the accompanying Patient Instructions for Use.

529 **If RELENZA is prescribed for children, it should be used only under adult**
530 **supervision and instruction, and the supervising adult should first be instructed by a**
531 **healthcare professional [see Dosage and Administration (2.1)].**

532 **17.5 Risk of Influenza Transmission to Others**

533 Patients should be advised that the use of RELENZA for treatment of influenza has not
534 been shown to reduce the risk of transmission of influenza to others.

535 **17.6 FDA-Approved Patient Labeling and Instructions for Use**

536 See separate leaflet.

537
538 RELENZA, DISKHALER, and ROTADISK are registered trademarks of GlaxoSmithKline.

539
540



541
542 GlaxoSmithKline
543 Research Triangle Park, NC 27709

544
545 ©2008, GlaxoSmithKline. All rights reserved.

546

Patient Labeling

RELENZA[®] (zanamivir) Inhalation Powder

This leaflet contains important patient information about RELENZA (zanamivir) Inhalation Powder, and should be read completely before beginning treatment. It does not, however, take the place of discussions with your healthcare provider about your medical condition or your treatment. This summary does not list all benefits and risks of RELENZA. The medication described here can only be prescribed and dispensed by a licensed healthcare provider, who has information about your medical condition and more information about the drug, including how to take it, what to expect, and potential side effects. If you have any questions about RELENZA, talk with your healthcare provider.

What is RELENZA?

RELENZA (ruh-LENS-uh) is a medicine for the treatment of influenza (flu, infection caused by influenza virus) and for reducing the chance of getting the flu in community and household settings. It belongs to a group of medicines called neuraminidase inhibitors. These medications attack the influenza virus and prevent it from spreading inside your body. RELENZA treats the cause of influenza at its source, rather than simply masking the symptoms.

Important Safety Information About RELENZA

Some patients have had bronchospasm (wheezing) or serious breathing problems when they used RELENZA. Many but not all of these patients had previous asthma or chronic obstructive pulmonary disease. RELENZA has not been shown to shorten the duration of influenza in people with these diseases. Because of the risk of side effects and because it has not been shown to help them, RELENZA is not recommended for people with chronic respiratory disease such as asthma or chronic obstructive pulmonary disease.

If you develop worsening respiratory symptoms such as wheezing or shortness of breath, stop using RELENZA and contact your healthcare provider right away.

If you have chronic respiratory disease such as asthma and chronic obstructive pulmonary disease and your healthcare provider has prescribed RELENZA, you should have a fast-acting, inhaled bronchodilator available for your use. If you are scheduled to use an inhaled bronchodilator at the same time as RELENZA, use the inhaled bronchodilator **before** using RELENZA.

Read the rest of this leaflet for more information about side effects and risks.

Other kinds of infections can appear like influenza or occur along with influenza, and need different kinds of treatment. Contact your healthcare provider if you feel worse or develop new symptoms during or after treatment, or if your influenza symptoms do not start to get better.

Who should not take RELENZA?

41 RELENZA is not recommended for people who have chronic lung disease such as
42 asthma or chronic obstructive pulmonary disease. RELENZA has not been shown to shorten the
43 duration of influenza in people with these diseases, and some people have had serious side
44 effects of bronchospasm and worsening lung function. (See the section of this Patient
45 Information entitled “**Important Safety Information About RELENZA.**”)

46 You should not take RELENZA if you are allergic to zanamivir or any other ingredient of
47 RELENZA. Also tell your healthcare provider if you have any type of chronic condition
48 including lung or heart disease, if you are allergic to any other medicines or food products, or if
49 you are pregnant.

50 RELENZA was not effective in reducing the chance of getting the flu in 2 studies in
51 nursing home patients.

52 RELENZA does not treat flu-like illness that is not caused by influenza virus.

53

54 **Who should consider taking RELENZA?**

55 Adult and pediatric patients at least 7 years of age who have influenza symptoms that
56 appeared within the previous day or two. Typical symptoms of influenza include sudden onset of
57 fever, cough, headache, fatigue, muscular weakness, and sore throat.

58 RELENZA can also help reduce the chance of getting the flu in adults and children at
59 least 5 years of age who have a higher chance of getting the flu because they spend time with
60 someone who has the flu. RELENZA can also reduce the chance of getting the flu if there is a flu
61 outbreak in the community.

62 The use of RELENZA for the treatment of flu has not been shown to reduce the risk of
63 spreading the virus to others.

64

65 **Can I take other medications with RELENZA?**

66 RELENZA has been shown to have an acceptable safety profile when used as labeled,
67 with minimal risk of drug interactions. Your healthcare provider may recommend taking other
68 medications, including over-the-counter medications, to reduce fever or other symptoms while
69 you are taking RELENZA. Before starting treatment, make sure that your healthcare provider
70 knows if you are taking other medicines. If you are scheduled to use an inhaled bronchodilator at
71 the same time as RELENZA, you should use the inhaled bronchodilator **before** using
72 RELENZA.

73 Before taking RELENZA, please let your healthcare provider know if you received live
74 attenuated influenza vaccine (FLUMIST[®]) intranasal in the past 2 weeks.

75

76 **How and when should I take RELENZA?**

77 RELENZA is packaged in medicine disks called ROTADISKS[®] and is inhaled by mouth
78 using a delivery device called a DISKHALER[®]. Each ROTADISK contains 4 blisters. Each
79 blister contains 5 mg of active drug and 20 mg of lactose powder (which contains milk proteins).

80 You should receive a demonstration on how to use RELENZA in the DISKHALER from
81 a healthcare provider. Before taking RELENZA, read the “Patient Instructions for Use.” Make
82 sure that you understand these instructions and talk to your healthcare provider if you have any
83 questions. Children who use RELENZA should always be supervised by an adult who
84 understands how to use RELENZA. Proper use of the DISKHALER to inhale the drug is
85 necessary for safe and effective use of RELENZA.

86 If you have the flu the usual dose for treatment is 2 inhalations of RELENZA (1 blister
87 per inhalation) twice daily (in the morning and evening) for 5 days. It is important that you begin
88 your treatment with RELENZA as soon as possible from the first appearance of your flu
89 symptoms. Take 2 doses on the first day of treatment whenever possible if there are at least
90 2 hours between doses.

91 To reduce the chance of getting the flu, the usual dose is 2 inhalations of RELENZA
92 (1 blister per inhalation) once daily for 10 or 28 days as prescribed by your healthcare provider.

93 Never share RELENZA with anyone, even if they have the same symptoms. If you feel
94 worse or develop new symptoms during treatment with RELENZA, or if your flu symptoms do
95 not start to get better, stop using the medicine and contact your healthcare provider.

96 **What if I miss a dose?**

97
98 If you forget to take your medicine at any time, take the missed dose as soon as you
99 remember, except if it is near the next dose (within 2 hours). Then continue to take RELENZA at
100 the usual times. You do not need to take a double dose. If you have missed several doses, inform
101 your healthcare provider and follow the advice given to you.

102 **What are important or common possible side effects of taking RELENZA?**

103 Some patients have had breathing problems while taking RELENZA. This can be very
104 serious and need treatment right away. Most of the patients who had this problem had asthma or
105 chronic obstructive pulmonary disease, but some did not. If you have trouble breathing or have
106 wheezing after your dose of RELENZA, stop taking RELENZA and get medical attention.

107
108 In studies, the most common side effects with RELENZA have been headaches; diarrhea;
109 nausea; vomiting; nasal irritation; bronchitis; cough; sinusitis; ear, nose, and throat infections;
110 and dizziness. Other side effects that have been reported, but were not as common, include
111 rashes and allergic reactions, some of which were severe.

112 People with influenza (the flu), particularly children and adolescents, may be at an
113 increased risk of seizures, confusion, or abnormal behavior early in their illness. These events
114 may occur after beginning RELENZA or may occur when flu is not treated. These events are
115 uncommon but may result in accidental injury to the patient. Therefore, patients should be
116 observed for signs of unusual behavior and a healthcare professional should be contacted
117 immediately if the patient shows any signs of unusual behavior.

118 This list of side effects is not complete. Your healthcare provider or pharmacist can
119 discuss with you a more complete list of possible side effects with RELENZA. Talk to your
120 healthcare provider promptly about any side effects you have.

121 Please refer to the section entitled "**Important Safety Information About RELENZA**"
122 for additional information.

123

124 **Should I get a flu shot?**

125 RELENZA is not a substitute for a flu shot. You should receive an annual flu shot
126 according to guidelines on immunization practices that your healthcare provider can share with
127 you.

128

129 **What if I am pregnant or nursing?**

130 If you are pregnant or planning to become pregnant while taking RELENZA, talk to your
131 healthcare provider before taking this medication. RELENZA is normally not recommended for
132 use during pregnancy or nursing, as the effects on the unborn child or nursing infant are
133 unknown.

134

135 **How and where should I store RELENZA?**

136 RELENZA should be stored at room temperature below 77°F (25°C). RELENZA is not
137 in a childproof container. Keep RELENZA out of the reach of children. Discard the
138 DISKHALER after finishing your treatment.

139

140

PATIENT INSTRUCTIONS FOR USE



141

142

**IMPORTANT: Read Step-by-Step Instructions
before using the DISKHALER®.**

143

144

145

Be sure to take the dose your healthcare provider has prescribed.

146

147

BEFORE YOU START:

148

149

Please read the entire Patient Labeling for important information about the effects of RELENZA including the section “Important Safety Information About RELENZA” for information about the risk of breathing difficulties.

150

151

152

153

If RELENZA is prescribed for a child, dosing should be supervised by an adult who understands how to use RELENZA and has been instructed in its use by a healthcare provider.

154

155



156 **Step-by-step instructions for using the DISKHALER®**

157


158 **Step A: Load the medicine into the DISKHALER**

159

- 160 1. Start by pulling off the blue cover.

161

- 162 2. **Always check inside the mouthpiece to make sure it is clear before each use. If foreign**
163 **objects are in the mouthpiece, they could be inhaled and cause serious harm.**
164
- 165 3. Pull the white mouthpiece by the edges to extend the white tray all the way.
166
- 167 4. Once the white tray is extended all the way, find the raised ridges on each side of it. Press in
168 these ridges, both sides at the same time, and **pull the whole white tray out of the**
169 **DISKHALER body.**
170
- 171 5. Place one silver medicine disk onto the dark brown wheel, flat side up. The four silver
172 blisters on the underside of the medicine disk will drop neatly into the four holes in the
173 wheel.
174
- 175 6. Push in the white tray as far as it will go. Now the DISKHALER is loaded with medicine.
176



177

178 **Step B: Puncture the blister**

179

180 **Be sure to keep the DISKHALER level.**

181

182 **The DISKHALER punctures one blister of medicine at a time so you can inhale the right**
183 **amount. It does not matter which blister you start with. Check to make sure that the silver**
184 **foil is unbroken.**

185

186 1. Be sure to keep the DISKHALER level so the medicine does not spill out.

187

188 2. Locate the half-circle flap with the name “RELENZA” on top of the DISKHALER.

189

190 3. Lift this flap from the outer edge until it cannot go any farther. Flap must be **straight up** for
191 the plastic needle to puncture both the **top** and **bottom** of the silver medicine disk inside.

192

193 4. Keeping the DISKHALER level, click the flap down into place.



194

195

196 **Step C: Inhale**

197

198 1. Before putting the white mouthpiece into your mouth, breathe all the way out (exhale).

199

200 **Then put the white mouthpiece into your mouth. Be sure to keep the DISKHALER level so**
201 **the medicine does not spill out.**

202

203 2. Close your lips firmly around the mouthpiece. Be sure not to cover the small holes on either
204 side of it.

205

206 3. Breathe in through your mouth steadily and as deeply as you can. Your breath pulls the
207 medicine into your airways and lungs.

208


209 4. Hold your breath for a few seconds to help RELENZA stay in your lungs where it can work.

210

211 **To take another inhalation, move to the next blister by following Step D below.**

212

213 **Once you've inhaled the number of blisters prescribed by your healthcare provider,**
214 **replace the cover until your next dose.**




215
216
217
218
219
220
221
222
223
224
225
226
227
228

Step D: Move the medicine disk to the next blister

1. **Pull** the mouthpiece to extend the white tray, without removing it.
2. Then **push** it back until it clicks. This pull-push motion rotates the medicine disk to the next blister.
3. To take your next inhalation, repeat Steps B and C.

If all four blisters in the medicine disk have been used, you are ready to start a new medicine disk (see Step A). Check to make sure that the silver foil is unbroken each time you are ready to puncture the next blister.



229

IMPORTANT INSTRUCTIONS

Read this entire leaflet before using RELENZA. Even if you have had a previous prescription for RELENZA, read this leaflet to see if any information has changed.

If you have the flu, the usual dose is 2 inhalations twice daily. To reduce the chance of getting the flu, the usual dose is 2 inhalations once daily. However, you must take the

number of inhalations your healthcare provider has prescribed.

If you feel worse or develop new symptoms during or after treatment, or if your flu symptoms do not start to improve, stop using the medicine and contact your healthcare provider.

Keep out of reach of children.

Always check inside the mouthpiece to make sure it is clear before each use. If foreign objects are in the mouthpiece, they could be inhaled and cause serious harm.

Always replace the cover after each use.

Throw away the DISKHALER after treatment is completed.

This DISKHALER is for use only with RELENZA. Do not use the RELENZA DISKHALER device with FLOVENT[®] (fluticasone propionate) and do not use RELENZA with the FLOVENT DISKHALER device.

Store at 25°C (77°F); excursions permitted to 15° to 30°C (59° to 86°F) (see USP Controlled Room Temperature).

REMEMBER: This medicine has been prescribed for you by your healthcare provider. **DO NOT** give this medicine to anyone else.

230

231 RELENZA, FLOVENT, ROTADISK, and DISKHALER are registered trademarks of
232 GlaxoSmithKline.

233 FLUMIST is a registered trademark of MedImmune, Inc.

234

235

 **GlaxoSmithKline**

236

237 GlaxoSmithKline

238 Research Triangle Park, NC 27709

239

240 ©2008, GlaxoSmithKline. All rights reserved.

241

242 February 2008

243

RLZ:3PIL