

128 CODE

MA

RID[®] MOUSSE

Drug Facts

Active ingredients (calculated without propellant) Purpose
 Piperonyl butoxide (4%) _____ lice treatment
 Pyrethrum extract* (equivalent to 0.55% pyrethrins) _____ lice treatment

Uses treats head, pubic (crab), and body lice

Warnings

- For external use only
- Flammable: Keep away from fire or flame

Do not use near the eyes

Ask a doctor before use if you have an allergy to ragweed

When using this product

- keep out of eyes when rinsing hair
- close eyes tightly and do not open eyes until product is rinsed out
- protect eyes with washcloth, towel or other suitable method
- if product gets into the eyes, immediately flush with water
- do not use inside the nose, mouth, or vagina. Irritation may occur.
- do not inhale; use in a well ventilated area
- do not puncture or incinerate. Contents under pressure.

Stop use and ask a doctor if

- skin irritation or infection is present or develops
- infestation of eyebrows or eyelashes occurs

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions

- Important: Read warnings before using
- shake well before using
- holding the can upside down, apply RID[®] Mousse to dry hair or other affected area. Massage until all the hair is thoroughly wet with product. For best results against head lice, see complete Directions in the Consumer Information insert.
- allow product to remain on the hair for 10 minutes but no longer
- wash area thoroughly with warm water and soap or shampoo
- a fine-toothed comb or a special lice/it removing comb may be used to help remove dead lice or their eggs (nits) from hair
- a second treatment must be done in 7 to 10 days to kill any newly hatched lice


Other Information

- store at 20°-25°C (68°-77°F)
- do not store at temperature above 45°C (110°F)
- keep in a cool place out of the sun

Inactive Ingredients cetylalcohol, isobutane, PEG-20 etherate, propane, propylene glycol, purified water, quaternium-52, SD Alcohol 3-C (26.5% w/w)

Questions? call 1-800-RID LICE
 (1-800-743-5423)

MAXIMUM
STRENGTH




Distributed By
 CONSUMER HEALTH CARE GROUP,
 PFIZER INC, NEW YORK, NY 10017

U.S. Pat. 5,783,202 SM 40120005P
 *50% extract (not USP)

EXP

LOT



Instruc

M
S



KEEP THIS AND A
See box

ENLARGED FOR PROOFREADING

11111

MAXIMUM STRENGTH



MOUSSE

Instrucciones En Español Incluidas

MAXIMUM STRENGTH

MAXIMUM STRENGTH


MAXIMUM STRENGTH



MOUSSE

MOUSSE

MOUSSE



Manufactured By
JANUARY HEALTH CARE GROUP,
INC., NEW YORK, NY 10017

L 5,783,202 SM 40120005P
Extract (not USP)

NET WT.
5.5 OZ (156 g)

KEEP THIS AND ALL DRUGS OUT OF THE REACH OF CHILDREN.
See back panel for additional warnings.

Plus Patented
Egg Removal
Comb




Visit us at <http://www.licerid.com>

ENLARGED FOR PROOFREADING

MAXIMUM STRENGTH



MOISTURE



LED THE WAY ALL OTHERS ON THE MARKET CHILDREN, PREGNANT AND LACTATING WOMEN
DISCONTINUED BY CONSUMER HEALTHCARE GROUP, PETERLIFE, INC., NEW YORK, NY 10017

Drug Facts

Active Ingredients
Featuralized without propylparan)
Piperonyl butoxide (4%), Pyrethrum extract
(equivalent to 0.33% pyrethrins)

Purpose lice treatment

Uses treats head, pubic (crab), and body lice

Warnings

- For external use only
- Flammable; Keep away from fire or flame
- Do not use near the eyes
- Ask a doctor before use if you have an allergy to pyrethroid

When using this product

- keep out of eyes when rinsing hair
- close eyes tightly and do not open eyes until product is rinsed out
- protect eyes with washcloth, towel or other suitable method
- if product gets into the eyes, immediately flush with water

- do not use inside the nose, mouth, or vagina. Irritation may occur.
- do not inhale; use in a well ventilated area
- do not puncture or incinerate. Contents under pressure.

Stop use and ask a doctor if

- skin irritation or infection is present or develops
- inhalation of eye drops or eye ointment occurs

Keep out of reach of children.
If swallowed, get medical help or contact a Poison Control Center right away.

NET WT. 5.5 OZ (156 g)

Drug Facts (continued)

Directions

- Important: Read warnings before using
- shake well before using
- holding the can upside down, apply RIDE Moisture to dry hair or other affected area. Massage until all the hair is thoroughly wet with product. For best results against head lice, see complete Directions in the Consumer Information Insert.
- allow product to remain on the hair for 10 minutes, but no longer
- wash area thoroughly with warm water and soap or shampoo
- a fine-toothed comb or a special lice/all removing comb may be used to help remove dead lice or their eggs (nits) from hair
- a second treatment must be done in 7 to 10 days to kill any newly hatched lice

Other Information

- store at 20°-25°C (68°-77° F)
- do not store at temperature above 43°C (110° F)
- keep this cool place out of the sun

Inactive Ingredients cetylalcohol, isobutane, PEG-20 stearate, propylene glycol, purified water, quaternium-15, SD Alcohol-3-C (26.5% w/w)

Questions? call 1-800-RID LICE
(1-800-743-6423)
Visit us at <http://www.lice.rid.com>
50% extra (not USP)
U.S. Pat. 5,783,202

EXP
LOT



SN 26120001P

ENLARGED FOR PROOFREADING

attach
desk C

