This label may not be the latest approved by FDA.
For current labeling information, please visit https://www.fda.gov/drugsatfda

Drug Facts (continued)

Stop use and ask a doctor if
- you experience any of the following signs of stomach bleeding:
  - feel faint
  - vomit blood
  - have bloody or black stools
  - have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
  - chest pain
  - trouble breathing
  - weakness in one part or side of body
  - slurred speech
  - leg swel ing
- redness or swelling is present in the painful area
- any new symptoms appear
- fever gets worse or lasts more than 3 days
- you have difficulty swallowing or the caplet feels stuck in your throat
- you get nervous, dizzy, or sleepless
- nasal congestion lasts more han 7 days

If pregnant or breast-feeding, ask a hea th professional before use. It is especia ly important not to use naproxen sodium during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Drug Facts (continued)

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions
- **do not take more than directed**
- **the smallest effective dose should be used**
- **swa low whole;** do not crush or chew
- **drink a full glass of water with each dose**
- adu ts and ch ldren 12 years and older: **1 caplet every 12 hours;** do not take more than 2 caplets in 24 hours
- ch ldren under 12 years: do not use

Other information
- **each caplet contains:** sodium 22 mg
- store at 20-25°C (68-77°F)
- store in a dry place

Inactive ingredients colloidal s licon dioxide, hypromellose, lactose, magnesium stearate, microcrystal ine cellulose, polyethylene glycol, povidone, talc, titanium dioxide

Questions or comments? 1-800-986-0369
(Mon-Fri 9AM – 5PM EST)

Bayer

Do not use if the blister seal is cut or broken.

Dist. by:
Bayer HealthCare LLC
Whippany, NJ 07981

Bayer, the Bayer Cross, Aleve-D and All Day Strong are registered trademarks of Bayer.

XXXXX

3 25866 56952 3

1 CAPLET - 12 HOURS

10 CAPLETS
CAPSULE-SHAPED TABLETS

ALEVE-D
SINUS & COLD
ALL DAY STRONG®

BAYER

12 HOUR MULTI-SYMPTOM RELIEF OF
SINUS PRESSURE
HEADACHE • SINUS CONGESTION
NASAL CONGESTION • BODY ACHES

naproxen sodium 220 mg (NSAID)*
Pseudoephedrine HCl 120 mg extended-release tablet
Pain reliever-fever reducer/nasal decongestant

NON-DROWSY

ALL DAY STRONG®
ALEVE-D®
SINUS & COLD

Drug Facts

Active ingredients (in each caplet)	Purposes
Naproxen sodium 220 mg (naproxen 200 mg) (NSAID)*	Pain reliever/fever reducer
Pseudoephedrine HCl 120 mg, extended-release	Nasal decongestant

*nonsteroidal anti-inflammatory drug

Uses
temporar ly relieves these cold, sinus, and flu symptoms:
- sinus pressure
- minor body aches and pains
- headache
- nasal and sinus congestion (promotes sinus drainage and restores freer brea hing hrough the nose)
- fever

Warnings
Allergy alert: Naproxen sodium may cause a severe allergic reaction, especia ly in people a lergic to aspirin. Symptoms may include:
- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:
- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood hinning (an icoagulant) or steroid drug
- take other drugs containing prescription or nonprescrip ion NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Drug Facts (continued)

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher f you use more than directed or for longer than directed.

Do not use
- if you have ever had an allergic reac ion to any other pain reliever/fever reducer
- right before or after heart surgery
- if you are now taking a prescription monoamine oxidase inh bitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- in children under 12 years of age

Ask a doctor before use f
- the stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, thyroid disease, diabetes, have trouble urinating due to an enlarged prostate gland, or had a stroke
- you are taking a diure ic
- you have problems or serious side e fects from taking pain relievers or fever reducers

Ask a doctor or pharmacist before use if you are
- under a doctor's care for any serious condition
- taking any other drug

When using this product
- take w th food or m lk f stomach upset occurs

LIFT HERE

LIFT HERE

This label may not be the latest approved by FDA.
For current labeling information, please visit https://www.fda.gov/drugsatfda

Drug Facts (continued)

Stop use and ask a doctor if
- you experience any of the following signs of stomach bleeding:
  - feel faint
  - vomit blood
  - have bloody or black stools
  - have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
  - chest pain
  - trouble breathing
  - weakness in one part or side of body
  - slurred speech
  - leg swe ling
- redness or swel ing is present in he painful area
- any new symptoms appear
- fever gets worse or lasts more than 3 days
- you have di ficulty swallowing or he caplet feels stuck in your throat
- you get nervous, dizzy, or sleepless
- nasal congestion lasts more than 7 days

If pregnant or breast-feeding, ask a hea th profess onal before use. It is especially important not to use naproxen sodium during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn ch ld or complications during delivery.

Drug Facts (continued)

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions
- **do not take more than directed**
- **the smallest effective dose should be used**
- **swallow whole;** do not crush or chew
- **drink a full glass of water w th each dose**
- adults and ch ldren 12 years and older: **1 caplet every 12 hours;** do not take more than 2 caplets in 24 hours
- children under 12 years: do not use

Other information
- **each caplet contains:** sodium 22 mg
- store at 20-25°C (68-77°F)
- store in a dry place

Inactive ingredients co loidal silicon dioxide, hyprome lose, lactose, magnesium stearate, microcrystal ine cellulose, polyethylene glycol, povidone, talc, itanium dioxide

Questions or comments? 1-800-986-0369 (Mon-Fri 9AM – 5PM EST)

Bayer

Distl. by:
Bayer HealthCare LLC
Whippany, NJ 07981

Bayer, the Bayer Cross, Aleve-D and All Day Strong are registered trademarks of Bayer.

XXXXX

3 258661569530

Do not use if the blister seal is cut or broken.

1 CAPLET - 12 HOURS

20 CAPLETS
CAPSULE-SHAPED TABLETS

ALL DAY STRONG®
ALEVE-D®
SINUS & COLD

12 HOUR MULTI-SYMPTOM RELIEF OF
SINUS PRESSURE
HEADACHE • SINUS CONGESTION
NASAL CONGESTION • BODY ACHES

naproxen sodium 220 mg (NSAID)*
Pseudoephedrine HCl 120 mg extended-release tablet
Pain reliever-fever reducer/nasal decongestant

BAYER

NON-DROWSY

ALL DAY STRONG®
ALEVE-D®
SINUS & COLD

: 2534D BY C.L

Drug Facts

Active ingredients (in each caplet)	Purposes
Naproxen sodium 220 mg (naproxen 200 mg) (NSAID)*	Pain reliever/fever reducer
Pseudoephedrine HCl 120 mg, extended-release	Nasal decongestant

*nonsteroidal anti-inflammatory drug

Uses
temporarily re ieves these cold, sinus, and flu symptoms:
- sinus pressure
- minor body aches and pains
- headache
- nasal and sinus conges ion (promotes sinus drainage and restores freer breathing hrough the nose)
- fever

Warnings
Allergy alert: Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:
- hives
- facial swe ling
- asthma (wheezing)
- shock
- skin reddening
- rash
- b isters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:
- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take o her drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoho ic drinks every day while using this product
- take more or for a longer time than directed

Drug Facts (continued)

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use
- f you have ever had an a lergic reaction to any o her pain re iever/fever reducer
- right before or after heart surgery
- f you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- in children under 12 years of age

Ask a doctor before use if
- the stomach bleeding warning app ies to you
- you have a history of stomach problems, such as hear burn
- you have high blood pressure, heart disease, iver cirrhosis, kidney disease, asthma, thyroid disease, diabetes, have trouble urinating due to an enlarged prostate gland, or had a stroke
- you are taking a diuretic
- you have problems or serious side effects from taking pain relievers or fever reducers

Ask a doctor or pharmacist before use f you are
- under a doctor's care for any serious cond tion
- taking any other drug

When using this product
- take with food or m lk f stomach upset occurs

LIFT HERE

LIFT HERE

This label may not be the latest approved by FDA.
For current labeling information, please visit https://www.fda.gov/drugsatfda

Drug Facts (continued)

Stop use and ask a doctor f

- you experience any of the following signs of stomach bleeding:
  - feel faint
  - vomit blood
  - have bloody or black stools
  - have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
  - chest pain
  - trouble breathing
  - weakness in one part or side of body
  - slurred speech
  - leg swelling
- redness or swe ling is present in the painful area
- any new symptoms appear
- fever gets worse or lasts more than 3 days
- you have d fficulty swa lowing or the caplet feels stuck in your throat
- you get nervous, dizzy, or sleepless
- nasal congestion lasts more than 7 days

If pregnant or breast-feeding, ask a hea th professional before use. It is especia ly important not to use naproxen sodium during the last 3 months of pregnancy unless de initely directed to do so by a doctor because t may cause problems in he unborn child or complica ions during delivery.

Drug Facts (continued)

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- **do not take more than directed**
- **the smallest effective dose should be used**
- **swallow whole;** do not crush or chew
- **drink a full glass of water with each dose**
- adults and children 12 years and older: **1 caplet every 12 hours;** do not take more than 2 caplets in 24 hours
- ch ldren under 12 years: do not use

Other information

- **each caplet contains:** sodium 22 mg
- store at 20-25°C (68-77°F)
- store in a dry place

Inactive ingredients colloidal s licon dioxide, hypromellose, lactose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, povidone, talc, titanium dioxide

Questions or comments? 1-800-986-0369
(Mon – Fri 9AM – 5PM EST)

Bayer

Do not use if the blister seal is cut or broken.

Dist. by:
Bayer HealthCare LLC
Whippany, NJ 07981

Bayer, the Bayer Cross, Aleve-D and All Day Strong are registered trademarks of Bayer.

XXXXX

3 25866 56951 6

1 CAPLET - 12 HOURS

10 CAPLETS
(CAPSULE-SHAPED TABLETS)

ALEVE-D®
SINUS & HEADACHE
ALL DAY STRONG®

12 HOUR MULTI-SYMPTOM RELIEF OF
SINUS PRESSURE
HEADACHE • SINUS CONGESTION
NASAL CONGESTION

BAYER

Naproxen sodium 220 mg (NSAID)
Pseudoephedrine HCl 120 mg extended-release tablet
Pain reliever-fever reducer/nasal decongestant

NON-DROWSY

ALL DAY STRONG®
ALEVE-D®
SINUS & HEADACHE

ALL DAY STRONG®
ALEVE-D®
SINUS & HEADACHE

C3 AB 25 h 25

Drug Facts

Active ingredients (in each caplet)	Purposes
Naproxen sodium 220 mg (naproxen 200 mg) (NSAID)*	Pain reliever/fever reducer
Pseudoephedrine HCl 120 mg, extended-release	Nasal decongestant

*nonsteroidal anti-inflammatory drug

Uses

temporar ly relieves hese cold, sinus, and flu symptoms:

- sinus pressure
- minor body aches and pains
- headache
- nasal and sinus congestion (promotes sinus drainage and restores freer breathing through the nose)
- fever

Warnings

Allergy alert: Naproxen sodium may cause a severe a lergic reaction, especia ly in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an a lergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher f you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Drug Facts (continued)

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart fa lure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- if you have ever had an allergic reac ion to any other pain reliever/fever reducer
- right before or after heart surgery
- if you are now taking a prescrip ion monoamine oxidase inhib tor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know f your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- in ch ldren under 12 years of age

Ask a doctor before use if

- the stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, thyroid disease, diabetes, have trouble urinating due to an enlarged prostate gland, or had a stroke
- you are taking a diure ic
- you have problems or serious side effects from taking pain relievers or fever reducers

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condi ion
- taking any other drug

When using this product

- take with food or m lk if stomach upset occurs

LIFT HERE

LIFT HERE

This label may not be the latest approved by FDA.
For current labeling information, please visit https://www.fda.gov/drugsatfda

---

This is a representation of an electronic record that was signed electronically and this page is the manifestation of the electronic signature.

/s/

VALERIE S PRATT
09/12/2017