




Labeling (prior approved)


Drug Facts (continued)	
Stop use and ask a doctor if ■ symptoms get worse ■ diarrhea lasts for more than 2 days	
If pregnant or breast-feeding, ask a health professional before use.	
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.	
Directions Drink plenty of clear fluids to help prevent dehydration caused by diarrhea. ■ Find right dose on chart. If possible, use weight to dose; otherwise, use age.	
adults and children 12 years and over	2 capsules after the first loose stool; 1 caplet after each subsequent loose stool; but no more than 4 capsules in 24 hours
children 9–11 years (50–95 lbs)	1 caplet after the first loose stool; 1/2 caplet after each subsequent loose stool; but no more than 2 capsules in 24 hours
children 5–8 years (40–55 lbs)	1 caplet after the first loose stool; 1/2 caplet after each subsequent loose stool; but no more than 2 capsules in 24 hours
children under 5 years (up to 47 lbs)	ask a doctor
Other Information Store between 20–25C (68–77F) ■ protect from light ■ do not use if pouch is open or torn	
Inactive Ingredients: polyethylene K, cellulose, dibasic calcium phosphate, flavor, sodium starch glycolate, white wax	
Questions or comments? Call 1-800-922-6343	
© 2000 Pharmacia Corporation All rights reserved. Pharmacia Corporation, Waukegan, IL 60087 Pharmacia Corporation is an Equal Opportunity Employer M/F/H/V	



150% OF SIZE



NDA 21-140
SLR 002
10/14/02
2-count single
pouch labeling
(Proposed)



Labelina Format Information:			
Fonts: Helvetica regular, bold and bold italic.			
Drug Facts:	NA	Leading:	4.7 pt
Header:	4.5 pt	Bullets:	3.5 pt
Subheader:	4.5 pt	Barlines:	NA
Body Text:	4.5 pt	Hairlines:	NA
Drug Facts (continued):	NA		
Horizontal Scale: 78%		Average Kerning: -10	

3003898 A- 1
AUGUST 27, 2002