



0 00000-00000-0000

Questions or comments? call 1-877-895-3665 (English) or 1-888-466-8746 (Spanish)

Inactive ingredients acesulfame potassium, croscarmellose sodium, dibasic calcium phosphate, flavor, microcrystalline cellulose, stearic acid

Other information ■ each caplet contains: calcium 165 mg, sodium 4 mg

■ see side panel for expiration date and lot number

■ do not use if carton is open or if blister unit is open or torn

■ store between 20-25°C (68-77°F). Protect from light.

Drug Facts (continued)

Directions ■ drink plenty of clear fluids to help prevent dehydration caused by diarrhea

■ take only on an empty stomach (1 hour before or 2 hours after a meal)

■ find right dose on chart. If possible, use weight to dose; otherwise, use age.

adults and children 12 years and over but no more than 4 caplets in 24 hours

children 9-11 years (60-95 lbs) 1 caplet after the first loose stool; 1/2 caplet after each subsequent loose stool; but no more than 3 caplets in 24 hours

children 6-8 years (45-59 lbs) 1 caplet after the first loose stool; 1/2 caplet after each subsequent loose stool; but no more than 2 caplets in 24 hours

children under 6 years (up to 47 lbs) ask a doctor

Warnings Do not use if you have bloody or black stool

Do not use if you have ever had a rash or other allergic reaction to loperamide HCl

Ask a doctor before use if you have

■ fever ■ mucus in the stool ■ a history of liver disease

Ask a doctor or pharmacist before use if you are taking antibiotics

When using this product tiredness, drowsiness or dizziness may occur. Be careful when driving or operating machinery.

Stop use and ask a doctor if

■ symptoms get worse ■ diarrhea lasts for more than 2 days

■ you get abdominal swelling or bloating. These may be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions ■ drink plenty of clear fluids to help prevent dehydration caused by diarrhea

■ take only on an empty stomach (1 hour before or 2 hours after a meal)

■ find right dose on chart. If possible, use weight to dose; otherwise, use age.

adults and children 12 years and over but no more than 4 caplets in 24 hours

children 9-11 years (60-95 lbs) 1 caplet after the first loose stool; 1/2 caplet after each subsequent loose stool; but no more than 3 caplets in 24 hours

children 6-8 years (45-59 lbs) 1 caplet after the first loose stool; 1/2 caplet after each subsequent loose stool; but no more than 2 caplets in 24 hours

children under 6 years (up to 47 lbs) ask a doctor

LOT

EXP

EASY TO OPEN



Drug Facts (continued)

Uses controls symptoms of diarrhea plus bloating, pressure, and cramps commonly referred to as gas

Warnings Allergy alert: Do not use if you have ever had a rash or other allergic reaction to loperamide HCl

Do not use if you have bloody or black stool

Ask a doctor before use if you have

■ fever ■ mucus in the stool ■ a history of liver disease

Ask a doctor or pharmacist before use if you are taking antibiotics

When using this product tiredness, drowsiness or dizziness may occur. Be careful when driving or operating machinery.

Stop use and ask a doctor if

■ symptoms get worse ■ diarrhea lasts for more than 2 days

■ you get abdominal swelling or bloating. These may be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Drug Facts

Active ingredients (in each caplet)

Loperamide HCl 2 mg

Simethicone 125 mg

Antidiarrheal/Anti-Gas

XXXXXX

000000000

Imodium

ADVANCED

Loperamide HCl, 2 mg/
Simethicone, 125 mg
Antidiarrheal/Anti-Gas

Controls the symptoms of

• diarrhea • bloating

• pressure and cramps

18 Caplets Easy to Swallow

000000000

000000000

000000000

000000000

Labeling Format Information:

Fonts: Custom regular, bold and bold italic

Drug Facts:	8.5 pt	Leading:	6.5 pt
Header:	8 pt	Bullets:	5 pt
Subheader:	6 pt	Barlines:	2.5 pt
Body Text:	6 pt	Hairlines:	.5 pt
Drug Facts (continued):	8 pt		

Horizontal Scale: 90%

Average Kerning: 0

DESCRIPTION CTN IMODIUM Advanced Caplet 18

FILE # SNDA 3007441

VERSION B-4 (SNDA)

DATE 09/06/07

Drug Facts	
Active ingredients (in each caplet)	Purposes
Loperamide HCl 2 mg Simethicone 125 mg	Anti-diarrhea Anti-gas
Uses controls symptoms of diarrhea plus bloating, pressure, and cramps commonly related to gas	
Warnings	
Allergy alert: Do not use if you have ever had a rash or other allergic reaction to loperamide HCl	
Do not use if you have bloody or black stool	
Ask a doctor before use if you have:	
■ fever	■ a history of liver disease
Ask a doctor or pharmacist before use if you are taking antibiotics	
When using this product, tiredness, drowsiness or dizziness may occur. Be careful when driving or operating machinery.	
Stop use and ask a doctor if:	
■ symptoms get worse	■ diarrhea lasts for more than 2 days
These may be signs of a serious condition.	
If pregnant or breast-feeding, ask a health professional before use.	
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away: (1-800-322-1222).	
Directions	
■ Drink plenty of clear fluids to help prevent dehydration caused by diarrhea	
■ Take only on empty stomach (1 hour before or 2 hours after a meal)	
■ Use only on empty stomach. If possible, use weight to dose. Otherwise, use age	
adults and children 12 years and over	2 caplets after the first loose stool, 1 caplet after each subsequent loose stool, but no more than 4 caplets in 24 hours
children 9-11 years (60-95 lbs)	1 caplet after the first loose stool, 1/2 caplet after each subsequent loose stool, but no more than 3 caplets in 24 hours
children 6-8 years (48-59 lbs)	1 caplet after the first loose stool, 1/2 caplet after each subsequent loose stool, but no more than 2 caplets in 24 hours
children under 6 years (up to 47 lbs)	ask a doctor

Drug Facts (continued)	
Other information	
■ Each caplet contains: loperamide 165 mg, sodium 4 mg	
■ Store between 20-25°C (68-77°F). Protect from light	
■ Do not use if caplet is open or if printed foil seal under bottle cap is open or torn	
■ See bottom panel for expiration date and lot number	
Inactive ingredients	
sodium lauryl sulfate, croscarmellose sodium, dicalcium phosphate, lactose, microcrystalline cellulose, stearic acid	
Questions or comments?	
call 1-877-895-5655 (English) or 1-888-466-6746 (Spanish)	
The makers of Imodium® do not manufacture store brands.	

Imodium®	
ADVANCED	
Controls the symptoms of • diarrhea • bloating • pressure and cramps	
30 Caplets Easy to Swallow	
THIS PACKAGE FOR HOUSEHOLDS WITHOUT YOUNG CHILDREN NDC 50586-XXX-XX	
Easy to Open Bottle	
Loperamide HCl 2 mg/ Simethicone 125 mg Antidiarrheal/Anti-Gas	

Imodium®	
ADVANCED	
Controls the symptoms of • diarrhea • bloating • pressure and cramps	
PATENTED FORMULA	
0 0000-0000-00 0	
Distributed by: McNeil Consumer Healthcare DIVISION OF MCNEIL-PPC, INC. FORT WASHINGTON, PA 19074 USA © MCNEIL-PPC, Inc. 07 www.imodium.com U.S. Patent No. 6,105,260	

Labeling Format Information:

Fonts: Custom condensed medium, bold and bold italic

Drug Facts: 8.25 pt

Header: 8 pt

Subheader: 6 pt

Body Text: 6 pt

Drug Facts (continued): 8 pt

Leading: 6.5 pt

Bullets: 5 pt

Barlines: 2.5 pt

Hairlines: .5 pt

DESCRIPTION CTN IMODIUM Advanced Caplet 30 NCR

FILE # SMDA 3007441

DATE 9/06/07

THIS PACKAGE FOR HOUSEHOLDS WITHOUT YOUNG CHILDREN

NDC 50580-XXX-XX

Imodium®

ADVANCED

Loperamide HCl, 2 mg/
Simethicone, 125 mg
Antidiarrheal/Anti-Gas

30 Caplets

Purposes

Active ingredients
(in each caplet)
Loperamide HCl 2 mg.....Anti-diarrheal
Simethicone 125 mg.....Anti-gas
Uses controls symptoms of diarrhea plus
bloating, pressure, and cramps commonly
referred to as gas

LOT

EXP

00000000

Ask a doctor or pharmacist before use if you are taking antibiotics. When using this product tiredness, drowsiness or dizziness may occur. Be careful when driving or operating machinery. Stop use and ask a doctor if - symptoms get worse - diarrhea lasts for more than 2 days - you get abdominal swelling or bulging. These may be signs of a serious condition. If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222) Directions • drink plenty of clear fluids to help prevent dehydration caused by diarrhea • take only on an empty stomach (1 hour before or 2 hours after a meal) • find right dose on chart. If possible, use weight to dose; otherwise, use age

adults and children 12 years and over	2 caplets after the first loose stool; 1 caplet after each subsequent loose stool; but no more than 4 caplets in 24 hours
children 9-11 years (60-95 lbs)	1 caplet after the first loose stool; 1/2 caplet after each subsequent loose stool; but no more than 3 caplets in 24 hours
children 6-8 years (48-59 lbs)	1 caplet after the first loose stool; 1/2 caplet after each subsequent loose stool; but no more than 2 caplets in 24 hours
children under 6 years (up to 47 lbs)	ask a doctor

Other information • each caplet contains: calcium 165 mg, sodium 4 mg • store between 20-25°C (68-77°F). Protect from light. • do not use if printed foil seal under bottle cap is open or torn

Questions or comments? call 1-877-895-3665 (English) or 1-888-466-8746 (Spanish)

Dist. by: MCNEIL-PPC, INC., FORT WASHINGTON, PA 19034 USA © MCN-PPC, Inc. 07 U.S. Patent No. 6,103,260

200% OF SIZE

THIS PACKAGE FOR HOUSEHOLDS WITHOUT YOUNG CHILDREN

NDC 50580-XXX-XX

Imodium®

ADVANCED

Loperamide HCl, 2 mg/
Simethicone, 125 mg
Antidiarrheal/Anti-Gas

30 Caplets

Purposes

Active ingredients
(in each caplet)
Loperamide HCl 2 mg.....Anti-diarrheal
Simethicone 125 mg.....Anti-gas
Uses controls symptoms of diarrhea plus
bloating, pressure, and cramps commonly
referred to as gas

LOT

EXP

00000000

Ask a doctor or pharmacist before use if you are taking antibiotics. When using this product tiredness, drowsiness or dizziness may occur. Be careful when driving or operating machinery. Stop use and ask a doctor if - symptoms get worse - diarrhea lasts for more than 2 days - you get abdominal swelling or bulging. These may be signs of a serious condition. If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222) Directions • drink plenty of clear fluids to help prevent dehydration caused by diarrhea • take only on an empty stomach (1 hour before or 2 hours after a meal) • find right dose on chart. If possible, use weight to dose; otherwise, use age

adults and children 12 years and over	2 caplets after the first loose stool; 1 caplet after each subsequent loose stool; but no more than 4 caplets in 24 hours
children 9-11 years (60-95 lbs)	1 caplet after the first loose stool; 1/2 caplet after each subsequent loose stool; but no more than 3 caplets in 24 hours
children 6-8 years (48-59 lbs)	1 caplet after the first loose stool; 1/2 caplet after each subsequent loose stool; but no more than 2 caplets in 24 hours
children under 6 years (up to 47 lbs)	ask a doctor

Other information • each caplet contains: calcium 165 mg, sodium 4 mg • store between 20-25°C (68-77°F). Protect from light. • do not use if printed foil seal under bottle cap is open or torn

Questions or comments? call 1-877-895-3665 (English) or 1-888-466-8746 (Spanish)

Dist. by: MCNEIL-PPC, INC., FORT WASHINGTON, PA 19034 USA © MCN-PPC, Inc. 07 U.S. Patent No. 6,103,260

Labeling Format Information:

Fonts: Custom condensed medium, bold and bold italic

Drug Facts: NA

Header: 4.5 pt

Subheader: 4.5 pt

Body Text: 4.5 pt

Drug Facts (continued): NA

Horizontal Scale: 80%

Average Kerning: 0

DESCRIPTION

FILE #

DATE

LBL IMODIUM Advanced Caplet 30 NCR

SNDA 3007441

9/07/07

VERSION

E-5 (SNDA)