

TEAR OFF A SECTION

FOLD CORNER INWARD AND PEEL BACK FOIL

Drug Facts (continued)

Other Information

- each caplet contains: calcium 165 mg, sodium 3 mg
- store between 20-25°C (68-77°F). Protect from light.

Drug Facts

Active Ingredients (in each caplet)

Loperamide HCl, 2 mg
Simethicone 125 mg

Purposes

Antidiarrheal
Anti-gas

Directions

Drink plenty of clear fluids to help prevent dehydration caused by diarrhea

Take only on an empty stomach (1 hour before or 2 hours after a meal)

Take with a full (8 oz) glass of water

Find right dose on chart below. If possible, use weight to dose.

adults and children 17 years and over

2 caplets after the first loose stool; 1 caplet after each subsequent loose stool but no more than 3 caplets in 24 hours

children 9-11 years (60-95 lbs)

1 caplet after the first loose stool; 1/2 caplet after each subsequent loose stool; but no more than 3 caplets in 24 hours

children 6-8 years (44-59 lbs)

1 caplet after the first loose stool; 1/2 caplet after each subsequent loose stool; but no more than 2 caplets in 24 hours

ask a doctor

children under 2 years (14 to 47 lbs)

do not use

Drug Facts (continued)

do not use if carton is open or if blister unit is open or torn

Inactive Ingredients

acetylsalicylic acid, croscarmellose sodium, dibasic calcium phosphate, flavor, microcrystalline cellulose, stearic acid

Questions or comments?

call 1-877-895-3665 (toll-free) or 215-273-8753 (collect)

Imodium
How can we help?
1-877-895-3665

See New Warning and Directions

Imodium
Loperamide HCl, 2 mg/Simethicone, 125 mg Antidiarrheal/Anti-gas

Multi-Symptom Relief
Caplet

Relieves symptoms of diarrhea *plus*

- Cramps & Pressure
- Bloating
- Gas

12 Caplets*

*capsule-shaped tablets

Actual Size

Care To Recycle™

Lot

EXP

Distributed by:
JOHNSON & JOHNSON CONSUMER INC.
McNeil Consumer Healthcare Division
Fort Washington, PA 19034 USA
© J&J 2016
www.imodium.com
U.S. Patent No. 6,033,260

TEAR OFF SECTION

FOLD CORNER INWARD AND PEEL BACK FOIL

Drug Facts (continued)

Active ingredients (in each caplet)

Loperamide HCl 2 mg
Simethicone 125 mg
Anti-diarrheal
Anti-gas

Purposes

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions

Drink plenty of clear fluids to help prevent dehydration caused by diarrhea.

Take only on an empty stomach (1 hour before or 2 hours after a meal).

Take with a full (8 oz.) glass of water.

Find right dose on chart below. If possible, use weight to dose; otherwise, use age.

Age	Dose
Adults and children 12 years and over	2 caplets after the first loose stool; 1 caplet after each subsequent loose stool; but no more than 4 caplets in 24 hours.
Children 9-11 years (60-95 lbs)	1 caplet after the first loose stool; 1/2 caplet after each subsequent loose stool; but no more than 3 caplets in 24 hours.
Children 6-8 years (48-59 lbs)	1 caplet after the first loose stool; 1/2 caplet after each subsequent loose stool; but no more than 3 caplets in 24 hours.
Children 2-5 years (24 to 47 lbs)	ask a doctor.
Children under 2 years (up to 33 lbs)	do not use.

Other information

- Each caplet contains: calcium 165 mg, sodium 3 mg.
- Store between 20-25°C (68-77°F). Protect from light.
- Do not use if carton is open or if blister unit is open or torn.

Drug Facts (continued)

Inactive ingredients

acetylsalicylic acid, croscarmellose sodium, dibasic calcium phosphate, lactose, microcrystalline cellulose, stearic acid.

Questions or comments?

Call 1-877-895-3665 (toll-free) or 215-273-8755 (collect).

00000000

Imodium®

How can we help?

1-877-895-3665

See New Warning and Directions

Imodium®

Loperamide HCl, 2 mg/Simethicone, 125 mg Antidiarrheal/Anti-gas

Multi-Symptom Relief Caplet

Relieves symptoms of diarrhea *plus*

- Cramps & Pressure
- Bloating
- Gas

18 Caplets*

*capsule-shaped tablets

Actual Size

00000000

TEAR OFF A SECTION

FOLD AND TEAR AT NOTCH

Drug Facts (continued)

Active ingredients (in each caplet)
Loperamide HCl 2 mg
Simethicone 125 mg
Anti-diarrheal
Anti-gas

Purposes
In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions
● Drink plenty of clear fluids to help prevent dehydration caused by diarrhea
● Take only on an empty stomach (1 hour before or 2 hours after a meal)
● Take with a full (8 oz.) glass of water
● Indigestion, use age appropriate dose on chart below. If possible, use weight to dose.

Warnings
● Heart alert: Taking more than directed can cause serious heart problems or death
● Allergy alert: Do not use if you have ever had a rash or other allergic reaction to loperamide HCl
● Do not use if you have difficulty swallowing
● If you have bloody or black stool
● If you have a history of liver disease
● Ask a doctor or pharmacist before use if you are taking antibiotics

When using this product tiredness, drowsiness or dizziness may occur. Be careful when driving or operating machinery.

Stop use and ask a doctor if
● symptoms get worse
● you get abdominal swelling or bloating. These may be signs of a serious condition
● you have difficulty swallowing the caplet

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Other information
● Each caplet contains calcium 165 mg, sodium 5 mg
● Store between 20-25°C (68-77°F). Protect from light.
● Do not use if carton is open or if blister unit is open or torn

Drug Facts (continued)

Inactive ingredients
acetylsalicylic acid, croscarmellose sodium, dibasic calcium phosphate, lactose, microcrystalline cellulose, stearic acid

Questions or comments?
Call 1-877-895-3665 (toll-free) or 215-273-8755 (collect)

Care To Recycle

Distributed by:
**JOHNSON & JOHNSON
CONSUMER INC.**
McNeil Consumer Healthcare Division
Fort Washington, PA 19034 USA
©J&JCI 2016
www.imodium.com
U.S. Patent No. 6,103,260

00000000

Imodium®
How can we help?
1-877-895-3665

See New Warning and Directions

Imodium®
Loperamide HCl, 2 mg/Simethicone, 125 mg Antidiarrheal/Anti-gas

Multi-Symptom Relief
Caplet

Relieves symptoms of diarrhea *plus*

- Cramps & Pressure
- Bloating
- Gas

18 Caplets*

*capsule-shaped tablets Actual Size

00000000

00000000

NDC 50580-338-30

Loperamide HCl, 2 mg/Simethicone, 125 mg Antidiarrheal/Anti-gas

*capsule-shaped tablets

Actual Size

Loperamide HCl, 2 mg/Simethicone, 125 mg Anti-diarrheal/Anti-gas



Imodium®
How can we help?
1-877-895-3665

Imodium

 Care To Recycle™

Drug facts (continued)

Inactive ingredients
acetaminophen potassium, croscarmellose sodium, croscarmellose, stearic acid

Questions or comments?
call 1-877-895-3665 (toll-free) or
215-273-8755 (collect)

Other information

- each caplet contains: calcium 165 mg, sodium 3 mg
- store between 20-25°C (68-77°F). Protected from light
- do not use if carton is open or if printed foil seal under bottle cap is open or torn

ask a doctor	children 2-5 years
do not use	children under 2 years (up to 33 lbs)



3 0045-0212-306

3 0045-0212-306

Drug Facts (continued)	adults and children	children 12 years and over	children 9-11 years (60-95 lbs)	children 6-5 years (40-59 lbs)
2 capsules after the first loose stool.	1 capsule after each subsequent loose stool, but no more than 4 capsules in 24 hours.	1 capsule after the first loose stool.	1 capsule after each subsequent loose stool, but no more than 2 capsules in 24 hours.	1 capsule after the first loose stool.
2 capsules after the first loose stool.	1 capsule after each subsequent loose stool, but no more than 4 capsules in 24 hours.	1 capsule after the first loose stool.	1 capsule after each subsequent loose stool, but no more than 2 capsules in 24 hours.	2 capsules in 24 hours

Do not use (continued)

- If you have bloody or black stool
- If you have difficulty swallowing
- Ask a doctor before use if you have:
 - a history of liver disease
 - muscle in the stool
 - Ask a doctor or pharmacist before use if you are taking antibiotics
- When using this product (tiredness, drowsiness or dizziness may occur. Be careful when driving or operating machinery).
- Stop use and ask a doctor if:**
 - Symptoms get worse
 - Diarrhea lasts for more than 2 days
 - You get bad abdominal cramping or bloating
 - These men get signs of a serious condition.
 - You have difficulty swallowing the caplet.

Drug Facts (continued)

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222).

Directions

• drink plenty of clear fluids to help prevent dehydration caused by diarrhea

• take only on an empty stomach (1 hour before or 2 hours after a meal)

• take with a full glass (8 oz.) of water

• limit night use (if possible, use weight to dose otherwise use age)

Drug Facts	Active ingredients	Purposes
Oprenamide HCl (in each caplet) Smethicone 125 mg Anti-diemetic Anti-gas	Uses relieves symptoms of diarrhea plus bloating, pressure and cramps, commonly referred to as gas	Warnings Allergy alert: Do not use if you have ever had a allergic reaction to oprenamide HCl Heart alert: Taking more than directed can cause serious heart problems or death

00000000

LOT:

EXP:

00000000

LOT:

EXP:

Active ingredients (in each caplet):
Loperamide HCl 2 mg.....Anti-diarrheal
Simethicone 25 mg.....Anti-gas
Uses: relieves symptoms of diarrhea plus bloating,
pressure and cramps. Commonly referred to as gas.
Warnings: Allergy alert: Do not use if you have

Purposes:
Anti-diarrheal
Anti-gas

NDC 50580-338-30

Imodium

Loperamide HCl 2mg/Simethicone 125mg

Antidiarrheal/Anti-gas

Multi-Symptom Relief

30 Caplets*

*capsule-shaped tablets

Ask a doctor or pharmacist before use. If you are taking anti-biotics. When using this product, dizziness, weakness or dizziness may occur. Be careful when driving or operating machinery. Stop use and ask a doctor if symptoms get worse. Diarrhea lasts for more than 2 days. You get abdominal swelling or bulging. These may be signs of a serious condition. You have difficulty swallowing the caplet. If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222) Directions: Drink plenty of clear fluids to help prevent dehydration caused by diarrhea. Take only on an empty stomach (1 hour before or 2 hours after a meal). Take with a full (8 oz.) glass of water. If possible, use weight to dose; otherwise, use age. Adults and children 12 years and over: 2 caplets after the first loose stool; 1 caplet after each subsequent loose stool, but no more than 4 caplets over 24 hours. Children 9-11 years (60-95 lbs): 1 caplet after the first loose stool; 1/2 caplet after each subsequent loose stool, but no more than 3 caplets in 24 hours. Children 6-8 years (48-59 lbs): 1 caplet after the first loose stool; 1/2 caplet after each subsequent loose stool, but no more than 2 caplets in 24 hours. Children 2-5 years (34 to 47 lbs): ask a doctor. Children under 2 years (up to 33 lbs) do not use. Other information: Each caplet contains calcium 165 mg, sodium 3 mg. Store between 20-25°C (68-77°F). Protect from light. Do not use if printed foil seal on bottle is broken or missing. Questions or comments? Call 1-877-895-3665 (toll-free) or 215-273-8755 (collect). Distributed by: JOHNSON & JOHNSON CONSUMER INC., McNeil Consumer Healthcare Division, Fort Washington, PA 19034 USA ©J&JCI 2016 www.imodium.com U.S. Patent No. 6,103,260

Reference ID: 4083274

Loperamide HCl, 2 mg/Simethicone, 125 mg Antidiarrheal/Anti-gas

30 Caplets*

*capsule-shaped tablets

Actual Size



Loperamide HCl, 2 mg/Simethicone, 125 mg Antidiarrheal/Anti-gas



Imodium

How can we help?
1-877-895-3665

 Care To Recycle™

Questions or comments?
Call 1-877-895-3665 (toll-free) or
215-273-8755 (collect)

[illegible]

- each caplet contains: calcium 165 mg, sodium 5 mg
- store between 20-25°C (68-77°F); protect from light
- do not use if carton is open or if printed foil seal under bottle cap is open or torn

ask a doctor	children 2-5 years (34 to 47 bs)	children under 2 years (up to 33 bs)
do not use		



3 0045-0212-306

[illegible]

Drug Facts (continued)

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-232-1222).

Directions

• drink plenty of clear fluids to help prevent dehydration caused by diarrhea

• take only on an empty stomach (1 hour before or 2 hours after a meal)

• take with at least 8 (oz.) glass of water

• may fight off some on-chest / flu viruses, use weight to dosage. (check with your doctor)

• to dosage. (check with your doctor)

Do not use if: • If you have a bloody or black stool
• If you have difficulty swallowing
• If you have an ulcer
• If you have a history of liver disease
• If you are taking other drugs
• If you are taking this product (headache, drowsiness or dizziness may occur. Be careful when driving or operating machinery.)

Stop use and ask a doctor if:
• Symptoms get worse
• Diarrhea lasts for more than 2 days
• You get double vision or blurring
• These may be signs of a serious condition.
• You have difficulty swallowing the caplet.

Drug Facts	Active ingredients Loperamide HCl 2 mg (in each capsule) Simethicone 125 mg, Anti-diemetic Anti-gas
Purposes	Uses relieves symptoms of diarrhea plus bloating, pressure and cramps, commonly relieved to gas
Warnings	Alerts Do not use if you have ever had a heart or other allergic reaction to loperamide HCl or other heart products that directed can cause serious heart problems or death

Active ingredients (in each caplet)
Loperamide HCl 2 mg
Simethicone 125 mg

Purposes
Anti-diarrheal
Anti-gas

Uses • relieves symptoms of diarrhea plus bloating, pressure and cramping, commonly referred to as gas

Warnings: Allergy alert: Do not use if you have ever had a rash or other allergic reaction to loperamide HCl. **Heart alert:** Taking more than directed can cause serious heart problems or death. **Do not use** • if you have bloody or black stool • if you have difficulty swallowing. **Ask a doctor before use if you have** • a fever • muscle pain in the stool • a history of liver disease. **Ask a doctor or pharmacist before use if you are** taking antibiotics. **When using this product** (tiredness, drowsiness or dizziness may occur. Be careful when driving or operating machinery. **Stop use and ask a doctor if symptoms get worse** • diarrhea lasts for more than 2 days • you get abdominal swelling or bulging. These may be signs of a serious condition. • you have difficulty swallowing the caplet. **If pregnant or breastfeeding**, ask a health professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-2222)

Multi-Symptom Relief
30 Caplets*
*capsule-shaped tablets

Directions • drink plenty of clear fluids to help prevent dehydration caused by diarrhea • take only on an empty stomach (1 hour before or 2 hours after a meal) • take with a full (8 oz.) glass of water • find right dose on chart. If possible, use weight dose; otherwise, use age.

adults and children 12 years and over	2 caplets after the first loose stool; 1 caplet after each subsequent loose stool; but no more than 4 caplets in 24 hours
children 9-11 years (60-95 lbs)	1 caplet after the first loose stool; 1/2 caplet after each subsequent loose stool; but no more than 3 caplets in 24 hours
children 6-8 years (48-99 lbs)	1 caplet after the first loose stool; 1/2 caplet after each subsequent loose stool; but no more than 2 caplets in 24 hours
children 2-5 years (34 to 47 lbs)	ask a doctor
children under 2 years (up to 33 lbs)	do not use

Other information • each caplet contains: calcium 165 mg, sodium 3 mg • store between 20-25°C (68-77°F). Protect from light. • **do not use if printed foil seal on bottle is broken or missing**
Questions or comments? call 1-877-895-3665 (toll-free) or 215-273-8755 (collect)

EXP
LOT

Distributed by: **JOHNSON & JOHNSON CONSUMER INC.**
Mandel Center for Healthcare Division
First Avenue, PA 19044 USA © J&J 2016 USN 032480



LOT:

EXP:

0.000000

Acting ingredients (in each caplet)

Simethicone 2 mg.....Anti-dar healing

Simethicone 25 mg.....Anti-gas

Uses: relieves symptoms of dar hea plus bloating, pressure and cramps. commonly referred to as gas

Warnings: Allergy alert: Do not use if you have

Purposes

Anti-dar healing

Anti-gas

Uses: relieves symptoms of dar hea plus bloating, pressure and cramps. commonly referred to as gas

Warnings: Allergy alert: Do not use if you have

NDC 50580 338 42

Imodium

Loperamide HCl 2mg/Simethicone, 125 mg

Antidiarrheal/Anti-gas

Multi-Symptom Relief

42 Caplets*

*capsule-shaped tablets

ever had a rash or other allergic reaction to loperamide HCl. Heart alert: Taking more than directed can cause serious heart problems or death. Do not use if you have bloody or black stool or if you have difficulty swallowing. Ask a doctor before use if you have liver problems or a history of liver disease.

Ask a doctor or pharmacist before use. If you are taking anti-biotics. When using this product, drowsiness or dizziness may occur. Be careful when driving or operating machinery. Stop use and ask a doctor if symptoms get worse or if diarrhea lasts for more than 2 days or if you get abdominal swelling or bloating. These may be signs of a serious condition. If you have difficulty swallowing the caplet, if you are pregnant or breastfeeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-2222) Directions: Drink plenty of clear fluids to help prevent dehydration caused by diarrhea. Take only on an empty stomach (1 hour before or 2 hours after a meal). Take with a full (8 oz.) glass of water. If possible, use weight to dose; otherwise, use age. Adults and children 12 years and older: 2 caplets after the first loose stool; 1 caplet after each subsequent loose stool; but no more than 4 caplets in 24 hours. Children 9-11 years (60-95 lbs): 1 caplet after the first loose stool; 1/2 caplet after each subsequent loose stool; but no more than 3 caplets in 24 hours. Children 6-8 years (48-59 lbs): 1 caplet after the first loose stool; 1/2 caplet after each subsequent loose stool; but no more than 2 caplets in 24 hours. Children 2-5 years (34 to 47 lbs) ask a doctor. Children under 2 years (up to 33 lbs) do not use.

Other information: Each caplet contains calcium 165 mg, sodium 3 mg. Store between 20-25°C (68-77°F). Protect from light. Do not use if printed foil seal on bottle is broken or missing.

Questions or comments? Call 1-877-895-5665 (toll-free) or 215-273-0755 (collect).

Distributed by: JOHNSON & JOHNSON CONSUMER INC.

McNeil Consumer Healthcare Division, Fort Washington, PA 19034 USA ©J&JCI 2016

www.imodium.com U.S. Patent No. 6,103,280

Reference ID: 4083274



Reference ID: 4083274

This is a representation of an electronic record that was signed electronically and this page is the manifestation of the electronic signature.

/s/

VALERIE S PRATT
04/12/2017