

page 1

Trivagizole 3™

Clotrimazole Vaginal Cream (2%)
 Vaginal Antifungal

Vaginal Cream

3 Day Treatment

1 Reusable Applicator

Trivagizole 3™

Clotrimazole Vaginal Cream (2%)
 Vaginal Antifungal

Vaginal Cream

Cures Most Vaginal Yeast Infections

Trivagizole 3™

Clotrimazole Vaginal Cream (2%)
 Vaginal Antifungal

Vaginal Cream

One 25 g (0.88 oz) Tube Of Clotrimazole Vaginal Cream (2%)
 and 1 Reusable Applicator

Cures Most
 Vaginal Yeast
 Infections

3 Day Treatment



Educational Brochure Enclosed



Drug Facts (continued)

Other information

- Store at 20-25°C (68-77°F)
- See and keep cap of carton and tube for number and expiration date
- The tube opening should be sealed. If seal has been broken, do not use product.

Inactive ingredients benzyl alcohol, cetostearyl alcohol, cetyl esters wax, 2-octyldodecanol, polyorbate-60, purified water, sorbitan monooleate

Questions or comments? Call toll-free, 1-888-827-8222, for our 24 hour automated response system.

PK-0000-0
 M000

TARO

TARO

TARO

Distributed by:
 TARO Pharmaceuticals,
 U.S.A., Inc.
 Hawthorne, NY 10532
 Made in Canada.
 TARO Pharmaceuticals U.S.A., Inc.
 is a registered trademark of
 TARO Pharmaceuticals U.S.A., Inc.
www.taropharma.com

Drug Facts

Active ingredient (in each applicator)
 Clotrimazole 100 mg (2%)
 Vaginal antifungal

Purpose
 Vaginal antifungal

Use
 Use treats vaginal yeast infections

Warnings
 For vaginal use only

Do not use

- If you have never had a vaginal yeast infection diagnosed by a doctor
- If you have a fever (higher than 100°F), pain in the lower abdomen, back, or other shoulder, or a four-smelling vaginal discharge. You should see a doctor for treatment of these symptoms.
- Ask a doctor before use if you have:
 - vaginal itching and discomfort for the first time
 - symptoms that return within 2 months. You could be pregnant, or have a serious underlying medical cause for your symptoms, including diabetes or a weakened immune system (which may be due to HIV - the virus that causes AIDS).

When using this product

- do not use tampons, douches, or spermicides
- do not use condoms or diaphragms, as this product may damage them and they may not prevent sexually transmitted disease or pregnancy

Stop use and ask a doctor if

- symptoms do not get better in 3 days
- you get a rash, fever, abdominal pain, or a four-smelling vaginal discharge

Directions

- Before using, read and keep the enclosed educational brochure for complete instructions and information
- adults and children 12 years and over:
 - insert one applicator of cream into the vagina at bedtime for 3 days in a row
 - children under 12 years: ask a doctor

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Graphic leading to ne

2.5 point box bar
 0.5 point barline
 2.5 point barline

5 pt. Helvetica Regular

5 pt. square

Helvetica Bold

Helvetica Bold

6.5 pt. leading, left justified


left justified

Helvetica Bold

APPROVAL		
DATE	3/22/00	FIG
		ANT COPY CTRL. 0

Trivagizole 3™ Clotrimazole Vaginal Cream (2%) Vaginal Antifungal 3 Day Treatment 1 Reusable Applicator Cures Most Vaginal Yeast Infections

page



TARO

Trivagizole³™

Clotrimazole Vaginal Cream (2%)
 Vaginal Antifungal

3 Day Treatment (Weight 25g (0.88oz))

For vaginal use only. Do not use in eyes or take by mouth.
 If this is the first time you have had vaginal itching and discomfort, consult your doctor. If you have had a doctor diagnose a vaginal yeast infection before and have the same symptoms now, use this cream as directed for 3 days in a row.
Warnings: Read the Warnings and Directions on the carton and in the educational brochure before using this product.
 Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.
Directions: To Open: Unseal the cap, use pointed end on cap to break seal. Fill the applicator and insert one applicatorful of cream into vagina at bedtime for 3 days in a row.
Active Ingredient: Clotrimazole 2% (100 mg per applicator)
 Store at 20-25 °C (68-77 °F).
 See end of tube for lot number and expiration date.
Safety Sealed: The tube opening should be sealed. If seal has been broken, do not use product.
 Mfd. by: Taro Pharmaceuticals Inc.
 Bramaleo, Ontario, Canada L6T 1C3 Made in Canada PK-0000-0

● PMS 281

●

APPROVAL					
DATE		LPE#		ART COPY CTRL.#	
PMS#		UPC#		PHARMACIST	
DEPARTMENT	SIGNATURE	NO	YES	YES W CORR	DATE
PROOFING					
ART					
MARKETING					
ENGINEERING					
Q.C.					
REG.					
CLIENT					

THIS COLOR PROOF INDICATES COLOR BREAK ONLY AND MAY NOT ACCURATELY REFLECT ACTUAL PRODUCTION COLOR. IF YOU HAVE ANY QUESTIONS OR COMMENTS PLEASE CALL TARO PHARMACEUTICALS-ART DEPARTMENT 800-844-1448.

100%

ID BAR CODE

EDUCATIONAL BROCHURE

Read This Before Using

Trivagizole 3™

Clotrimazole

Vaginal

Cream (2%)

Vaginal Antifungal

Clotrimazole 2% (200mg per applicator)

3-day treatment of vaginal yeast infections.

Use:

• For the treatment of vaginal yeast infections.

If you have symptoms such as vaginal itching, burning or discharge, and if at some time in the past your doctor has told you that these symptoms were due to a vaginal yeast infection, then Trivagizole 3 Vaginal Cream should work for you. If, however, you have never had these symptoms or a vaginal yeast infection before, you should see your doctor before using this product.

What is a Vaginal Yeast Infection?

A yeast infection is a common type of vaginal infection. Your doctor may call it candidiasis. This condition is caused by an

organism called Candida, which is a type of yeast.

Healthy women usually have this yeast on the skin, in the mouth, in the digestive tract and in the vagina. At times, the yeast can grow very quickly. In fact, the infection is sometimes called yeast (Candida) overgrowth. Some women may have a yeast infection on the skin outside the vagina (yuck), associated with the vaginal infection.

A yeast infection can occur at any age. It is most common during the childbearing years. Women who are pregnant or diabetic, taking antibiotics, birth control pills or steroids, or who have a weakened immune system are more likely to get repeated yeast infections.

Various medical conditions can damage the body's normal defenses against

infection. One of the most serious of these conditions is

infection with the human immunodeficiency virus (HIV), the virus that causes AIDS. The HIV virus causes the body to be more likely to get infections, including vaginal yeast infections. Therefore, women with HIV infection may have frequent vaginal yeast infections that do not clear up easily with proper treatment. If you may have been exposed to HIV and are experiencing repeated vaginal yeast infections, you should see your doctor right away. For additional information on HIV infection, please contact your doctor or the CDC National AIDS Hotline. The CDC phone numbers are: 1-800-342-AIDS (English), 1-800-344-7432 (Spanish), or 1-800-243-7889 (hearing impaired, TDD).

Symptoms of a Vaginal Yeast Infection

There are many symptoms of a vaginal yeast infection. They can include:

- Vaginal itching (ranging from mild to severe)
- A clumpy, vaginal discharge that may look like cottage cheese.
- Vaginal soreness, irritation or burning.

• Rash or redness on the skin outside the vagina (usually irritation).

Note: Vaginal discharge that is different from above, for example, a yellow/green discharge or a foul-smelling discharge may mean that you have a different infection. If this is the case, you should talk to your doctor before using Trivagizole 3 Vaginal Cream.

Warnings

• For vaginal use only. Do not use in eyes or take by mouth.

• This product is effective only in treating vaginal yeast infections.

• Do not use Trivagizole 3 Vaginal Cream if you have any of the following symptoms:

- Fever (higher than 100°F).
- Pain in the lower abdomen, back, or either shoulder.
- A foul-smelling vaginal discharge.

• While using Trivagizole 3 Vaginal Cream, if you get a fever, abdominal pain, or a foul-smelling vaginal discharge, stop

using the product & contact your doctor at once. You may have more serious disease

• If your symptoms do not improve in 3 days, you still have symptoms after 7 days, you should call your doctor.

• If your symptoms (e.g. within 2 months) should talk to your doctor. You could be pregnant or there could be a serious underlying medical cause for your infection, including diabetes or a weakened immune system (which may be due to HIV, the virus that causes AIDS).

• Contact your doctor if you get a fever, abdominal pain, or a foul-smelling vaginal discharge, stop

4-10 CLOTRIMAZOLE 2% CR, 3-DAY (TARO) - INSERT - FRONT 4 X 12 FOLDED TO 1 X DIE IS AT 100%

COLORS...
PMS-291
Date 3/21/00

page 3

pink

Warnings

- For vaginal use only. Do not use in eyes or take by mouth.
- This product is effective only in treating vaginal yeast infections.
- Do not use Trivagizole 3 Vaginal Cream if you have any of the following symptoms:
 - fever (higher than 100°F);
 - pain in the lower abdomen, back, or either shoulder;
 - a foul-smelling vaginal discharge.
- While using Trivagizole 3 Vaginal Cream, if you get a fever, abdominal pain, or a foul-smelling vaginal discharge, stop using the product and contact your doctor right away. You may have a more serious disease.
- If your symptoms do not improve in 3 days, or you still have symptoms after 7 days, you should call your doctor.
- If your symptoms return within 2 months, you should talk to your doctor. You could be pregnant or there could be a serious underlying medical cause for your infections, including diabetes or a weakened immune system (which may be due to HIV - the virus that causes AIDS).
- Contact your doctor if you get hives or a skin rash while using this product.
- Do not use tampons, douches, or spermicides while using this medicine.
- Do not rely on condoms or diaphragms to prevent sexually transmitted diseases or pregnancy. This product may damage condoms and diaphragms and cause them to fail.
- If pregnant or breastfeeding, ask a health professional before use.
- Do not use in girls less than 12 years of age.
- Keep this and all drugs out of the reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

INSERT - FRONT 4 X 12 FOLDED TO 1 X 1/2

COLORS ...
 PMS 291
 Date 3/21/00

Boxes print
 8% BD
 PMS 281


page 4

ID BAR CODE


2. Attach the applicator to the tube of cream by placing end of the applicator firmly onto the orifice of cream. (See picture.)




3. Gently squeeze the cream into the applicator. Continue squeezing until the cream is fully loaded. (See picture.)




4. Hold the barrel as shown in the picture.



5. The Trivagrole 3 Vaginal Cream applicator is inserted into the vagina as much as the same size as a tampon. Gently insert the applicator into the vagina as far back as it will go comfortably. This can be done while lying on your back with your knees bent (as shown in the picture), or while standing with your feet apart and your knees bent.



6. With one hand holding the barrel, use the other hand to push the plunger into the barrel as far as it will go. Remove the applicator from the vagina.



7. Clean the applicator after each use. Pull the two pieces apart. Wash them with soap and water.

8. Lie down as soon as possible to keep the cream from leaking out.
 9. Repeat steps 2 through 8 before bedtime for the next 2 days.

To protect your clothing from leakage or discharge, you may want to use deodorant-lined pads or panty shields while you are using Trivagrole 3 Vaginal Cream. Do not use tampons, douches, or spermicides until after you have completed the treatment and your symptoms are gone.

Mfd. by:
 Taro Pharmaceuticals Inc.
 Bloomfield, Ontario
 Canada L6T 1C3
www.taropharma.com

FPD

10) - INSERT - BACK VIEW 4 X 12 FOLDED TO 1 X 1/2

COLORS...
 PMS 661 Date 3/17/00

page 6