





**ZYRTEC-D ALLERGY + SINUS**

Original Prescription Strength

**ZYRTEC-D**  
NDC 000000 000 000 00

Cetirizine HCl 5 mg / antihistamine  
Pseudoephedrine HCl 120 mg / nasal decongestant  
Extended Release Tablets

**ALLERGY + SINUS**

**NASAL CONGESTION + SINUS PRESSURE**

  
12 hours  
relief of

  
Sneezing

  
Runny Nose

  
Itchy, Watery Eyes

  
Itchy Throat or Nose

  
Actual Size

**24 Extended Release Tablets**





**Drug Facts** (continued)

**Indications** This product is indicated for the relief of the signs and symptoms of seasonal allergic rhinitis and allergic conjunctivitis.

**Warnings**

- Do not use if you are taking MAO inhibitors or have taken MAO inhibitors within the last 14 days.
- Do not use if you are taking other sympathomimetic amines such as pseudoephedrine or phenylephrine.
- Do not use if you are taking other antihistamines.
- Do not use if you are taking other decongestants.
- Do not use if you are taking other sedatives or tranquilizers.
- Do not use if you are taking other drugs that may cause drowsiness.
- Do not use if you are taking other drugs that may cause dizziness.
- Do not use if you are taking other drugs that may cause dry mouth.
- Do not use if you are taking other drugs that may cause blurred vision.
- Do not use if you are taking other drugs that may cause difficulty swallowing.
- Do not use if you are taking other drugs that may cause constipation.
- Do not use if you are taking other drugs that may cause urinary retention.
- Do not use if you are taking other drugs that may cause increased heart rate.
- Do not use if you are taking other drugs that may cause increased blood pressure.
- Do not use if you are taking other drugs that may cause increased risk of bleeding.
- Do not use if you are taking other drugs that may cause increased risk of infection.
- Do not use if you are taking other drugs that may cause increased risk of liver damage.
- Do not use if you are taking other drugs that may cause increased risk of kidney damage.
- Do not use if you are taking other drugs that may cause increased risk of pancreas damage.
- Do not use if you are taking other drugs that may cause increased risk of gallbladder disease.
- Do not use if you are taking other drugs that may cause increased risk of stomach ulcers.
- Do not use if you are taking other drugs that may cause increased risk of heart disease.
- Do not use if you are taking other drugs that may cause increased risk of stroke.
- Do not use if you are taking other drugs that may cause increased risk of death.

**Directions**

Take 1 tablet twice daily with water. Do not take more than 2 tablets in 24 hours.

**Other Information**

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-322-1222)

Keep this and all other medicines out of the reach of children. See important information about prescription medicines on the back of this package.

© 2000 Schering-Plough HealthCare Products, Inc.

www.zyrtec.com

**Drug Facts** (continued)

**Warnings**

- Do not use if you are taking MAO inhibitors or have taken MAO inhibitors within the last 14 days.
- Do not use if you are taking other sympathomimetic amines such as pseudoephedrine or phenylephrine.
- Do not use if you are taking other antihistamines.
- Do not use if you are taking other decongestants.
- Do not use if you are taking other sedatives or tranquilizers.
- Do not use if you are taking other drugs that may cause drowsiness.
- Do not use if you are taking other drugs that may cause dizziness.
- Do not use if you are taking other drugs that may cause dry mouth.
- Do not use if you are taking other drugs that may cause blurred vision.
- Do not use if you are taking other drugs that may cause difficulty swallowing.
- Do not use if you are taking other drugs that may cause constipation.
- Do not use if you are taking other drugs that may cause urinary retention.
- Do not use if you are taking other drugs that may cause increased heart rate.
- Do not use if you are taking other drugs that may cause increased blood pressure.
- Do not use if you are taking other drugs that may cause increased risk of bleeding.
- Do not use if you are taking other drugs that may cause increased risk of infection.
- Do not use if you are taking other drugs that may cause increased risk of liver damage.
- Do not use if you are taking other drugs that may cause increased risk of kidney damage.
- Do not use if you are taking other drugs that may cause increased risk of pancreas damage.
- Do not use if you are taking other drugs that may cause increased risk of gallbladder disease.
- Do not use if you are taking other drugs that may cause increased risk of stomach ulcers.
- Do not use if you are taking other drugs that may cause increased risk of heart disease.
- Do not use if you are taking other drugs that may cause increased risk of stroke.
- Do not use if you are taking other drugs that may cause increased risk of death.

**Directions**

Take 1 tablet twice daily with water. Do not take more than 2 tablets in 24 hours.

**Other Information**

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-322-1222)

Keep this and all other medicines out of the reach of children. See important information about prescription medicines on the back of this package.

© 2000 Schering-Plough HealthCare Products, Inc.

www.zyrtec.com

**Drug Facts** (continued)

**Warnings**

- Do not use if you are taking MAO inhibitors or have taken MAO inhibitors within the last 14 days.
- Do not use if you are taking other sympathomimetic amines such as pseudoephedrine or phenylephrine.
- Do not use if you are taking other antihistamines.
- Do not use if you are taking other decongestants.
- Do not use if you are taking other sedatives or tranquilizers.
- Do not use if you are taking other drugs that may cause drowsiness.
- Do not use if you are taking other drugs that may cause dizziness.
- Do not use if you are taking other drugs that may cause dry mouth.
- Do not use if you are taking other drugs that may cause blurred vision.
- Do not use if you are taking other drugs that may cause difficulty swallowing.
- Do not use if you are taking other drugs that may cause constipation.
- Do not use if you are taking other drugs that may cause urinary retention.
- Do not use if you are taking other drugs that may cause increased heart rate.
- Do not use if you are taking other drugs that may cause increased blood pressure.
- Do not use if you are taking other drugs that may cause increased risk of bleeding.
- Do not use if you are taking other drugs that may cause increased risk of infection.
- Do not use if you are taking other drugs that may cause increased risk of liver damage.
- Do not use if you are taking other drugs that may cause increased risk of kidney damage.
- Do not use if you are taking other drugs that may cause increased risk of pancreas damage.
- Do not use if you are taking other drugs that may cause increased risk of gallbladder disease.
- Do not use if you are taking other drugs that may cause increased risk of stomach ulcers.
- Do not use if you are taking other drugs that may cause increased risk of heart disease.
- Do not use if you are taking other drugs that may cause increased risk of stroke.
- Do not use if you are taking other drugs that may cause increased risk of death.

**Directions**

Take 1 tablet twice daily with water. Do not take more than 2 tablets in 24 hours.

**Other Information**

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-322-1222)

Keep this and all other medicines out of the reach of children. See important information about prescription medicines on the back of this package.

© 2000 Schering-Plough HealthCare Products, Inc.

www.zyrtec.com

**Net Content Statement**

PDP Area: (width x length) **13.5 in<sup>2</sup>**  
Net Content Size: **0.125"**

NET CONTENT	AREA OF PDP
1/8 IN.	5 SQ. IN. OR LESS
3/16 IN.	MORE THAN 5 SQ. IN. BUT NOT MORE THAN 25 SQ. IN.
1/4 IN.	MORE THAN 25 SQ. IN. BUT NOT MORE THAN 100 SQ. IN.
3/8 IN.	MORE THAN 100 SQ. IN. BUT NOT MORE THAN 400 SQ. IN.
1/2 IN.	OVER 400 SQ. IN.

**Labeling Format Information:** Font: Helvetica LT Pro-Narrow, Bold Cond, Bold Cond Obi

Drug Facts: 6.00 pt | Body Text: 6.00 pt | Bullets: 6.00 pt | Horizontal Scale: 80%

Headers: 6.00 pt | Leadings: 6.00 pt | Barlines: 2.00 pt

Subheaders: 6.00 pt | Drug Facts (continued): 8.00 pt | Halflines: 0.75 pt | Average Tracking: 0

**ZYRTEC-D**

Cetirizine HCl  
Pseudoephedrine HCl  
Cetirizine HCl



-----  
**This is a representation of an electronic record that was signed electronically. Following this are manifestations of any and all electronic signatures for this electronic record.**  
-----

/s/  
-----

MARTHA K LENHART  
09/12/2024 04:34:58 PM