

28 Count Outer Carton



100% SIZE
12 digit UPC
(non-suppressed)
For Position Only
0 37000 35906 7

Prilosec
OTC

Safety Feature—
Do not use
if tablet blister
unit is open
or torn.

NDC 37000-455-03

Easy Push-Through Package

Treats Frequent
Heartburn!

Prilosec
omeprazole delayed-release tablets
20 mg / acid reducer **OTC**

THIS PACKAGE FOR HOUSEHOLDS WITHOUT YOUNG CHILDREN

28 TABLETS
Two 14-day courses of treatment

Dist. by Preter & Gamble, Cincinnati, OH 45222
Product of Sweden 95227133

Prilosec
OTC

Prilosec
OTC

Questions? 1-800-289-9181

Inactive Ingredients hydroxypropyl cellulose, hypromellose, iron oxide, magnesium stearate, methacrylic acid copolymer, microcrystalline cellulose, paraffin, polyethylene glycol 6000, polysorbate 80, polyvinylpyrrolidone, sodium stearoyl fumarate, starch, sucrose, talc, titanium dioxide, triethyl citrate

■ protect product from moisture
■ keep product out of high heat and humidity
■ store at 20-25°C (68-77°F)
■ keep the carton and package insert. They contain important information.

Drug Facts (continued)

Warnings
Do not use if you have:
■ trouble or pain swallowing food
■ vomiting or black stools
■ bloody or black stools
These may be signs of a serious condition. See your doctor.

Ask a doctor before use if you have
■ had heartburn over 3 months. This may be a sign of a more serious condition.
■ heartburn with **lightheadedness, sweating or dizziness**
■ chest pain or shoulder pain with shortness of breath, sweating, pain spreading to arms, neck or shoulders, or lightheadedness

Ask a doctor or pharmacist before use if you are taking
■ warfarin (blood-thinning medicine)
■ prescription antifungal or anti-yeast medicines
■ diazepam (anxiety medicine), ■ digoxin (heart medicine)

Use
■ treats frequent heartburn (occurs 2 or more days a week)
■ not intended for immediate relief of heartburn. This drug may take 1 to 4 days for full effect.

Active Ingredient (in each tablet)
Omeprazole magnesium delayed-release tablet, 20.5 mg (equivalent to 20 mg omeprazole).....Acid reducer

Purpose
■ you need to take this product for more than 14 days
■ your heartburn continues or worsens
■ you need to take more than 1 course of treatment every 4 months

Stop use and ask a doctor if
■ pregnant or breast-feeding, ask a health professional before use.
■ you need to take more than 1 course of treatment every 4 months

Directions
■ adults 18 years of age and older
■ this product is to be used once a day (every 24 hours), every day for 14 days
■ it may take 1 to 4 days for full effect, although some people get complete relief of symptoms within 24 hours

14-Day Course of Treatment
■ swallow 1 tablet with a glass of water before eating in the morning
■ take every day for 14 days
■ do not take more than 1 tablet a day

Repeated 14-Day Courses (if needed)
■ do not chew or crush the tablets
■ do not crush tablets in food
■ do not use for more than 14 days unless directed by your doctor

Other information
■ read the directions, warnings and package insert before use

■ children under 18 years of age: ask a doctor
■ every 4 months unless directed by a doctor
■ you may repeat a 14-day course every 4 months
■ do not take for more than 14 days or more often than

Keep out of reach of children. In case of overdose, get medical help or contact a poison Control Center right away.

Drug Facts (continued)

Prilosec OTC
FPO 128 code

Prilosec OTC
FPO 128 code

14 Count Inner Carton

Other information

- read the directions, warnings and package insert before use
- children under 18 years of age: ask a doctor
- do not take for more than 14 days or more often than every 4 months unless directed by a doctor
- you may repeat a 14-day course every 4 months

Repeated 14-Day Courses (if needed)

- do not use for more than 14 days unless directed by your doctor
- do not crush tablets in food
- do not chew or crush the tablets
- do not take more than 1 tablet a day
- take every day for 14 days
- swallow 1 tablet with a glass of water before eating in the morning
- complete relief of symptoms within 24 hours
- it may take 1 to 4 days for full effect, although some people get 14 days
- this product is to be used once a day (every 24 hours), every day for adults 18 years of age and older

Directions

medical help or contact a Poison Control Center right away.

Keep out of reach of children. In case of overdose, get before use.

If pregnant or breast-feeding, ask a health professional

- you need to take more than 1 course of treatment every 4 months
- you need to take this product for more than 14 days
- your heartburn continues or worsens

Stop use and ask a doctor if

Drug Facts (continued)

Warnings

Do not use if you have

- trouble or pain swallowing food
- vomiting with blood
- bloody or black stools

These may be signs of a serious condition. See your doctor.

Ask a doctor before use if you have

- had heartburn over 3 months. This may be a sign of a more serious condition.
- heartburn with **lightheadedness, sweating or dizziness**
- chest pain or shoulder pain with shortness of breath, sweating, pain spreading to arms, neck or shoulders, or lightheadedness
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain

Ask a doctor or pharmacist before use if you are taking

- wafarin (blood-thinning medicine)
- prescription antifungal or anti-yeast medicines
- diazepam (anxiety medicine)
- digoxin (heart medicine)

Active ingredient (in each tablet) Purpose

Omeprazole magnesium delayed-release tablet 20.6 mg (equivalent to 20 mg omeprazole).....Acid reducer

Use

- treats frequent heartburn (occurs 2 or more days a week)
- not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Warnings

Allergy alert: Do not use if you are allergic to omeprazole

FPO 128 code

FPO 128 code

NOT FOR RESALE **Prilosec OTC** Safety Feature—Do not use if tablet blister unit is open or torn.

Drug Facts (continued)

- keep the carton and package insert. They contain important information.
- store at 20-25°C (68-77°F)
- keep product out of high heat and humidity
- protect product from moisture

Easy Push-Through Package **Treats Frequent Heartburn!**

Prilosec

omeprazole delayed-release tablets **OTC**[®]
20 mg / acid reducer

THIS PACKAGE FOR HOUSEHOLDS WITHOUT YOUNG CHILDREN

14 TABLETS
One 14-day course of treatment

Drug Facts (continued)

Inactive ingredients glyceryl monostearate, hydroxypropyl cellulose, hypromellose, iron oxide, magnesium stearate, methacrylic acid copolymer, microcrystalline cellulose, paraffin, polyethylene glycol 6000, polysorbate 80, polyvinylpyrrolidone, sodium stearyl fumarate, starch, sucrose, talc, titanium dioxide, triethyl citrate

Dist. by
Procter
& Gamble,
Cincinnati,
OH 45202

Drug Facts (continued)
Questions? 1-800-289-9181

Product of Sweden 98227631

NDC 37000-455-02

Blister Card (Non-CR)

Do not
chew or
crush tablets

Prilosec OTC[®]

Do not
crush tablets
in food

1 omeprazole delayed-release tablet
20 mg / acid reducer

Push tablet through foil.

Dist. by Procter & Gamble, Cincinnati, OH 45202

PACKAGE NOT CHILD-RESISTANT

95437426

Prilosec OTC[®]

1 omeprazole delayed-release tablet
20 mg / acid reducer

Push tablet through foil.

Dist. by Procter & Gamble,
Cincinnati, OH 45202

Prilosec OTC[®]

1 omeprazole delayed-release tablet
20 mg / acid reducer

Push tablet through foil.

Dist. by Procter & Gamble,
Cincinnati, OH 45202

Prilosec OTC[®]

1 omeprazole delayed-release tablet
20 mg / acid reducer

Push tablet through foil.

Dist. by Procter & Gamble,
Cincinnati, OH 45202

Prilosec OTC[®]

1 omeprazole delayed-release tablet
20 mg / acid reducer

Push tablet through foil.

Dist. by Procter & Gamble,
Cincinnati, OH 45202

Prilosec OTC[®]

1 omeprazole delayed-release tablet
20 mg / acid reducer

Push tablet through foil.

Dist. by Procter & Gamble,
Cincinnati, OH 45202

Prilosec OTC[®]

1 omeprazole delayed-release tablet
20 mg / acid reducer

Push tablet through foil.

Dist. by Procter & Gamble,
Cincinnati, OH 45202